

Resonant electron-molecular cation collisions in the edge plasmas of fusion devices: new state-to-state cross sections and rate coefficients

J. Zs. Mezei^{1,2}, A. Abdoulanziz², Y. Moulane^{2,3}, S. Niyonzima⁴, M. D. Epée Epée⁵, F. Iacob⁶, D. A. Little⁷, N. Pop⁸, K. Chakrabarti⁹, O. Motapon⁵, J. Tennyson⁷, and I. F. Schneider^{2,10}

¹(Presenting author underlined) *Institute of Nuclear Research, Hungarian Academy of Sciences, Debrecen, Hungary*

²*Lab. Ondes et Milieux Complexes, CNRS, Université du Havre, France*

³*High Energy Physics and Astrophysics Lab., Cadi Ayyad University, Marrakech, Morocco*

⁴*Dépt. de Physique, Université du Burundi, Bujumbura, Burundi*

⁵*Dept. of Physics, Faculty of Sciences, University of Douala, Cameroon*

⁶*Physics Faculty, West University of Timișoara, Romania*


⁷*Dept. of Physics and Astronomy, University College London, United Kingdom*

⁸*Dept. of Physical Foundation of Engineering, Politehnica University Timișoara, Romania*

⁹*Dept. of Mathematics, Scottish Church College, Calcutta, India*

¹⁰*Lab. Aimé Cotton, CNRS, ENS Cachan and Université Paris-Sud, Orsay, France*

Dissociative recombination, elastic scattering, (ro-)vibrational excitation, vibrational de-excitation and dissociative excitation [1]:



are dominant elementary processes in numerous cold ionized gases. Here N^+/v^+ stand for the rotational/vibrational quantum numbers of the cation, AB^* for a bound excited (mostly Rydberg) state of the neutral, and AB^{**} for a dissociative (mostly doubly- or multiply-excited) state of the neutral.

The quantum interference between the direct mechanism - capture into the doubly-excited states AB^{**} - and the indirect one - temporary capture into a Rydberg state AB^* - induces resonances in the cross section.

The Multichannel Quantum Defect Theory (MQDT) [2, 3, 4, 5, 6] is the most suitable approach for these processes, efficiently handling channels - open for the direct process and closed for the indirect one - and the corresponding channel mixing via electronic and vibronic interactions. We will provide new cross sections and rate coefficients for H_2^+ , HD^+ [2], BeH^+ [3, 4], BeD^+ [5], CH^+ , N_2^+ [6].

- [1] I. F. Schneider, O. Dulieu, J. Robert, editors, *Proceedings of DR2013: The 9th International Conference on Dissociative Recombination: Theory, Experiment and Applications*, Paris, July 7-12, 2013, EPJ Web of Conferences **84** (2015).
- [2] K. Chakrabarti *et al*, Phys. Rev. A **87** 022702 (2013).
- [3] S. Niyonzima *et al*, At. Data Nucl. Data Tables **115-116** 287 (2017).
- [4] V. Laporta *et al*, Plasma Phys. Control. Fusion **59** 045008 (2017).
- [5] N. Pop *et al*, AIP Conf. Proc. **1916** 020013 (2017).
- [6] D. A. Little *et al*, Phys. Rev. A **90** 052705 (2014).