

# $G_2$ -manifolds

Sebastian Goette

Mathematisches Institut der Universität Freiburg

23. 5. 2018

# Outline

- ▶ Holonomy  
Intro and general concepts
- ▶  $G_2$ -geometry  
Properties, constructions, questions
- ▶ The  $\nu$ -invariant  
Definition, properties, first examples
- ▶ Extra twisted connected sums  
Computation of the  $\nu$ -invariant, consequences

# Holonomy

Let  $(M, g)$  be a Riemannian manifold,  $\dim M = m$ , with Levi-Civita connection  $\nabla$ .

Fix  $p \in M$ . For each loop  $\gamma: [a, b] \rightarrow M$  at  $p$ , **parallel translation** with respect to  $\nabla$  defines an automorphism of the tangent space at  $p$ ,

$$\Pi_\gamma: T_p M \rightarrow T_p M$$

# Holonomy

Let  $(M, g)$  be a Riemannian manifold,  $\dim M = m$ , with Levi-Civita connection  $\nabla$ .

Fix  $p \in M$ . For each loop  $\gamma: [a, b] \rightarrow M$  at  $p$ , **parallel translation** with respect to  $\nabla$  defines an automorphism of the tangent space at  $p$ ,

$$\Pi_\gamma: T_p M \rightarrow T_p M$$

The **holonomy group** is the group of all these parallel translations

$$\text{Hol}_p(M, g) = (\{ \Pi_\gamma \mid \gamma \text{ loop at } p \}, \circ, \text{id})$$

# Holonomy

Let  $(M, g)$  be a Riemannian manifold,  $\dim M = m$ , with Levi-Civita connection  $\nabla$ .

Fix  $p \in M$ . For each loop  $\gamma: [a, b] \rightarrow M$  at  $p$ , **parallel translation** with respect to  $\nabla$  defines an automorphism of the tangent space at  $p$ ,


$$\Pi_\gamma: T_p M \rightarrow T_p M$$

The **holonomy group** is the group of all these parallel translations

$$\begin{aligned} \text{Hol}_p(M, g) &= (\{ \Pi_\gamma \mid \gamma \text{ loop at } p \}, \circ, \text{id}) \\ &\subset \text{Aut}(T_p M, g_p) \cong O(m) \end{aligned}$$


# Holonomy—an example

Consider parallel translation  
along a spherical triangle




# Holonomy—an example

Consider parallel translation  
along a spherical triangle



# Holonomy—an example


Consider parallel translation  
along a spherical triangle





# Holonomy—an example

Consider parallel translation  
along a spherical triangle




# Holonomy—an example

Consider parallel translation  
along a spherical triangle

A vector is rotated by an  
angle equal to the spherical  
area of the triangle

(Gauß-Bonnet theorem)




# Holonomy—an example

Consider parallel translation  
along a spherical triangle

A vector is rotated by an  
angle equal to the spherical  
area of the triangle

(Gauß-Bonnet theorem)

$$\text{Hol}(S^2, g^{\text{rd}}) \cong SO(2)$$



# Holonomy—Kähler manifolds

Let  $(M, g, J)$  be a **Kähler manifold**,  $\dim M = 2n$ , so  $J$  is a parallel complex structure on the tangent bundle  $TM$

In particular,  $(M, J)$  is a complex manifold

Examples: complex projective space,  
smooth complex projective varieties

Parallel translation commutes with  $J$ , hence

$$\mathrm{Hol}_p(M, g) \subset \mathrm{Aut}(T_p M, g_p, J_p) \cong U(n) \subset SO(2n)$$

# Holonomy—Kähler manifolds

Let  $(M, g, J)$  be a **Kähler manifold**,  $\dim M = 2n$ , so  $J$  is a parallel complex structure on the tangent bundle  $TM$ .  
In particular,  $(M, J)$  is a complex manifold.

Examples: complex projective space,  
smooth complex projective varieties

Parallel translation commutes with  $J$ , hence

$$\mathrm{Hol}_p(M, g) \subset \mathrm{Aut}(T_p M, g_p, J_p) \cong U(n) \subset SO(2n)$$

**Hodge decomposition** of de Rham cohomology on  $(M, g, J)$

$$\Omega^\ell(M; \mathbb{C}) = \bigoplus_{p+q=\ell} \Omega^{p,q}(M)$$

$$H_{\mathrm{dR}}^\ell(M; \mathbb{C}) = \bigoplus_{p+q=\ell} H^{p,q}(M)$$

# Holonomy—Calabi-Yau manifolds

Let  $(M, g, J)$  be a Kähler manifold,  $\dim_{\mathbb{C}} M = n$

Then  $(M, g, J)$  is called **Calabi-Yau manifold**

if  $M$  has a parallel complex volume form  $\Omega \in \Omega^{n,0}(M)$

$$\mathrm{Hol}_p(M, g) \subset \mathrm{Aut}(T_p M, g_p, J_p, \Omega_p) \cong SU(n) \subset U(n)$$

# Holonomy—Calabi-Yau manifolds

Let  $(M, g, J)$  be a Kähler manifold,  $\dim_{\mathbb{C}} M = n$

Then  $(M, g, J)$  is called **Calabi-Yau manifold**

if  $M$  has a parallel complex volume form  $\Omega \in \Omega^{n,0}(M)$

$$\mathrm{Hol}_p(M, g) \subset \mathrm{Aut}(T_p M, g_p, J_p, \Omega_p) \cong SU(n) \subset U(n)$$

Calabi-Yau manifolds are **Ricci flat**,  $\mathrm{ric} = 0$

# Holonomy—Calabi-Yau manifolds

Let  $(M, g, J)$  be a Kähler manifold,  $\dim_{\mathbb{C}} M = n$

Then  $(M, g, J)$  is called **Calabi-Yau manifold**

if  $M$  has a parallel complex volume form  $\Omega \in \Omega^{n,0}(M)$

$$\mathrm{Hol}_p(M, g) \subset \mathrm{Aut}(T_p M, g_p, J_p, \Omega_p) \cong SU(n) \subset U(n)$$

Calabi-Yau manifolds are **Ricci flat**,  $\mathrm{ric} = 0$

## Theorem (Yau)

*A compact Kähler manifold admits a Calabi-Yau metric if and only if  $c_1(TM) = 0$ .*



# Holonomy—Calabi-Yau manifolds

Let  $(M, g, J)$  be a Kähler manifold,  $\dim_{\mathbb{C}} M = n$

Then  $(M, g, J)$  is called **Calabi-Yau manifold**

if  $M$  has a parallel complex volume form  $\Omega \in \Omega^{n,0}(M)$

$$\mathrm{Hol}_p(M, g) \subset \mathrm{Aut}(T_p M, g_p, J_p, \Omega_p) \cong SU(n) \subset U(n)$$

Calabi-Yau manifolds are **Ricci flat**,  $\mathrm{ric} = 0$

## Theorem (Yau)

*A compact Kähler manifold admits a Calabi-Yau metric if and only if  $c_1(TM) = 0$ .*

Methods in / Motivation for Calabi-Yau geometry:

**Geometric Analysis** and **Algebraic Geometry**

# Holonomy—Calabi-Yau manifolds

Let  $(M, g, J)$  be a Kähler manifold,  $\dim_{\mathbb{C}} M = n$

Then  $(M, g, J)$  is called **Calabi-Yau manifold**

if  $M$  has a parallel complex volume form  $\Omega \in \Omega^{n,0}(M)$

$$\mathrm{Hol}_p(M, g) \subset \mathrm{Aut}(T_p M, g_p, J_p, \Omega_p) \cong SU(n) \subset U(n)$$

Calabi-Yau manifolds are **Ricci flat**,  $\mathrm{ric} = 0$

## Theorem (Yau)

*A compact Kähler manifold admits a Calabi-Yau metric if and only if  $c_1(TM) = 0$ .*

Methods in / Motivation for Calabi-Yau geometry:

**Geometric Analysis** and **Algebraic Geometry**

**Mathematical Physics** (String Theory / Conformal Field Theory)

# Holonomy—Representation theory

Let  $P_{SO}$  denote the manifold of oriented orthonormal bases of  $TM$ . Then  $SO(n)$  acts on  $P_{SO}$  by change of bases, and

$$M \cong P_{SO}/SO(n) \quad \text{and} \quad TM \cong P_{SO} \times_{SO(n)} \mathbb{R}^n$$

# Holonomy—Representation theory

Let  $P_{SO}$  denote the manifold of oriented orthonormal bases of  $TM$ . Then  $SO(n)$  acts on  $P_{SO}$  by change of bases, and

$$M \cong P_{SO}/SO(n) \quad \text{and} \quad TM \cong P_{SO} \times_{SO(n)} \mathbb{R}^n$$

If  $(M, g)$  has holonomy  $G$ , there is a submanifold  $P_G \subset P_{SO}$  with

$$M \cong P_G/G \quad \text{and} \quad TM \cong P_G \times_G \mathbb{R}^n$$

# Holonomy—Representation theory

Let  $P_{SO}$  denote the manifold of oriented orthonormal bases of  $TM$ . Then  $SO(n)$  acts on  $P_{SO}$  by change of bases, and

$$M \cong P_{SO}/SO(n) \quad \text{and} \quad TM \cong P_{SO} \times_{SO(n)} \mathbb{R}^n$$

If  $(M, g)$  has holonomy  $G$ , there is a submanifold  $P_G \subset P_{SO}$  with

$$M \cong P_G/G \quad \text{and} \quad TM \cong P_G \times_G \mathbb{R}^n$$

Differential forms on  $M$  live in the bundle

$$\Lambda^\bullet T^*M \cong P_{SO} \times_{SO(n)} \Lambda^\bullet \mathbb{R}^n \cong P_G \times_G \Lambda^\bullet \mathbb{R}^n$$

If  $\Lambda^\bullet \mathbb{R}^n$  is a direct sum of  $G$ -invariant subspaces, then  $\Lambda^\bullet T^*M$  is a direct sum of parallel subbundles

# Holonomy—Representation theory

Let  $P_{SO}$  denote the manifold of oriented orthonormal bases of  $TM$ . Then  $SO(n)$  acts on  $P_{SO}$  by change of bases, and

$$M \cong P_{SO}/SO(n) \quad \text{and} \quad TM \cong P_{SO} \times_{SO(n)} \mathbb{R}^n$$

If  $(M, g)$  has holonomy  $G$ , there is a submanifold  $P_G \subset P_{SO}$  with

$$M \cong P_G/G \quad \text{and} \quad TM \cong P_G \times_G \mathbb{R}^n$$

Differential forms on  $M$  live in the bundle

$$\Lambda^\bullet T^*M \cong P_{SO} \times_{SO(n)} \Lambda^\bullet \mathbb{R}^n \cong P_G \times_G \Lambda^\bullet \mathbb{R}^n$$

If  $\Lambda^\bullet \mathbb{R}^n$  is a direct sum of  $G$ -invariant subspaces, then  $\Lambda^\bullet T^*M$  is a direct sum of parallel subbundles

**Example.** Hodge decomposition on Kähler manifolds

# Holonomy—Parallel spinors

If the holonomy group  $G$  is connected and simply connected, then  $(M, g)$  is a **spin manifold** with spinor bundle

$$SM = P_G \times_G S\mathbb{R}^n \quad \text{with} \quad S\mathbb{R}^n \cong \mathbb{R}^{2\left[\frac{n}{2}\right]}$$

# Holonomy—Parallel spinors

If the holonomy group  $G$  is connected and simply connected, then  $(M, g)$  is a **spin manifold** with spinor bundle

$$SM = P_G \times_G S\mathbb{R}^n \quad \text{with} \quad S\mathbb{R}^n \cong \mathbb{R}^{2\left[\frac{n}{2}\right]}$$

If the  $G$ -invariant subspace  $(S\mathbb{R}^n)^G \subset S\mathbb{R}^n$  has dimension  $N$  then  $(M, g)$  admits an  $N$ -dimensional space of **parallel spinors**



# Holonomy—Parallel spinors

If the holonomy group  $G$  is connected and simply connected, then  $(M, g)$  is a **spin manifold** with spinor bundle

$$SM = P_G \times_G S\mathbb{R}^n \quad \text{with} \quad S\mathbb{R}^n \cong \mathbb{R}^{2\left[\frac{n}{2}\right]}$$

If the  $G$ -invariant subspace  $(S\mathbb{R}^n)^G \subset S\mathbb{R}^n$  has dimension  $N$  then  $(M, g)$  admits an  $N$ -dimensional space of **parallel spinors**

If  $(M, g)$  admit a parallel spinor  $\sigma \neq 0$ , so  $N \geq 1$ , then **ric = 0**

# Holonomy—Parallel spinors

If the holonomy group  $G$  is connected and simply connected, then  $(M, g)$  is a **spin manifold** with spinor bundle

$$SM = P_G \times_G S\mathbb{R}^n \quad \text{with} \quad S\mathbb{R}^n \cong \mathbb{R}^{2\left[\frac{n}{2}\right]}$$

If the  $G$ -invariant subspace  $(S\mathbb{R}^n)^G \subset S\mathbb{R}^n$  has dimension  $N$  then  $(M, g)$  admits an  $N$ -dimensional space of **parallel spinors**

If  $(M, g)$  admit a parallel spinor  $\sigma \neq 0$ , so  $N \geq 1$ , then **ric = 0**

**Example.** Let  $(M, g)$  be a Calabi-Yau manifold. Then

$$SM \cong \Lambda^{0,\bullet} T^*M$$

The forms  $1 \in \Omega^{0,0}(M)$  and  $\bar{\Omega} \in \Omega^{0,n}(M)$  are parallel spinors and  $\text{ric} = 0$

# Holonomy—Berger's list

## Theorem (Berger)

*Let  $(M, g)$  be a complete, simply connected Riemannian manifold and neither a product nor a symmetric space*

# Holonomy—Berger's list

## Theorem (Berger)

*Let  $(M, g)$  be a complete, simply connected Riemannian manifold and neither a product nor a symmetric space*

*Then these are the only possibilities for its holonomy*

$\text{Hol}(M, g)$	dim	ric	Structure	$N$	Name
$SO(n)$	$n$			0	general case
$U(k)$	$2k$		$J$	0	Kähler
$SU(k)$	$2k$	0	$J, \Omega$	2	Calabi-Yau
$Sp(\ell) \cdot Sp(1)$	$4\ell$	const	$\langle I, J, K \rangle$	0	Quat. Kähler
$Sp(\ell)$	$4\ell$	0	$I, J, K, \Omega$	$\ell + 1$	hyper Kähler
$G_2$	7	0	$\varphi \in \Omega^3$	1	exceptional
$\text{Spin}(7)$	8	0	$\psi \in \Omega^4$	1	exceptional

# Holonomy—Berger's list

## Theorem (Berger)

*Let  $(M, g)$  be a complete, simply connected Riemannian manifold and neither a product nor a symmetric space*

*Then these are the only possibilities for its holonomy*

$\text{Hol}(M, g)$	dim	ric	Structure	$N$	Name
$SO(n)$	$n$			0	general case
$U(k)$	$2k$		$J$	0	Kähler
$SU(k)$	$2k$	0	$J, \Omega$	2	Calabi-Yau
$Sp(\ell) \cdot Sp(1)$	$4\ell$	const	$\langle I, J, K \rangle$	0	Quat. Kähler
$Sp(\ell)$	$4\ell$	0	$I, J, K, \Omega$	$\ell + 1$	hyper Kähler
$G_2$	7	0	$\varphi \in \Omega^3$	1	exceptional
$\text{Spin}(7)$	8	0	$\psi \in \Omega^4$	1	exceptional

# $G_2$ -Geometry—Some motivation

Why consider  $G_2$ -manifolds?

# $G_2$ -Geometry—Some motivation

Why consider  $G_2$ -manifolds?

## Mathematical motivation

- ▶ Only special holonomy group for odd dimensions
- ▶ Only  $G_2$  and  $\text{Spin}(7)$  holonomy have no direct relation to algebraic geometry

Hence, new methods are needed

# $G_2$ -Geometry—Some motivation

Why consider  $G_2$ -manifolds?

## Mathematical motivation

- ▶ Only special holonomy group for odd dimensions
- ▶ Only  $G_2$  and  $\text{Spin}(7)$  holonomy have no direct relation to algebraic geometry

Hence, new methods are needed

## Physical motivation

- ▶ In string theory, spacetime takes the form  $\mathbb{R}^{3,1} \times V$ , where  $V$  is a Calabi-Yau manifold
- ▶ In **M-theory**, spacetime takes the form  $\mathbb{R}^{3,1} \times M$ , where  $M$  is a  $G_2$ -manifold
- ▶ Possible relations to other physical theories

Hence, many fruitful interactions possible



# $G_2$ -Geometry—The group $G_2$

Recall the Cayley-Dickson construction

$$\mathbb{R} \subset \mathbb{C}$$

# $G_2$ -Geometry—The group $G_2$

Recall the **Cayley-Dickson** construction

$$\mathbb{R} \subset \mathbb{C} \subset \mathbb{H}$$

The quaternions  $\mathbb{H}$  are not commutative

# $G_2$ -Geometry—The group $G_2$

Recall the **Cayley-Dickson** construction

$$\mathbb{R} \subset \mathbb{C} \subset \mathbb{H} \subset \mathbb{O}$$

The quaternions  $\mathbb{H}$  are not commutative

The octonions  $\mathbb{O}$  are neither commutative nor associative

# $G_2$ -Geometry—The group $G_2$

Recall the **Cayley-Dickson** construction

$$\mathbb{R} \subset \mathbb{C} \subset \mathbb{H} \subset \mathbb{O}$$

The quaternions  $\mathbb{H}$  are not commutative

The octonions  $\mathbb{O}$  are neither commutative nor associative

The octonions split as  $\mathbb{O} \cong \mathbb{R} \oplus \text{Im } \mathbb{O}$  with  $\text{Im } \mathbb{O} \cong \mathbb{R}^7$

The octonion multiplication induces a **product** and a **norm**

$$u \times v = \text{Im}(u \cdot v) \quad \text{and} \quad \|u\| = \sqrt{-u^2} \quad \text{for } u, v \in \text{Im } \mathbb{O}$$

# $G_2$ -Geometry—The group $G_2$

Recall the **Cayley-Dickson** construction

$$\mathbb{R} \subset \mathbb{C} \subset \mathbb{H} \subset \mathbb{O}$$

The quaternions  $\mathbb{H}$  are not commutative

The octonions  $\mathbb{O}$  are neither commutative nor associative

The octonions split as  $\mathbb{O} \cong \mathbb{R} \oplus \text{Im } \mathbb{O}$  with  $\text{Im } \mathbb{O} \cong \mathbb{R}^7$

The octonion multiplication induces a **product** and a **norm**

$$u \times v = \text{Im}(u \cdot v) \quad \text{and} \quad \|u\| = \sqrt{-u^2} \quad \text{for } u, v \in \text{Im } \mathbb{O}$$

The group  $G_2$  is defined as

$$G_2 = \text{Aut}(\mathbb{O}) = \text{Aut}(\text{Im } \mathbb{O}, \times, \|\cdot\|)$$

# $G_2$ -geometry—The 3-form $\varphi$

Assume  $\text{Hol}(M, g) \subset G_2$ . Then “ $\times$ ” induces a parallel product on  $TM$  and a parallel 3-form  $\varphi \in \Omega^3(M)$  with

$$\varphi(u, v, w) = \langle u \times v, w \rangle$$

Note that

$$\varphi \wedge \iota_u \varphi \wedge \iota_v \varphi = -6 \langle u, v \rangle d\text{vol}_g \quad (*)$$

# $G_2$ -geometry—The 3-form $\varphi$

Assume  $\text{Hol}(M, g) \subset G_2$ . Then “ $\times$ ” induces a parallel product on  $TM$  and a parallel 3-form  $\varphi \in \Omega^3(M)$  with

$$\varphi(u, v, w) = \langle u \times v, w \rangle$$

Note that

$$\varphi \wedge \iota_u \varphi \wedge \iota_v \varphi = -6 \langle u, v \rangle d\text{vol}_g \quad (*)$$

Conversely, call an arbitrary  $\varphi \in \Omega^3(M)$  **positive** if there exists a Riemannian metric  $g_\varphi$  on  $M$  such that  $(*)$  holds

Being positive is a point-wise, open condition on  $\Lambda^3 T^*M$

Each positive 3-form  $\varphi$  defines a  **$G_2$ -structure** on  $M$

## $G_2$ -geometry—The 3-form $\varphi$

Assume  $\text{Hol}(M, g) \subset G_2$ . Then “ $\times$ ” induces a parallel product on  $TM$  and a parallel 3-form  $\varphi \in \Omega^3(M)$  with

$$\varphi(u, v, w) = \langle u \times v, w \rangle$$

Note that

$$\varphi \wedge \iota_u \varphi \wedge \iota_v \varphi = -6 \langle u, v \rangle d\text{vol}_g \quad (*)$$

Conversely, call an arbitrary  $\varphi \in \Omega^3(M)$  **positive** if there exists a Riemannian metric  $g_\varphi$  on  $M$  such that  $(*)$  holds

Being positive is a point-wise, open condition on  $\Lambda^3 T^*M$

Each positive 3-form  $\varphi$  defines a  **$G_2$ -structure** on  $M$

If  $\varphi$  above is closed and coclosed with respect to  $g_\varphi$ , then  $\varphi$  is parallel, also called **torsion free**, and  $\text{Hol}(M, g) \subset G_2$   
Write  $(M, g)$  for  $(M, \varphi)$  if  $(M, g)$  has  **$G_2$ -holonomy**



# $G_2$ -geometry—Representation theory

If  $M$  has a  $G_2$ -structure, we have “Hodge decompositions”

$$\Omega^0(M) \cong \Omega^7(M) \cong \Omega_1(M)$$

$$\Omega^1(M) \cong \Omega^6(M) \cong \Omega_7(M)$$

$$\Omega^2(M) \cong \Omega^5(M) \cong \Omega_7(M) \oplus \Omega_{14}(M)$$

$$\Omega^3(M) \cong \Omega^4(M) \cong \Omega_1(M) \oplus \Omega_7(M) \oplus \Omega_{27}(M)$$

# $G_2$ -geometry—Representation theory

If  $M$  has a  $G_2$ -structure, we have “Hodge decompositions”

$$\Omega^0(M) \cong \Omega^7(M) \cong \Omega_1(M)$$

$$\Omega^1(M) \cong \Omega^6(M) \cong \Omega_7(M)$$

$$\Omega^2(M) \cong \Omega^5(M) \cong \Omega_7(M) \oplus \Omega_{14}(M)$$

$$\Omega^3(M) \cong \Omega^4(M) \cong \Omega_1(M) \oplus \Omega_7(M) \oplus \Omega_{27}(M)$$

If  $\text{Hol}(M, g) \subset G_2$  we have similar decompositions of  $H_{\text{dR}}^\bullet(M)$

# $G_2$ -geometry—Representation theory

If  $M$  has a  $G_2$ -structure, we have “Hodge decompositions”

$$\Omega^0(M) \cong \Omega^7(M) \cong \Omega_1(M)$$

$$\Omega^1(M) \cong \Omega^6(M) \cong \Omega_7(M)$$

$$\Omega^2(M) \cong \Omega^5(M) \cong \Omega_7(M) \oplus \Omega_{14}(M)$$

$$\Omega^3(M) \cong \Omega^4(M) \cong \Omega_1(M) \oplus \Omega_7(M) \oplus \Omega_{27}(M)$$

If  $\text{Hol}(M, g) \subset G_2$  we have similar decompositions of  $H_{\text{dR}}^\bullet(M)$

If  $\text{Hol}(M, g) = G_2$  then  $H_7^k(M) = 0$  for  $1 \leq k \leq 6$  and

$$H^3(M) \cong \langle \varphi \rangle \oplus H_{27}^3(M) \quad \text{and} \quad H^4(M) \cong \langle *\varphi \rangle \oplus H_{27}^4(M)$$

# $G_2$ -geometry—Representation theory

If  $M$  has a  $G_2$ -structure, we have “Hodge decompositions”

$$\Omega^0(M) \cong \Omega^7(M) \cong \Omega_1(M)$$

$$\Omega^1(M) \cong \Omega^6(M) \cong \Omega_7(M)$$

$$\Omega^2(M) \cong \Omega^5(M) \cong \Omega_7(M) \oplus \Omega_{14}(M)$$

$$\Omega^3(M) \cong \Omega^4(M) \cong \Omega_1(M) \oplus \Omega_7(M) \oplus \Omega_{27}(M)$$

If  $\text{Hol}(M, g) \subset G_2$  we have similar decompositions of  $H_{\text{dR}}^\bullet(M)$

If  $\text{Hol}(M, g) = G_2$  then  $H_7^k(M) = 0$  for  $1 \leq k \leq 6$  and

$$H^3(M) \cong \langle \varphi \rangle \oplus H_{27}^3(M) \quad \text{and} \quad H^4(M) \cong \langle *\varphi \rangle \oplus H_{27}^4(M)$$

$G_2$ -manifolds are spin, and the spinor bundle decomposes

$$SM \cong TM \oplus \underline{\mathbb{R}}$$

There exists a distinguished parallel spinor  $\sigma \neq 0$ , and  $\text{ric} = 0$

# $G_2$ -geometry—Obstructions

Let  $p_1(TM) \in H^4(M; \mathbb{R})$  denote the first Pontryagin class

# $G_2$ -geometry—Obstructions

Let  $p_1(TM) \in H^4(M; \mathbb{R})$  denote the first Pontryagin class

Let  $\varphi \in \Omega^3(M)$  be positive and closed. There exists a bilinear form on  $H^2(M; \mathbb{R})$  given for closed forms  $\alpha, \beta \in \Omega^2(M)$  by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

# $G_2$ -geometry—Obstructions

Let  $p_1(TM) \in H^4(M; \mathbb{R})$  denote the first Pontryagin class

Let  $\varphi \in \Omega^3(M)$  be positive and closed. There exists a bilinear form on  $H^2(M; \mathbb{R})$  given for closed forms  $\alpha, \beta \in \Omega^2(M)$  by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

If  $M$  is compact and  $\text{Hol}(M, \varphi) \subset G_2$  then

- ▶  $M$  is **oriented** and **spin** and  **$\dim H^3(M; \mathbb{R}) \geq 1$**

# $G_2$ -geometry—Obstructions

Let  $p_1(TM) \in H^4(M; \mathbb{R})$  denote the first Pontryagin class

Let  $\varphi \in \Omega^3(M)$  be positive and closed. There exists a bilinear form on  $H^2(M; \mathbb{R})$  given for closed forms  $\alpha, \beta \in \Omega^2(M)$  by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

If  $M$  is compact and  $\text{Hol}(M, \varphi) \subset G_2$  then

- ▶  $M$  is **oriented** and **spin** and  $\dim H^3(M; \mathbb{R}) \geq 1$
- ▶  $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$  (Cheeger-Gromoll)



# $G_2$ -geometry—Obstructions

Let  $p_1(TM) \in H^4(M; \mathbb{R})$  denote the first Pontryagin class

Let  $\varphi \in \Omega^3(M)$  be positive and closed. There exists a bilinear form on  $H^2(M; \mathbb{R})$  given for closed forms  $\alpha, \beta \in \Omega^2(M)$  by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

If  $M$  is compact and  $\text{Hol}(M, \varphi) \subset G_2$  then

- ▶  $M$  is **oriented** and **spin** and  $\dim H^3(M; \mathbb{R}) \geq 1$
- ▶  $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$  (Cheeger-Gromoll)
- ▶  $\text{Hol}(M, g) = G_2 \iff B$  is negative definite

# $G_2$ -geometry—Obstructions

Let  $p_1(TM) \in H^4(M; \mathbb{R})$  denote the first Pontryagin class

Let  $\varphi \in \Omega^3(M)$  be positive and closed. There exists a bilinear form on  $H^2(M; \mathbb{R})$  given for closed forms  $\alpha, \beta \in \Omega^2(M)$  by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

If  $M$  is compact and  $\text{Hol}(M, \varphi) \subset G_2$  then

- ▶  $M$  is **oriented** and **spin** and  $\dim H^3(M; \mathbb{R}) \geq 1$
- ▶  $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$  (Cheeger-Gromoll)
- ▶  $\text{Hol}(M, g) = G_2 \iff B$  is negative definite
- ▶ If  $\text{Hol}(M, g) = G_2$  then  $(p_1(TM) \smile [\varphi])[M] < 0$

# $G_2$ -geometry—Obstructions

Let  $p_1(TM) \in H^4(M; \mathbb{R})$  denote the first Pontryagin class

Let  $\varphi \in \Omega^3(M)$  be positive and closed. There exists a bilinear form on  $H^2(M; \mathbb{R})$  given for closed forms  $\alpha, \beta \in \Omega^2(M)$  by

$$B([\alpha], [\beta]) = ([\alpha] \smile [\beta] \smile [\varphi])[M] = \int_M \alpha \wedge \beta \wedge \varphi$$

If  $M$  is compact and  $\text{Hol}(M, \varphi) \subset G_2$  then

- ▶  $M$  is **oriented** and **spin** and  $\dim H^3(M; \mathbb{R}) \geq 1$
- ▶  $\text{Hol}(M, g) = G_2 \iff \#\pi_1(M) < \infty$  (Cheeger-Gromoll)
- ▶  $\text{Hol}(M, g) = G_2 \iff B$  is negative definite
- ▶ If  $\text{Hol}(M, g) = G_2$  then  $(p_1(TM) \smile [\varphi])[M] < 0$

These are all **known** obstructions against holonomy  $G_2$

# $G_2$ -geometry—The moduli space

Let  $M$  be a compact oriented 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

# $G_2$ -geometry—The moduli space

Let  $M$  be a compact oriented 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

Let  $\mathcal{D} \subset \text{Diff}(M)$  be the connected component of  $\text{id}_M$ . Then

$$\mathcal{M} = \mathcal{X}/\mathcal{D}$$

is called the  $G_2$ -moduli space of  $M$

# $G_2$ -geometry—The moduli space

Let  $M$  be a compact oriented 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

Let  $\mathcal{D} \subset \text{Diff}(M)$  be the connected component of  $\text{id}_M$ . Then

$$\mathcal{M} = \mathcal{X}/\mathcal{D}$$

is called the  $G_2$ -moduli space of  $M$

## Theorem (Joyce)

*The  $G_2$ -moduli space is a manifold, and the map*

$$\mathcal{M} \longrightarrow H^3(M; \mathbb{R}) \quad \text{with} \quad [\varphi] \longmapsto [\varphi]$$

*is a **local** diffeomorphism*

# $G_2$ -geometry—The moduli space

Let  $M$  be a compact oriented 7-manifold and define

$$\mathcal{X} = \{ \varphi \in \Omega^3(M) \mid \varphi \text{ is positive and torsion free} \}$$

Let  $\mathcal{D} \subset \text{Diff}(M)$  be the connected component of  $\text{id}_M$ . Then

$$\mathcal{M} = \mathcal{X}/\mathcal{D}$$

is called the  $G_2$ -moduli space of  $M$

## Theorem (Joyce)

*The  $G_2$ -moduli space is a manifold, and the map*

$$\mathcal{M} \longrightarrow H^3(M; \mathbb{R}) \quad \text{with} \quad [\varphi] \longmapsto [\varphi]$$

*is a **local** diffeomorphism*

Not much is known about the **global** structure of  $\mathcal{M}$

# $G_2$ -geometry—Constructions I

Constructions of  $(M, g)$  with  $\text{Hol}(M, g) = G_2$

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **compact** examples



# $G_2$ -geometry—Constructions I

Constructions of  $(M, g)$  with  $\text{Hol}(M, g) = G_2$

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **compact** examples

Joyce's construction: let a “rich enough” finite subgroup  $\Gamma \subset G_2$  act on **flat**  $T^7$  with “sufficiently many” fixpoints, preserving a parallel positive 3-form  $\varphi \in \Omega^3(T^7)$

# $G_2$ -geometry—Constructions I

Constructions of  $(M, g)$  with  $\text{Hol}(M, g) = G_2$

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **compact** examples

Joyce's construction: let a “rich enough” finite subgroup  $\Gamma \subset G_2$  act on **flat**  $T^7$  with “sufficiently many” fixpoints, preserving a parallel positive 3-form  $\varphi \in \Omega^3(T^7)$

The stabilisers of fixpoints  $p \in T^7$  are isomorphic to subgroups of  $SU(2)$  or  $SU(3)$

# $G_2$ -geometry—Constructions I

Constructions of  $(M, g)$  with  $\text{Hol}(M, g) = G_2$

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **compact** examples

Joyce's construction: let a “rich enough” finite subgroup  $\Gamma \subset G_2$  act on **flat**  $T^7$  with “sufficiently many” fixpoints, preserving a parallel positive 3-form  $\varphi \in \Omega^3(T^7)$

The stabilisers of fixpoints  $p \in T^7$  are isomorphic to subgroups of  $SU(2)$  or  $SU(3)$

By gluing in suitable **noncompact Calabi-Yau manifolds** in place of the singularities, Joyce constructs a desingularisation

$$M \longrightarrow T^7/\Gamma$$

# $G_2$ -geometry—Constructions I

Constructions of  $(M, g)$  with  $\text{Hol}(M, g) = G_2$

- ▶ Bryant '87: first **non-complete** examples
- ▶ Bryant and Salamon '89: first **complete** examples
- ▶ Joyce '96: first **compact** examples

Joyce's construction: let a “rich enough” finite subgroup  $\Gamma \subset G_2$  act on **flat**  $T^7$  with “sufficiently many” fixpoints, preserving a parallel positive 3-form  $\varphi \in \Omega^3(T^7)$

The stabilisers of fixpoints  $p \in T^7$  are isomorphic to subgroups of  $SU(2)$  or  $SU(3)$

By gluing in suitable **noncompact Calabi-Yau manifolds** in place of the singularities, Joyce constructs a desingularisation

$$M \longrightarrow T^7/\Gamma$$


The gluing metric on  $M$  is close to one with holonomy  $G_2$

# $G_2$ -geometry—Constructions II

- ▶ Kovalev '03, Corti-Haskins-Nordström-Pacini '15:  
Twisted connected sums

# $G_2$ -geometry—Constructions II


- ▶ Kovalev '03, Corti-Haskins-Nordström-Pacini '15:  
Twisted connected sums



Let  $V_+$ ,  $V_-$  be asymptotically cylindrical Calabi-Yau manifolds

# $G_2$ -geometry—Constructions II


- Kovalev '03, Corti-Haskins-Nordström-Pacini '15:  
Twisted connected sums



Let  $V_+$ ,  $V_-$  be asymptotically cylindrical Calabi-Yau manifolds  
The ends are asymptotic to  $\Sigma_{\pm} \times S^1 \times \mathbb{R}$ ,  $\Sigma_{\pm}$  are K3 surfaces

# $G_2$ -geometry—Constructions II

- Kovalev '03, Corti-Haskins-Nordström-Pacini '15:  
Twisted connected sums




Let  $V_+$ ,  $V_-$  be asymptotically cylindrical Calabi-Yau manifolds  
 The ends are asymptotic to  $\Sigma_{\pm} \times S^1 \times \mathbb{R}$ ,  $\Sigma_{\pm}$  are K3 surfaces  
 Glue to  $V_- \times S^1$  to  $V_+ \times S^1$ , flipping the circles



# $G_2$ -geometry—Constructions II

- ▶ Kovalev '03, Corti-Haskins-Nordström-Pacini '15:  
Twisted connected sums



Let  $V_+$ ,  $V_-$  be asymptotically cylindrical Calabi-Yau manifolds  
The ends are asymptotic to  $\Sigma_{\pm} \times S^1 \times \mathbb{R}$ ,  $\Sigma_{\pm}$  are K3 surfaces  
Glue to  $V_- \times S^1$  to  $V_+ \times S^1$ , flipping the circles  
The gluing metric on  $M$  is close to one with holonomy  $G_2$

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space—the  $G_2$ -metric is close to one with smaller “local” holonomy groups  $SU(2)$  or  $SU(3)$

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space—the  $G_2$ -metric is close to one with smaller “local” holonomy groups  $SU(2)$  or  $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against  $G_2$ -metrics

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space—the  $G_2$ -metric is close to one with smaller “local” holonomy groups  $SU(2)$  or  $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against  $G_2$ -metrics
- ▶ Construct  $G_2$ -metrics in the interior of the moduli space



# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space—the  $G_2$ -metric is close to one with smaller “local” holonomy groups  $SU(2)$  or  $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against  $G_2$ -metrics
- ▶ Construct  $G_2$ -metrics in the interior of the moduli space
- ▶ How can families of  $G_2$ -metrics become singular?

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space—the  $G_2$ -metric is close to one with smaller “local” holonomy groups  $SU(2)$  or  $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against  $G_2$ -metrics
- ▶ Construct  $G_2$ -metrics in the interior of the moduli space
- ▶ How can families of  $G_2$ -metrics become singular?  
How far can one deform a given  $G_2$ -metric?

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space—the  $G_2$ -metric is close to one with smaller “local” holonomy groups  $SU(2)$  or  $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against  $G_2$ -metrics
- ▶ Construct  $G_2$ -metrics in the interior of the moduli space
- ▶ How can families of  $G_2$ -metrics become singular?  
How far can one deform a given  $G_2$ -metric?
- ▶ Construct  $G_2$ -metrics with prescribed singularities

# $G_2$ -geometry—Some open questions

Let us summarise first

- ▶ Only a few obstructions against  $G_2$ -holonomy are known
- ▶ Only a few examples are known—only  $\sim 10^7$  families
- ▶ Known compact examples represent points close to the boundary of the moduli space—the  $G_2$ -metric is close to one with smaller “local” holonomy groups  $SU(2)$  or  $SU(3)$

Important open problems / questions

- ▶ Find more invariants for / obstructions against  $G_2$ -metrics
- ▶ Construct  $G_2$ -metrics in the interior of the moduli space
- ▶ How can families of  $G_2$ -metrics become singular?  
How far can one deform a given  $G_2$ -metric?
- ▶ Construct  $G_2$ -metrics with prescribed singularities  
Singularities represent matter and forces in  $M$ -theory

# The $\nu$ -invariant—Spinors and $G_2$ -structures

Recall: every  $G_2$ -structure leads  
to a distinguished **nowhere vanishing spinor**  $\sigma$   
It is **parallel** if and only if  $\text{Hol}(M, g) \subset G_2$

# The $\nu$ -invariant—Spinors and $G_2$ -structures

Recall: every  $G_2$ -structure leads  
to a distinguished **nowhere vanishing spinor**  $\sigma$

It is **parallel** if and only if  $\text{Hol}(M, g) \subset G_2$

Conversely, if  $(M, g)$  is an oriented, spin 7-manifold, then it has  
nowhere vanishing spinor fields because

$$\text{rk } SM = 8 \quad > \quad 7 = \dim M$$

# The $\nu$ -invariant—Spinors and $G_2$ -structures

Recall: every  $G_2$ -structure leads  
to a distinguished **nowhere vanishing spinor**  $\sigma$

It is **parallel** if and only if  $\text{Hol}(M, g) \subset G_2$

Conversely, if  $(M, g)$  is an oriented, spin 7-manifold, then it has  
nowhere vanishing spinor fields because

$$\text{rk } SM = 8 > 7 = \dim M$$

Let  $0 \neq \sigma \in S\mathbb{R}^7$ , then

$$\{ g \in \text{Spin}(7) \mid g \cdot \sigma = \sigma \} \cong G_2$$

# The $\nu$ -invariant—Spinors and $G_2$ -structures

Recall: every  $G_2$ -structure leads  
to a distinguished **nowhere vanishing spinor**  $\sigma$

It is **parallel** if and only if  $\text{Hol}(M, g) \subset G_2$

Conversely, if  $(M, g)$  is an oriented, spin 7-manifold, then it has  
nowhere vanishing spinor fields because

$$\text{rk } SM = 8 > 7 = \dim M$$

Let  $0 \neq \sigma \in S\mathbb{R}^7$ , then

$$\{ g \in \text{Spin}(7) \mid g \cdot \sigma = \sigma \} \cong G_2$$

A nowhere vanishing spinor  $\sigma$  determines a  $G_2$ -structure



# The $\nu$ -invariant—Spinors and $G_2$ -structures

Recall: every  $G_2$ -structure leads  
to a distinguished **nowhere vanishing spinor**  $\sigma$   
It is **parallel** if and only if  $\text{Hol}(M, g) \subset G_2$

Conversely, if  $(M, g)$  is an oriented, spin 7-manifold, then it has  
nowhere vanishing spinor fields because

$$\text{rk } SM = 8 > 7 = \dim M$$

Let  $0 \neq \sigma \in S\mathbb{R}^7$ , then

$$\{g \in \text{Spin}(7) \mid g \cdot \sigma = \sigma\} \cong G_2$$

A nowhere vanishing spinor  $\sigma$  determines a  $G_2$ -structure

**Idea.** Use nowhere vanishing spinors  
to describe and distinguish  $G_2$ -structures

# The $\nu$ -invariant—Comparing $G_2$ -structures

Let  $\sigma_0, \sigma_1$  be two nowhere vanishing spinors

Extend to a spinor  $\bar{\sigma} \in \Gamma(S^+(M \times [0, 1]))$

A generic  $\bar{\sigma}$  will have nondegenerate isolated zeros because

$$\mathrm{rk} S^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

# The $\nu$ -invariant—Comparing $G_2$ -structures

Let  $\sigma_0, \sigma_1$  be two nowhere vanishing spinors

Extend to a spinor  $\bar{\sigma} \in \Gamma(S^+(M \times [0, 1]))$

A generic  $\bar{\sigma}$  will have nondegenerate isolated zeros because

$$\operatorname{rk} S^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

Orient  $S^+(M \times [0, 1])$  and count with signs

$$\Delta\nu(M; \sigma_0, \sigma_1) = 2 \cdot \#\bar{\sigma}^{-1}(0) = 2 \cdot \sum_{p \in \bar{\sigma}^{-1}(0)} \operatorname{sign}(d_p \bar{\sigma})$$

# The $\nu$ -invariant—Comparing $G_2$ -structures

Let  $\sigma_0, \sigma_1$  be two nowhere vanishing spinors

Extend to a spinor  $\bar{\sigma} \in \Gamma(\mathcal{S}^+(M \times [0, 1]))$

A generic  $\bar{\sigma}$  will have nondegenerate isolated zeros because

$$\mathrm{rk} \mathcal{S}^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

Orient  $\mathcal{S}^+(M \times [0, 1])$  and count with signs

$$\Delta\nu(M; \sigma_0, \sigma_1) = 2 \cdot \#\bar{\sigma}^{-1}(0) = 2 \cdot \sum_{p \in \bar{\sigma}^{-1}(0)} \mathrm{sign}(d_p \bar{\sigma})$$

## Theorem (Crowley-Nordström)

Let  $F: M \rightarrow M$  be a spin diffeomorphism, then

$$\Delta\nu(M; \sigma, F^* \sigma) \in 48\mathbb{Z}$$

# The $\nu$ -invariant—Comparing $G_2$ -structures

Let  $\sigma_0, \sigma_1$  be two nowhere vanishing spinors

Extend to a spinor  $\bar{\sigma} \in \Gamma(\mathcal{S}^+(M \times [0, 1]))$

A generic  $\bar{\sigma}$  will have nondegenerate isolated zeros because

$$\mathrm{rk} \mathcal{S}^+(M \times [0, 1]) = 8 = \dim(M \times [0, 1])$$

Orient  $\mathcal{S}^+(M \times [0, 1])$  and count with signs

$$\Delta\nu(M; \sigma_0, \sigma_1) = 2 \cdot \#\bar{\sigma}^{-1}(0) = 2 \cdot \sum_{p \in \bar{\sigma}^{-1}(0)} \mathrm{sign}(d_p \bar{\sigma})$$

## Theorem (Crowley-Nordström)

Let  $F: M \rightarrow M$  be a spin diffeomorphism, then

$$\Delta\nu(M; \sigma, F^* \sigma) \in 48\mathbb{Z}$$

Can we write  $\Delta\nu(M; \sigma_0, \sigma_1) = \nu(M, \sigma_0) - \nu(M, \sigma_1) \in \mathbb{Z}/48$ ?

# The $\nu$ -invariant—Cobordism definition

**Idea.** If  $M$  is spin, then  $M$  is the spin coboundary of some compact 8-manifold  $W$  (because  $\Omega_{\text{Spin}}^7 = 0$ )

Extend  $\sigma$  to  $\bar{\sigma} \in \Gamma(\mathcal{S}^+ W)$ , count  $\bar{\sigma}^{-1}(0)$

# The $\nu$ -invariant—Cobordism definition

**Idea.** If  $M$  is spin, then  $M$  is the spin coboundary of some compact 8-manifold  $W$  (because  $\Omega_{\text{Spin}}^7 = 0$ )

Extend  $\sigma$  to  $\bar{\sigma} \in \Gamma(\mathcal{S}^+ W)$ , count  $\bar{\sigma}^{-1}(0)$

The result depends on  $W$ —**not well-defined** for  $(M, \sigma)$ !

# The $\nu$ -invariant—Cobordism definition

**Idea.** If  $M$  is spin, then  $M$  is the spin coboundary of some compact 8-manifold  $W$  (because  $\Omega_{\text{Spin}}^7 = 0$ )

Extend  $\sigma$  to  $\bar{\sigma} \in \Gamma(\mathcal{S}^+ W)$ , count  $\bar{\sigma}^{-1}(0)$

The result depends on  $W$ —**not well-defined** for  $(M, \sigma)$ !

## Definition (Crowley-Nordström)

Assume that  $M = \partial W$  with  $W$  spin, compact. Define

$$\nu(M, \sigma) = \chi(W) - 3 \operatorname{sign}(W) - 2\#\bar{\sigma}^{-1}(0) \pmod{48}$$

- ▶  $\chi(W)$ —Euler characteristic of  $W$
- ▶  $\operatorname{sign}(W)$ —signature of  $W$



# The $\nu$ -invariant—Cobordism definition

**Idea.** If  $M$  is spin, then  $M$  is the spin coboundary of some compact 8-manifold  $W$  (because  $\Omega_{\text{Spin}}^7 = 0$ )

Extend  $\sigma$  to  $\bar{\sigma} \in \Gamma(\mathcal{S}^+ W)$ , count  $\bar{\sigma}^{-1}(0)$

The result depends on  $W$ —**not well-defined** for  $(M, \sigma)$ !

## Definition (Crowley-Nordström)

Assume that  $M = \partial W$  with  $W$  spin, compact. Define

$$\nu(M, \sigma) = \chi(W) - 3 \operatorname{sign}(W) - 2\#\bar{\sigma}^{-1}(0) \pmod{48}$$

- ▶  $\chi(W)$ —Euler characteristic of  $W$
- ▶  $\operatorname{sign}(W)$ —signature of  $W$

## Theorem (Crowley-Nordström)

$$\Delta\nu(M; \sigma_0, \sigma_1) = \nu(M, \sigma_0) - \nu(M, \sigma_1) \in \mathbb{Z}/48$$

# The $\nu$ -invariant—Index theory

**Problem.** Given  $M$ , how to determine  $W$  with  $M = \partial W$ ?

# The $\nu$ -invariant—Index theory

**Problem.** Given  $M$ , how to determine  $W$  with  $M = \partial W$ ?  
Instead employ **index theorems** to define  $\nu(M, \sigma)$  on  $M$  itself

# The $\nu$ -invariant—Index theory

**Problem.** Given  $M$ , how to determine  $W$  with  $M = \partial W$ ?  
Instead employ **index theorems** to define  $\nu(M, \sigma)$  on  $M$  itself

## Theorem (Atiyah-Patodi-Singer)

*Assume that  $M = \partial W$  with  $W$  compact, then*

$$\text{sign}(W) = \int_W L(\nabla) - \eta(B_M)$$

- ▶ sign—signature
- ▶  $L(\nabla)$ —Hirzebruch  $L$ -form in  $\Omega^\bullet(W)$
- ▶  $B_M$ —odd signature operator  $*d \pm d*$  on  $\Omega^{\text{ev}}(M)$
- ▶  $\eta$ —Atiyah-Patodi-Singer  $\eta$ -invariant

# The $\nu$ -invariant—Index theory

**Problem.** Given  $M$ , how to determine  $W$  with  $M = \partial W$ ?  
Instead employ **index theorems** to define  $\nu(M, \sigma)$  on  $M$  itself

## Theorem (Atiyah-Patodi-Singer)

*Assume that  $M = \partial W$  with  $W$  compact, spin, then*

$$\text{ind}(D_W) = \int_W \hat{A}(\nabla) - \frac{\eta + h}{2}(D_M)$$

- ▶  $D_W, D_M$ —spin Dirac operators on  $\Gamma(S^\pm W)$  and  $\Gamma(SM)$
- ▶  $\text{ind}$ —Fredholm index
- ▶  $\hat{A}(\nabla)$ —Atiyah  $\hat{A}$ -form in  $\Omega^\bullet(W)$
- ▶  $\eta$ —Atiyah-Patodi-Singer  $\eta$ -invariant
- ▶  $h = \dim \ker$

# The $\nu$ -invariant—Index theory

**Problem.** Given  $M$ , how to determine  $W$  with  $M = \partial W$ ?  
Instead employ **index theorems** to define  $\nu(M, \sigma)$  on  $M$  itself

## Theorem (Gauß-Bonnet-Chern)

*Assume that  $M = \partial W$  with  $W$  compact, then*

$$\chi(W) = \int_W e(\nabla)$$

- ▶  $\chi$ —Euler characteristic
- ▶  $e(\nabla)$ —Euler form in  $\Omega^\bullet(W)$

# The $\nu$ -invariant—Index theory

**Problem.** Given  $M$ , how to determine  $W$  with  $M = \partial W$ ?  
Instead employ **index theorems** to define  $\nu(M, \sigma)$  on  $M$  itself

## Theorem (Matthai-Quillen)

*Assume that  $M = \partial W$  with  $W$  compact, spin, then*

$$\# \bar{\sigma}^{-1}(0) = \int_W e(\nabla^{S^+W}) - \int_M \sigma^* \psi(\nabla^{SM})$$

- ▶  $e(\nabla^{S^+W})$ —Euler form of the spinor bundle
- ▶  $\psi(\nabla^{SM}, g^{SM})$ —Mathai-Quillen form in  $\Omega^\bullet(SM)$

# The $\nu$ -invariant—Analytic description

Magic formula

$$2e(\nabla^{S^+W}) = e(\nabla) + 48\hat{A}(\nabla)^{[8]} - 3L(\nabla)^{[8]} \in \Omega^8(W)$$



# The $\nu$ -invariant—Analytic description

Magic formula

$$2e(\nabla^{S^+W}) = e(\nabla) + 48\hat{A}(\nabla)^{[8]} - 3L(\nabla)^{[8]} \in \Omega^8(W)$$

Theorem (Crowley-G-Nordström)

$$\nu(M, \sigma) = 2 \int_M \sigma^* \psi(\nabla^{SM}, g^{SM}) - 24(\eta + h)(D_M) + 3\eta(B_M) \in \mathbb{Z}/48$$

# The $\nu$ -invariant—Analytic description

Magic formula

$$2e(\nabla^{S^+W}) = e(\nabla) + 48\hat{A}(\nabla)^{[8]} - 3L(\nabla)^{[8]} \in \Omega^8(W)$$

Theorem (Crowley-G-Nordström)

$$\nu(M, \sigma) = 2 \int_M \sigma^* \psi(\nabla^{SM}, g^{SM}) - 24(\eta + h)(D_M) + 3\eta(B_M) \in \mathbb{Z}/48$$

- ▶  $\psi(\nabla^{SM}, g^{SM})$ —Mathai-Quillen form in  $\Omega^\bullet(SM)$
- ▶  $D_M$ —spin Dirac operator on  $\Gamma(SM)$
- ▶  $B_M$ —odd signature operator  $*d \pm d*$  on  $\Omega^{\text{ev}}(M)$
- ▶  $h$ —dimension of the kernel
- ▶  $\eta$ —Atiyah-Patodi-Singer  $\eta$ -invariant

$$\eta(A) = \left( \sum_{\lambda \in \text{spec}(A)} \text{sign } \lambda |\lambda|^t \right) \Big|_{t=0} = \int_0^\infty \text{tr}(A e^{-tA^2}) \frac{dt}{\sqrt{\pi t}}$$

# The $\nu$ -invariant—The extended $\nu$ -invariant

In the case of  $G_2$ -holonomy, things simplify

- ▶  $\sigma$  is parallel, so  $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$

# The $\nu$ -invariant—The extended $\nu$ -invariant

In the case of  $G_2$ -holonomy, things simplify

- ▶  $\sigma$  is parallel, so  $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶  $\ker D_M = \langle \sigma \rangle$ , so  $h(D_M) = 1$

# The $\nu$ -invariant—The extended $\nu$ -invariant

In the case of  $G_2$ -holonomy, things simplify

- ▶  $\sigma$  is parallel, so  $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶  $\ker D_M = \langle \sigma \rangle$ , so  $h(D_M) = 1$
- ▶  $\eta(D_M) \in \mathbb{R}$  is smooth on the  $G_2$ -moduli space  $\mathcal{M}$

# The $\nu$ -invariant—The extended $\nu$ -invariant

In the case of  $G_2$ -holonomy, things simplify

- ▶  $\sigma$  is parallel, so  $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶  $\ker D_M = \langle \sigma \rangle$ , so  $h(D_M) = 1$
- ▶  $\eta(D_M) \in \mathbb{R}$  is smooth on the  $G_2$ -moduli space  $\mathcal{M}$

## Definition (Crowley-G-Nordström)

Let  $(M, g)$  be a compact manifold with  $\text{Hol}(M, g) \cong G_2$

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

# The $\nu$ -invariant—The extended $\nu$ -invariant

In the case of  $G_2$ -holonomy, things simplify

- ▶  $\sigma$  is parallel, so  $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶  $\ker D_M = \langle \sigma \rangle$ , so  $h(D_M) = 1$
- ▶  $\eta(D_M) \in \mathbb{R}$  is smooth on the  $G_2$ -moduli space  $\mathcal{M}$

## Definition (Crowley-G-Nordström)

Let  $(M, g)$  be a compact manifold with  $\text{Hol}(M, g) \cong G_2$

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

- ▶  $\nu(M, \sigma) = \bar{\nu}(M, g) - 24 \pmod{48}$

# The $\nu$ -invariant—The extended $\nu$ -invariant

In the case of  $G_2$ -holonomy, things simplify

- ▶  $\sigma$  is parallel, so  $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶  $\ker D_M = \langle \sigma \rangle$ , so  $h(D_M) = 1$
- ▶  $\eta(D_M) \in \mathbb{R}$  is smooth on the  $G_2$ -moduli space  $\mathcal{M}$

## Definition (Crowley-G-Nordström)

Let  $(M, g)$  be a compact manifold with  $\text{Hol}(M, g) \cong G_2$

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

- ▶  $\nu(M, \sigma) = \bar{\nu}(M, g) - 24 \pmod{48}$
- ▶  $\bar{\nu}(M, g)$  is locally constant on  $\mathcal{M}$



# The $\nu$ -invariant—The extended $\nu$ -invariant

In the case of  $G_2$ -holonomy, things simplify

- ▶  $\sigma$  is parallel, so  $\sigma^*\psi(\nabla^{SM}, g^{SM}) = 0$
- ▶  $\ker D_M = \langle \sigma \rangle$ , so  $h(D_M) = 1$
- ▶  $\eta(D_M) \in \mathbb{R}$  is smooth on the  $G_2$ -moduli space  $\mathcal{M}$

## Definition (Crowley-G-Nordström)

Let  $(M, g)$  be a compact manifold with  $\text{Hol}(M, g) \cong G_2$

$$\bar{\nu}(M, g) = 3\eta(B_M) - 24\eta(D_M) \in \mathbb{Z}$$

- ▶  $\nu(M, \sigma) = \bar{\nu}(M, g) - 24 \pmod{48}$
- ▶  $\bar{\nu}(M, g)$  is locally constant on  $\mathcal{M}$
- ▶  $\bar{\nu}(M, g) = 0$  if  $M$  admits an orientation reversing isometry

# The $\nu$ -invariant—First examples

What about the known examples by Joyce and Kovalev?

- ▶  $\bar{\nu}(M, g) = 0$  for all twisted connected sums

# The $\nu$ -invariant—First examples

What about the known examples by Joyce and Kovalev?

- ▶  $\bar{\nu}(M, g) = 0$  for all twisted connected sums
- ▶  $\bar{\nu}(M, g) = 0$  for some of Joyce's examples
  - Some are twisted connected sums
  - Some have orientation reversing isometries

# The $\nu$ -invariant—First examples

What about the known examples by Joyce and Kovalev?

- ▶  $\bar{\nu}(M, g) = 0$  for all twisted connected sums
- ▶  $\bar{\nu}(M, g) = 0$  for some of Joyce's examples
  - Some are twisted connected sums
  - Some have orientation reversing isometries

**Question.** Is  $\bar{\nu}(M, g) = 0$  whenever  $\text{Hol}(M, g) = G_2$ ?

# The $\nu$ -invariant—First examples

What about the known examples by Joyce and Kovalev?

- ▶  $\bar{\nu}(M, g) = 0$  for all twisted connected sums
- ▶  $\bar{\nu}(M, g) = 0$  for some of Joyce's examples
  - Some are twisted connected sums
  - Some have orientation reversing isometries

**Question.** Is  $\bar{\nu}(M, g) = 0$  whenever  $\text{Hol}(M, g) = G_2$ ?

- ▶ If **yes**, then  $\bar{\nu}(M, g) \neq 0$  or  $\nu(M, \sigma) \neq 24$  is a new obstruction against  $G_2$ -holonomy

# The $\nu$ -invariant—First examples

What about the known examples by Joyce and Kovalev?

- ▶  $\bar{\nu}(M, g) = 0$  for all twisted connected sums
- ▶  $\bar{\nu}(M, g) = 0$  for some of Joyce's examples
  - Some are twisted connected sums
  - Some have orientation reversing isometries

**Question.** Is  $\bar{\nu}(M, g) = 0$  whenever  $\text{Hol}(M, g) = G_2$ ?

- ▶ If **yes**, then  $\bar{\nu}(M, g) \neq 0$  or  $\nu(M, \sigma) \neq 24$  is a new obstruction against  $G_2$ -holonomy
- ▶ If **no**, then  $\bar{\nu}(M, g)$  is a non-trivial new invariant

# The $\nu$ -invariant—First examples

What about the known examples by Joyce and Kovalev?

- ▶  $\bar{\nu}(M, g) = 0$  for all twisted connected sums
- ▶  $\bar{\nu}(M, g) = 0$  for some of Joyce's examples
  - Some are twisted connected sums
  - Some have orientation reversing isometries

**Question.** Is  $\bar{\nu}(M, g) = 0$  whenever  $\text{Hol}(M, g) = G_2$ ?

- ▶ If **yes**, then  $\bar{\nu}(M, g) \neq 0$  or  $\nu(M, \sigma) \neq 24$  is a new obstruction against  $G_2$ -holonomy
- ▶ If **no**, then  $\bar{\nu}(M, g)$  is a non-trivial new invariant

**Answer.** We will construct examples with  $\bar{\nu}(M, g) \neq 0$

# The $\nu$ -invariant—First examples

What about the known examples by Joyce and Kovalev?

- ▶  $\bar{\nu}(M, g) = 0$  for all twisted connected sums
- ▶  $\bar{\nu}(M, g) = 0$  for some of Joyce's examples
  - Some are twisted connected sums
  - Some have orientation reversing isometries

**Question.** Is  $\bar{\nu}(M, g) = 0$  whenever  $\text{Hol}(M, g) = G_2$ ?


- ▶ If **yes**, then  $\bar{\nu}(M, g) \neq 0$  or  $\nu(M, \sigma) \neq 24$  is a new obstruction against  $G_2$ -holonomy
- ▶ If **no**, then  $\bar{\nu}(M, g)$  is a non-trivial new invariant

**Answer.** We will construct examples with  $\bar{\nu}(M, g) \neq 0$   
Using  $\bar{\nu}(M, g)$ , we will show that for some particular  $M$ ,  
the  $G_2$ -moduli space  $\mathcal{M}$  has several connected components




# Extra twisted connected sums—Construction

Recall twisted connected sums




Gluing of tori at angle  $\vartheta = \frac{\pi}{2}$



# Extra twisted connected sums—Construction

## Extra twisted connected sums




Assume that  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  acts both on  $V_{\pm}$  and on  $S^1_{\pm,ext}$

The induced action on  $\partial V_{\pm}$  has to fix  $\Sigma_{\pm}$  pointwise

The actions on  $S^1_{\pm,int}$  and  $S^1_{\pm,ext}$  have to be free

# Extra twisted connected sums—Construction

## Extra twisted connected sums



Assume that  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  acts both on  $V_{\pm}$  and on  $S^1_{\pm, \text{ext}}$


The induced action on  $\partial V_{\pm}$  has to fix  $\Sigma_{\pm}$  pointwise

The actions on  $S^1_{\pm, \text{int}}$  and  $S^1_{\pm, \text{ext}}$  have to be free

Then  $(S^1_{\pm, \text{int}} \times S^1_{\pm, \text{ext}})/\Gamma_{\pm}$  is again a flat 2-torus

# Extra twisted connected sums—Construction

## Extra twisted connected sums



Assume that  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  acts both on  $V_{\pm}$  and on  $S^1_{\pm,ext}$

The induced action on  $\partial V_{\pm}$  has to fix  $\Sigma_{\pm}$  pointwise


The actions on  $S^1_{\pm,int}$  and  $S^1_{\pm,ext}$  have to be free

Then  $(S^1_{\pm,int} \times S^1_{\pm,ext})/\Gamma_{\pm}$  is again a flat 2-torus


If both the tori and the K3 surfaces are isometric,  
we can glue  $M_{\pm} = (V_{\pm} \times S^1_{\pm,ext})/\Gamma_{\pm}$  at various angles  $\vartheta$

# Extra twisted connected sums—Construction

## Extra twisted connected sums




## Modified gluing of tori at angle $\vartheta = \frac{3}{4}\pi$






# Extra twisted connected sums—Construction

## Extra twisted connected sums



## Modified gluing of tori at angle $\vartheta = \arccos(\frac{1}{\sqrt{6}})$



# Extra twisted connected sums—The gluing formula

Compute the extended  $\nu$ -invariant on  $M = M_+ \cup_X M_-$

## Theorem (Bunke, Kirk-Lesch)

*For suitable boundary conditions  $L_{\pm}$ ,*

$$\eta(D_M) = \eta_{\text{APS}}(D_{M_+}; L_+) + \eta_{\text{APS}}(D_{M_-}; L_-) + m_{\ker(D_X)}(L_+, L_-)$$

$$\eta(B_M) = \eta_{\text{APS}}(B_{M_+}; L_+) + \eta_{\text{APS}}(B_{M_-}; L_-) + m_{H^3(X)}(L_+, L_-)$$



# Extra twisted connected sums—The gluing formula

Compute the extended  $\nu$ -invariant on  $M = M_+ \cup_X M_-$

## Theorem (Bunke, Kirk-Lesch)

For suitable boundary conditions  $L_{\pm}$ ,

$$\eta(D_M) = \eta_{\text{APS}}(D_{M_+}; L_+) + \eta_{\text{APS}}(D_{M_-}; L_-) + m_{\ker(D_X)}(L_+, L_-)$$

$$\eta(B_M) = \eta_{\text{APS}}(B_{M_+}; L_+) + \eta_{\text{APS}}(B_{M_-}; L_-) + m_{H^3(X)}(L_+, L_-)$$

## Theorem (Crowley-G-Nordström)

Let  $(M, g)$  be an extra twisted connected sum with gluing angle  $\vartheta$  and  $\rho = \pi - 2\vartheta$ , then there exists  $N \in \mathbb{Z}$  such that

$$3m_{H^3(X)}(L_+, L_-) - 24m_{\ker(D_X)}(L_+, L_-) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$$

# Extra twisted connected sums—The gluing formula

Compute the extended  $\nu$ -invariant on  $M = M_+ \cup_X M_-$

## Theorem (Bunke, Kirk-Lesch)

For suitable boundary conditions  $L_{\pm}$ ,

$$\eta(D_M) = \eta_{\text{APS}}(D_{M_+}; L_+) + \eta_{\text{APS}}(D_{M_-}; L_-) + m_{\ker(D_X)}(L_+, L_-)$$

$$\eta(B_M) = \eta_{\text{APS}}(B_{M_+}; L_+) + \eta_{\text{APS}}(B_{M_-}; L_-) + m_{H^3(X)}(L_+, L_-)$$

## Theorem (Crowley-G-Nordström)

Let  $(M, g)$  be an extra twisted connected sum with gluing angle  $\vartheta$  and  $\rho = \pi - 2\vartheta$ , then there exists  $N \in \mathbb{Z}$  such that

$$3m_{H^3(X)}(L_+, L_-) - 24m_{\ker(D_X)}(L_+, L_-) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$$

One computes  $N$  from the maps  $H^2(V_{\pm}) \rightarrow H^2(\Sigma_+) \cong H^2(\Sigma_-)$

# Extra twisted connected sums—The gluing formula

Compute the extended  $\nu$ -invariant on  $M = M_+ \cup_X M_-$

## Theorem (Bunke, Kirk-Lesch)

For suitable boundary conditions  $L_{\pm}$ ,

$$\eta(D_M) = \eta_{\text{APS}}(D_{M_+}; L_+) + \eta_{\text{APS}}(D_{M_-}; L_-) + m_{\ker(D_X)}(L_+, L_-)$$

$$\eta(B_M) = \eta_{\text{APS}}(B_{M_+}; L_+) + \eta_{\text{APS}}(B_{M_-}; L_-) + m_{H^3(X)}(L_+, L_-)$$

## Theorem (Crowley-G-Nordström)

Let  $(M, g)$  be an extra twisted connected sum with gluing angle  $\vartheta$  and  $\rho = \pi - 2\vartheta$ , then there exists  $N \in \mathbb{Z}$  such that

$$3m_{H^3(X)}(L_+, L_-) - 24m_{\ker(D_X)}(L_+, L_-) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$$

Note that  $\frac{\rho}{\pi}$  can be **irrational**, e.g., for  $\vartheta = \arccos(\frac{1}{\sqrt{6}})$

# Extra twisted connected sums—First results

## Theorem (Crowley-G-Nordström)

*If  $\Gamma_{\pm} \cong \{0\}$  or  $\mathbb{Z}/2$  then*

$$\eta_{\text{APS}}(D_{V_{\pm} \times S^1}; L_{\pm}) = \eta_{\text{APS}}(B_{V_{\pm} \times S^1}; L_{\pm}) = 0$$

# Extra twisted connected sums—First results

## Theorem (Crowley-G-Nordström)

*If  $\Gamma_{\pm} \cong \{0\}$  or  $\mathbb{Z}/2$  then*

$$\eta_{\text{APS}}(D_{V_{\pm} \times S^1}; L_{\pm}) = \eta_{\text{APS}}(B_{V_{\pm} \times S^1}; L_{\pm}) = 0$$

From the gluing formula, we derive the value of  $\bar{\nu}$

## Corollary

*If  $\Gamma_{\pm} \cong \{0\}$  or  $\mathbb{Z}/2$  then*

$$\bar{\nu}(M, g) = -72 \frac{\rho}{\pi} - 3N \operatorname{sign} \rho$$

# Extra twisted connected sums—Examples I

## Example (Crowley-G-Nordström)

There exists a spin 7-manifold  $M$  with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{97}, \quad \operatorname{div} p_1(TM) = 4$$

admitting three different  $G_2$ -holonomy metrics  $g_1, g_2, g_3$  with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 36, \quad \bar{\nu}(M, g_3) = -36.$$

# Extra twisted connected sums—Examples I

## Example (Crowley-G-Nordström)

There exists a spin 7-manifold  $M$  with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{97}, \quad \operatorname{div} p_1(TM) = 4$$

admitting three different  $G_2$ -holonomy metrics  $g_1, g_2, g_3$  with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 36, \quad \bar{\nu}(M, g_3) = -36.$$

Hence, the  $G_2$ -moduli space of  $M$  is disconnected

# Extra twisted connected sums—Examples I

## Example (Crowley-G-Nordström)

There exists a spin 7-manifold  $M$  with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{97}, \quad \operatorname{div} p_1(TM) = 4$$

admitting three different  $G_2$ -holonomy metrics  $g_1, g_2, g_3$  with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 36, \quad \bar{\nu}(M, g_3) = -36.$$

Hence, the  $G_2$ -moduli space of  $M$  is disconnected

The metric  $g_1$  comes from a rectangular twisted connected sum

The metrics  $g_2, g_3$  come from extra twisted connected sums with  $\Gamma_+ \cong \mathbb{Z}/2$ ,  $\Gamma_- \cong \{0\}$  and with gluing angles  $\frac{\pi}{4}$



# Extra twisted connected sums—Examples II

## Example (Crowley-G-Nordström)

There exists a spin 7-manifold  $M$  with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{109}, \quad \operatorname{div} p_1(TM) = 4$$

admitting three different  $G_2$ -holonomy metrics  $g_1, g_2, g_3$  with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 48, \quad \bar{\nu}(M, g_3) = -48.$$

In particular  $\nu(M, \sigma_1) = \nu(M, \sigma_2) = \nu(M, \sigma_3) = 0$ , and one can show that the underlying  $G_2$ -structures are homotopic.

# Extra twisted connected sums—Examples II

## Example (Crowley-G-Nordström)

There exists a spin 7-manifold  $M$  with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{109}, \quad \operatorname{div} p_1(TM) = 4$$

admitting three different  $G_2$ -holonomy metrics  $g_1, g_2, g_3$  with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 48, \quad \bar{\nu}(M, g_3) = -48.$$

In particular  $\nu(M, \sigma_1) = \nu(M, \sigma_2) = \nu(M, \sigma_3) = 0$ , and one can show that the underlying  $G_2$ -structures are homotopic.

Hence, one  $G_2$ -structure can give rise to several connected components of the  $G_2$ -moduli space

# Extra twisted connected sums—Examples II

## Example (Crowley-G-Nordström)

There exists a spin 7-manifold  $M$  with

$$H^1(M) \cong H^2(M) \cong 0, \quad H^4(M) \cong \mathbb{Z}^{109}, \quad \operatorname{div} p_1(TM) = 4$$

admitting three different  $G_2$ -holonomy metrics  $g_1, g_2, g_3$  with

$$\bar{\nu}(M, g_1) = 0, \quad \bar{\nu}(M, g_2) = 48, \quad \bar{\nu}(M, g_3) = -48.$$

In particular  $\nu(M, \sigma_1) = \nu(M, \sigma_2) = \nu(M, \sigma_3) = 0$ , and one can show that the underlying  $G_2$ -structures are homotopic.

Hence, one  $G_2$ -structure can give rise to several connected components of the  $G_2$ -moduli space

The metric  $g_1$  comes from a rectangular twisted connected sum

The metrics  $g_2, g_3$  come from extra twisted connected sums with  $\Gamma_+ \cong \Gamma_- \cong \mathbb{Z}/2$  and with gluing angles  $\frac{\pi}{6}$

# Extra twisted connected sums— $G_2$ -Bordism

Let  $(M, g)$  be an extra twisted connected sum with  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  where  $k_+, k_- \in \{1, 2\}$  and with  $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$

In this case, 3 divides  $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$

# Extra twisted connected sums— $G_2$ -Bordism

Let  $(M, g)$  be an extra twisted connected sum with  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  where  $k_+, k_- \in \{1, 2\}$  and with  $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$   
In this case, 3 divides  $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$

Note that  $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

One can show that  $M$  is  $G_2$ -nullbordant if and only if  $3 \mid \nu(M, \sigma)$

Hence there exists a compact  $W$  with a  $G_2$ -structure such that  $\partial W = M$  if and only if  $3 \mid \nu(M, \sigma)$

# Extra twisted connected sums— $G_2$ -Bordism

Let  $(M, g)$  be an extra twisted connected sum with  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  where  $k_+, k_- \in \{1, 2\}$  and with  $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$   
In this case, 3 divides  $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$

Note that  $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

One can show that  $M$  is  **$G_2$ -nullbordant** if and only if  $3 \mid \nu(M, \sigma)$

Hence there exists a compact  $W$  with a  $G_2$ -structure such that  $\partial W = M$  if and only if  $3 \mid \nu(M, \sigma)$

**Question.** Is  $3 \mid \nu(M, \sigma)$  an obstruction against  $G_2$ -holonomy?  
Is the  $G_2$ -bordism class of  $(M, \sigma)$  an obstruction?

# Extra twisted connected sums— $G_2$ -Bordism

Let  $(M, g)$  be an extra twisted connected sum with  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  where  $k_+, k_- \in \{1, 2\}$  and with  $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$   
In this case, 3 divides  $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$

Note that  $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

One can show that  $M$  is  **$G_2$ -nullbordant** if and only if  $3 \mid \nu(M, \sigma)$

Hence there exists a compact  $W$  with a  $G_2$ -structure such that  $\partial W = M$  if and only if  $3 \mid \nu(M, \sigma)$

**Question.** Is  $3 \mid \nu(M, \sigma)$  an obstruction against  $G_2$ -holonomy?  
Is the  $G_2$ -bordism class of  $(M, \sigma)$  an obstruction?

**Answer.** No, there are examples with  $3 \nmid \nu(M, \sigma)$ .

# Extra twisted connected sums— $G_2$ -Bordism

Let  $(M, g)$  be an extra twisted connected sum with  $\Gamma_{\pm} \cong \mathbb{Z}/k_{\pm}$  where  $k_+, k_- \in \{1, 2\}$  and with  $\vartheta \in \{\pm\frac{\pi}{6}, \pm\frac{\pi}{4}, \pm\frac{\pi}{3}, \frac{\pi}{2}\}$   
In this case, 3 divides  $\bar{\nu}(M, g) = -72\frac{\rho}{\pi} - 3N \operatorname{sign} \rho$

Note that  $\Omega_{G_2}^7 \cong \mathbb{Z}/3$

One can show that  $M$  is  $G_2$ -nullbordant if and only if  $3 \mid \nu(M, \sigma)$

Hence there exists a compact  $W$  with a  $G_2$ -structure such that  $\partial W = M$  if and only if  $3 \mid \nu(M, \sigma)$

**Question.** Is  $3 \mid \nu(M, \sigma)$  an obstruction against  $G_2$ -holonomy?  
Is the  $G_2$ -bordism class of  $(M, \sigma)$  an obstruction?

**Answer.** No, there are examples with  $3 \nmid \nu(M, \sigma)$ .

**Note.** Recall that if  $k_+ \geq 3$  or  $k_- \geq 3$ , then there are gluing angles  $\vartheta$  such that  $\frac{\rho}{\pi} \notin \mathbb{Q}$

Because  $\bar{\nu}(M, g) \in \mathbb{Z}$ , expect more contributions to  $\bar{\nu}(M, g)$



# Extra twisted connected sums—Adiabatic limits

Let  $M_{\pm} = V_{\pm} \times S^1_{\pm, \text{ext}}$ , rescale  $S^1_{\pm, \text{ext}}$  by  $\varepsilon > 0$  to get  $M_{\pm, \varepsilon}$   
The limit  $\varepsilon \rightarrow 0$  is called **adiabatic limit**

# Extra twisted connected sums—Adiabatic limits

Let  $M_{\pm} = V_{\pm} \times S^1_{\pm, \text{ext}}$ , rescale  $S^1_{\pm, \text{ext}}$  by  $\varepsilon > 0$  to get  $M_{\pm, \varepsilon}$   
The limit  $\varepsilon \rightarrow 0$  is called **adiabatic limit**

**Theorem (Bismut-Cheeger, Dai, G-Nordström)**

*Let  $\gamma \in \Gamma_{\pm}$  be the generator that acts by  $\frac{2\pi}{k_{\pm}}$  on  $S^1_{\pm, \text{ext}}$*

*Let  $V_{\pm, j}$  be the set of isolated fixpoints of  $\gamma^j$  on  $V_{\pm}$*

*Let  $e^{i\alpha_{j,1}(p)}$ ,  $e^{i\alpha_{j,2}(p)}$ ,  $e^{i\alpha_{j,3}(p)}$  be the eigenvalues of  $\gamma^j$  on  $T_p V_{\pm}$*

$$\begin{aligned}\bar{\nu}(M_{\pm}) &= \lim_{\varepsilon \rightarrow 0} (3\eta(B_{M_{\pm, \varepsilon}}, L_{\pm}) - 24\eta(D_{M_{\pm, \varepsilon}}, L_{\pm})) \\ &= \frac{3}{k_{\pm}} \sum_{j=1}^{k_{\pm}-1} \cot \frac{\pi j}{k_{\pm}} \sum_{p \in V_{\pm, j}} \frac{\cos \frac{\alpha_{j,1}(p)}{2} \cos \frac{\alpha_{j,2}(p)}{2} \cos \frac{\alpha_{j,3}(p)}{2} - 1}{\sin \frac{\alpha_{j,1}(p)}{2} \sin \frac{\alpha_{j,2}(p)}{2} \sin \frac{\alpha_{j,3}(p)}{2}} \\ &\in \mathbb{Q}\end{aligned}$$

# Extra twisted connected sums—Adiabatic limits

Let  $M_{\pm} = V_{\pm} \times S^1_{\pm, \text{ext}}$ , rescale  $S^1_{\pm, \text{ext}}$  by  $\varepsilon > 0$  to get  $M_{\pm, \varepsilon}$   
The limit  $\varepsilon \rightarrow 0$  is called **adiabatic limit**

**Theorem (Bismut-Cheeger, Dai, G-Nordström)**

*Let  $\gamma \in \Gamma_{\pm}$  be the generator that acts by  $\frac{2\pi}{k_{\pm}}$  on  $S^1_{\pm, \text{ext}}$*

*Let  $V_{\pm, j}$  be the set of isolated fixpoints of  $\gamma^j$  on  $V_{\pm}$*

*Let  $e^{i\alpha_{j,1}(p)}$ ,  $e^{i\alpha_{j,2}(p)}$ ,  $e^{i\alpha_{j,3}(p)}$  be the eigenvalues of  $\gamma^j$  on  $T_p V_{\pm}$*

$$\begin{aligned}\bar{\nu}(M_{\pm}) &= \lim_{\varepsilon \rightarrow 0} (3\eta(B_{M_{\pm, \varepsilon}}, L_{\pm}) - 24\eta(D_{M_{\pm, \varepsilon}}, L_{\pm})) \\ &= \frac{3}{k_{\pm}} \sum_{j=1}^{k_{\pm}-1} \cot \frac{\pi j}{k_{\pm}} \sum_{p \in V_{\pm, j}} \frac{\cos \frac{\alpha_{j,1}(p)}{2} \cos \frac{\alpha_{j,2}(p)}{2} \cos \frac{\alpha_{j,3}(p)}{2} - 1}{\sin \frac{\alpha_{j,1}(p)}{2} \sin \frac{\alpha_{j,2}(p)}{2} \sin \frac{\alpha_{j,3}(p)}{2}} \\ &\in \mathbb{Q}\end{aligned}$$

There are examples with  $k_{\pm} \geq 3$  where  $\bar{\nu}(M_{\pm}) \neq 0$

# Extra twisted connected sums—Variational formula

The  $\eta$ -invariants of  $M_{\pm,\varepsilon}$  depend on  $\varepsilon$

## Theorem (Bismut-Cheeger, Dai-Freed)

*The variational formula for the  $\eta$ -invariant of a Dirac type operator a manifold with boundary consists of*

- ▶ *the integral of a Chern-Simons form over the interior*
- ▶ *the degree-1-component of an  $\eta$ -form on the boundary*

# Extra twisted connected sums—Variational formula

The  $\eta$ -invariants of  $M_{\pm,\varepsilon}$  depend on  $\varepsilon$

## Theorem (Bismut-Cheeger, Dai-Freed)

*The variational formula for the  $\eta$ -invariant of a Dirac type operator a manifold with boundary consists of*

- ▶ *the integral of a Chern-Simons form over the interior*
- ▶ *the degree-1-component of an  $\eta$ -form on the boundary*

The interior contribution vanishes because  $M_{\pm,\varepsilon}$  is a product

# Extra twisted connected sums—Variational formula

The  $\eta$ -invariants of  $M_{\pm,\varepsilon}$  depend on  $\varepsilon$

## Theorem (Bismut-Cheeger, Dai-Freed)

*The variational formula for the  $\eta$ -invariant of a Dirac type operator a manifold with boundary consists of*

- ▶ *the integral of a Chern-Simons form over the interior*
- ▶ *the degree-1-component of an  $\eta$ -form on the boundary*

The interior contribution vanishes because  $M_{\pm,\varepsilon}$  is a product

The boundary contribution comes from the  $\eta$ -form  $\tilde{\eta}(\mathbb{A})$  of the family of tori  $(S^1_{\pm,\text{int}} \times \varepsilon S^1_{\pm,\text{ext}})/\Gamma_{\pm}$  for  $\varepsilon \in (0, 1)$

# Extra twisted connected sums—Variational formula

The  $\eta$ -invariants of  $M_{\pm,\varepsilon}$  depend on  $\varepsilon$

## Theorem (Bismut-Cheeger, Dai-Freed)

*The variational formula for the  $\eta$ -invariant of a Dirac type operator on a manifold with boundary consists of*

- ▶ *the integral of a Chern-Simons form over the interior*
- ▶ *the degree-1-component of an  $\eta$ -form on the boundary*

The interior contribution vanishes because  $M_{\pm,\varepsilon}$  is a product


The boundary contribution comes from the  $\eta$ -form  $\tilde{\eta}(\mathbb{A})$  of the family of tori  $(S^1_{\pm,\text{int}} \times \varepsilon S^1_{\pm,\text{ext}})/\Gamma_{\pm}$  for  $\varepsilon \in (0, 1)$

Integrate over  $\varepsilon$  to get the missing last contribution to  $\bar{\nu}(M, g)$

# Extra twisted connected sums—Matching the tori

**Idea.** Represent adiabatic limits as rays in the hyperbolic plane

Lattices  $\Lambda_- \subset \Lambda \supset \Lambda_+$  in  $\mathbb{C}$







# Extra twisted connected sums—Matching the tori

**Idea.** Represent adiabatic limits as rays in the hyperbolic plane

Lattices  $\Lambda_- \subset \Lambda \supset \Lambda_+$  in  $\mathbb{C}$

Hyperbolic plane  $\mathcal{H} = \mathbb{R} + i\mathbb{R}_{>}$





# Extra twisted connected sums—Matching the tori

**Idea.** Represent adiabatic limits as rays in the hyperbolic plane

Lattices  $\Lambda_- \subset \Lambda \supset \Lambda_+$  in  $\mathbb{C}$

Hyperbolic plane  $\mathcal{H} = \mathbb{R} + i\mathbb{R}_{>}$





# Extra twisted connected sums—Matching the tori



**Idea.** Represent adiabatic limits as rays in the hyperbolic plane

Lattices  $\Lambda_- \subset \Lambda \supset \Lambda_+$  in  $\mathbb{C}$

Hyperbolic plane  $\mathcal{H} = \mathbb{R} + i\mathbb{R}_{>}$



Conformal change of basis





# Extra twisted connected sums—Matching the tori



**Idea.** Represent adiabatic limits as rays in the hyperbolic plane

Lattices  $\Lambda_- \subset \Lambda \supset \Lambda_+$  in  $\mathbb{C}$

Hyperbolic plane  $\mathcal{H} = \mathbb{R} + i\mathbb{R}_{>}$



Conformal change of basis





# Extra twisted connected sums—Matching the tori

**Idea.** Represent adiabatic limits as rays in the hyperbolic plane



Lattices  $\Lambda_- \subset \Lambda \supset \Lambda_+$  in  $\mathbb{C}$

Hyperbolic plane  $\mathcal{H} = \mathbb{R} + i\mathbb{R}_{>}$



Conformal change of basis

Möbius transformation



# Extra twisted connected sums—Modular functions

The logarithm of the **Dedekind  $\eta$ -function** is given by

$$L(\tau) = \frac{\pi i \tau}{12} - \sum_{n=1}^{\infty} \sum_{d|n} d^{-1} e^{2\pi i n \tau}$$

## Theorem (G-Nordström-Zagier)

*There exists a constant  $c_{k_{\pm}, \varepsilon_{\pm}} \in \mathbb{Q}$  such that*

$$\begin{aligned} & 3\eta(B_{M_{\pm}}; L_{\pm}) - 24\eta(D_{M_{\pm}}; L_{\pm}) \\ &= \bar{\nu}(M_{\pm}) + \frac{144}{\pi} \left( iL\left(\frac{s_{\pm}i + \varepsilon_{\pm}}{k_{\pm}}\right) - iL\left(\frac{s_{\pm}i - \varepsilon_{\pm}}{k_{\pm}}\right) + c_{k_{\pm}, \varepsilon_{\pm}} \right) \end{aligned}$$

# Extra twisted connected sums—Modular functions

The logarithm of the **Dedekind  $\eta$ -function** is given by

$$L(\tau) = \frac{\pi i \tau}{12} - \sum_{n=1}^{\infty} \sum_{d|n} d^{-1} e^{2\pi i n \tau}$$

## Theorem (G-Nordström-Zagier)

*There exists a constant  $c_{k_{\pm}, \varepsilon_{\pm}} \in \mathbb{Q}$  such that*

$$\begin{aligned} & 3\eta(B_{M_{\pm}}; L_{\pm}) - 24\eta(D_{M_{\pm}}; L_{\pm}) \\ &= \bar{\nu}(M_{\pm}) + \frac{144}{\pi} \left( iL\left(\frac{s_{\pm}i + \varepsilon_{\pm}}{k_{\pm}}\right) - iL\left(\frac{s_{\pm}i - \varepsilon_{\pm}}{k_{\pm}}\right) + c_{k_{\pm}, \varepsilon_{\pm}} \right) \end{aligned}$$

Compute the variational term using the functional equations

$$L(\tau + 1) = \frac{\pi i}{12} + L(\tau) \quad \text{and} \quad L\left(-\frac{1}{\tau}\right) = \frac{1}{2} \log \frac{\tau}{i} + L(\tau)$$

# Extra twisted connected sums—Hyperbolic geometry

The exterior differential of the  $\eta$ -form  $\tilde{\eta}(\mathbb{A})$  is

$$d\tilde{\eta}(\mathbb{A}) = -\frac{1}{4\pi} dA_{\text{hyp}}$$




# Extra twisted connected sums—Hyperbolic geometry

The exterior differential of the  $\eta$ -form  $\tilde{\eta}(\mathbb{A})$  is

$$d\tilde{\eta}(\mathbb{A}) = -\frac{1}{4\pi} dA_{\text{hyp}}$$

Integrate  $288\tilde{\eta}(\mathbb{A})$  over a contour in  $\mathcal{H}$  of the form




# Extra twisted connected sums—Hyperbolic geometry

The exterior differential of the  $\eta$ -form  $\tilde{\eta}(\mathbb{A})$  is

$$d\tilde{\eta}(\mathbb{A}) = -\frac{1}{4\pi} dA_{\text{hyp}}$$

Integrate  $288\tilde{\eta}(\mathbb{A})$  over a contour in  $\mathcal{H}$  of the form




The bottom arcs do not contribute for symmetry reasons

# Extra twisted connected sums—Hyperbolic geometry

The exterior differential of the  $\eta$ -form  $\tilde{\eta}(\mathbb{A})$  is

$$d\tilde{\eta}(\mathbb{A}) = -\frac{1}{4\pi} dA_{\text{hyp}}$$

Integrate  $288\tilde{\eta}(\mathbb{A})$  over a contour in  $\mathcal{H}$  of the form



The bottom arcs do not contribute for symmetry reasons

Compute the variational term using [Gauß-Bonnet](#)

The corner angle cancels the irrational contribution from  $-72\frac{\rho}{\pi}$

There are additional contributions from the cusps

# Extra twisted connected sums—Results and questions

**Example.** The example with  $\cos \vartheta = \frac{1}{\sqrt{6}}$  has  $\bar{\nu}(M, g) = -65$   
In particular,  $3 \nmid \bar{\nu}(M, g)$ , so it is not  $G_2$ -nullbordant

# Extra twisted connected sums—Results and questions

**Example.** The example with  $\cos \vartheta = \frac{1}{\sqrt{6}}$  has  $\bar{\nu}(M, g) = -65$   
In particular,  $3 \nmid \bar{\nu}(M, g)$ , so it is not  $G_2$ -nullbordant

## Conjecture

*All values in  $\mathbb{Z}/48$  occur as  $\nu$ -invariants of  $G_2$ -holonomy metrics*

# Extra twisted connected sums—Results and questions

**Example.** The example with  $\cos \vartheta = \frac{1}{\sqrt{6}}$  has  $\bar{\nu}(M, g) = -65$   
In particular,  $3 \nmid \bar{\nu}(M, g)$ , so it is not  $G_2$ -nullbordant

## Conjecture

*All values in  $\mathbb{Z}/48$  occur as  $\nu$ -invariants of  $G_2$ -holonomy metrics*

## Questions

- ▶ How many different  $G_2$ -metrics exist on one 7-manifold?
- ▶ Are different  $G_2$ -metrics on a fixed 7-manifold  $G_2$ -bordant?

# Extra twisted connected sums—Results and questions

**Example.** The example with  $\cos \vartheta = \frac{1}{\sqrt{6}}$  has  $\bar{\nu}(M, g) = -65$   
In particular,  $3 \nmid \bar{\nu}(M, g)$ , so it is not  $G_2$ -nullbordant

## Conjecture

*All values in  $\mathbb{Z}/48$  occur as  $\nu$ -invariants of  $G_2$ -holonomy metrics*

## Questions

- ▶ How many different  $G_2$ -metrics exist on one 7-manifold?
- ▶ Are different  $G_2$ -metrics on a fixed 7-manifold  $G_2$ -bordant?

## Construct more examples

- ▶ Find more asymptotically cylindrical Calabi-Yau manifolds
- ▶ Understand their moduli space, make the K3 surfaces match
- ▶ Consider other constructions

Thanks for your attention!