


Preliminaries

1) Convolution: consider two functions, f & g .



Their convolution is defined as

$$f * g(x) = \int_{-\infty}^{\infty} f(x') g(x-x') dx'$$

The convolution of f with g can be interpreted as a "blurring" of f with g . To see this, use the Riemann sum interpretation of the integral:

$$x' \rightarrow x_m = m \Delta x, \quad \text{for } \Delta x \rightarrow 0.$$

$$f * g = \lim_{\Delta x \rightarrow 0} \sum_m \frac{f(x_m) g(x-x_m) \Delta x}{\Delta x}$$

That is, we take each piece of f :



and "blur" each piece with a displaced version of g :




Notice that the convolution is commutative, i.e.

$$f * g(x) = \int_{-\infty}^{\infty} f(x') g(x-x') dx' = \int_{\infty}^{-\infty} g(x'') f(x-x'') dx'' = \int_{-\infty}^{\infty} g(x'') f(x-x'') dx'' = g * f(x).$$

$x'' = x - x'$, $dx' = -dx''$

Exercise:

1) Let $f_1(x) = \text{rect}(x) =$  $= \begin{cases} 1, & |x| \leq 1/2 \\ 0, & |x| > 1/2 \end{cases}$
find $f_1 * f_1$


2) Let $f_2(x) = \frac{e^{-\pi(x/a)^2}}{a}$
find $f_2 * f_2$

3) (Only for those who like maths!)
find $f_1 * f_2$


Hint: $\text{erf}(\tau) = \frac{2}{\sqrt{\pi}} \int_0^{\tau} e^{-t^2} dt$

2) Delta function (Dirac)

$$\delta(x) = \begin{cases} \infty & x=0 \\ 0 & x \neq 0 \end{cases} \quad \text{such that} \quad \int_{-\infty}^{\infty} \delta(x) dx = 1.$$




We can build $\delta(x)$ from a function $g(x)$ (say, a Gaussian or a rectangle function) of unit area:



$$\int_{-\infty}^{\infty} g(x) dx = 1.$$

Note that $\frac{1}{\Delta} g\left(\frac{x}{\Delta}\right)$, for $0 < \Delta < 1$, also has unit area:



$$\int_{-\infty}^{\infty} \frac{g\left(\frac{x}{\Delta}\right)}{\Delta} dx = 1$$

↑ this is thinner and taller, but with the same area. Then, we can build $\delta(x)$ as

$$\delta(x) = \lim_{\Delta \rightarrow 0} \frac{1}{\Delta} g\left(\frac{x}{\Delta}\right)$$

Properties:

- Units. since $\int \delta(x) dx$ has no units, δ has units of $\frac{1}{x}$.

- Note that, since $\delta(x-x_0)$ is zero except at $x=x_0$, then $f(x)\delta(x-x_0) = f(x_0)\delta(x-x_0)$ for any (well-behaved) $f(x)$. Therefore

$$\int f(x)\delta(x-x_0)dx = f(x_0)\int\delta(x-x_0)dx = f(x_0)$$

This is ^{the} so-called "sifting property" of the delta function.

Note then that

$$f * \delta = \int f(x')\delta(x-x')dx' = f(x)$$

so δ is the "unity" element for convolutions.

Finally let us show that we can write

$$\delta(x) = \int_{-\infty}^{\infty} e^{i2\pi\nu x} d\nu$$

to show this, we insert 1 in the integrand in the form

$$1 = \lim_{a \rightarrow 0} e^{-\pi a \nu^2} \cdot e^{-\pi a(\nu^2 - 2i x \nu / a)}$$

so

$$\int_{-\infty}^{\infty} e^{i2\pi\nu x} d\nu = \lim_{a \rightarrow 0} \int_{-\infty}^{\infty} \overbrace{e^{-\pi a \nu^2} e^{-\pi a(\nu^2 - 2i x \nu / a)}} e^{i2\pi\nu x} d\nu$$

but

$$v^2 - 2i\frac{x}{a}v = \left(v - i\frac{x}{a}\right)^2 + \frac{x^2}{a^2}, \text{ so}$$


$$\begin{aligned} \int_{-\infty}^{\infty} e^{i2\pi vx} dv &= \lim_{a \rightarrow 0} \int e^{-\pi a \left(v - i\frac{x}{a}\right)^2} e^{-\frac{\pi x^2}{a}} dv \\ &= \lim_{a \rightarrow 0} e^{-\frac{\pi x^2}{a}} \underbrace{\int e^{-\pi a v'^2} dv'}_{\frac{1}{\sqrt{a}}} = \lim_{a \rightarrow 0} \frac{e^{-\frac{\pi x^2}{a}}}{\sqrt{a}} \end{aligned}$$

let $a = \Delta^2$, so

$$\int_{-\infty}^{\infty} e^{i2\pi vx} dv = \lim_{\Delta \rightarrow 0} \frac{e^{-\pi \left(\frac{x}{\Delta}\right)^2}}{\Delta} = \delta(x)$$

Fourier Theory

Sines & cosines



Any (well-behaved) function can be expressed as a continuous superposition of sines and cosines with different amplitudes and periods (Δ).

It is more convenient, though, to use imaginary exponentials. Recall

$$e^{\pm i\theta} = \cos \theta \pm i \sin \theta$$

so, instead of $\cos \frac{2\pi x}{\Delta}$ and $\sin \frac{2\pi x}{\Delta}$, we use:

$$e^{i2\pi \nu x}, \text{ with } \nu = \pm \frac{1}{\Delta}$$

The Fourier theorem then states that $f(x)$ can be written as

$$f(x) = \int_{-\infty}^{\infty} \tilde{f}(\nu) e^{i2\pi \nu x} d\nu$$

where $\tilde{f}(\nu)$, known as the Fourier transform of $f(x)$, is the amplitude of the corresponding oscillation.

How do we find $\tilde{F}(v)$? Consider the integral

$$\begin{aligned} \int_{-\infty}^{\infty} f(x) e^{-i2\pi v x} dx &= \int_{-\infty}^{\infty} \tilde{F}(v') e^{i2\pi v' x} dv' e^{-i2\pi v x} dx \\ &\quad \uparrow \text{Substitute as } \int_{-\infty}^{\infty} \tilde{F}(v') e^{i2\pi v' x} dv' \\ &= \int_{-\infty}^{\infty} \tilde{F}(v') \underbrace{\int_{-\infty}^{\infty} e^{i2\pi(v'-v)x} dx}_{\delta(v'-v)} dv' = \int_{-\infty}^{\infty} \tilde{F}(v') \delta(v'-v) dv' \\ &= \tilde{F}(v), \end{aligned}$$

so

$$\tilde{F}(v) = \int_{-\infty}^{\infty} f(x) e^{-i2\pi v x} dx$$

So in summary

$$\begin{aligned} \text{Fourier Transformation } \tilde{F}(v) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi v x} dx \\ \text{Inverse Fourier Transformation } f(x) &= \int_{-\infty}^{\infty} \tilde{F}(v) e^{i2\pi v x} dv \end{aligned}$$

In what follows we use the notation:

$$\begin{aligned} \tilde{F}(v) &= \hat{\mathcal{F}}_{x \rightarrow v} f(x) \\ f(x) &= \hat{\mathcal{F}}_{v \rightarrow x} \tilde{F}(v) \end{aligned}$$

Properties

• Parseval-Plancherel theorem

In many physical applications, $|f(x)|^2$ is a physically significant (and observable) quantity, and the integral of this quantity corresponds to, for example, the total power or energy. Note that

$$\begin{aligned}\int_{-\infty}^{\infty} |f(x)|^2 dx &= \int_{-\infty}^{\infty} f^*(x) \underbrace{f(x)}_{=\int_{-\infty}^{\infty} \tilde{f}(v) e^{i2\pi vx} dv} dx \\ &= \int_{-\infty}^{\infty} f^*(x) \int_{-\infty}^{\infty} \tilde{f}(v) e^{i2\pi vx} dv dx = \int_{-\infty}^{\infty} \tilde{f}(v) \int_{-\infty}^{\infty} f^*(x) e^{i2\pi vx} dx dv \\ &= \int_{-\infty}^{\infty} \tilde{f}(v) \underbrace{\left[\int_{-\infty}^{\infty} f(x) e^{-i2\pi vx} dx \right]^*}_{\tilde{f}(v)} dv = \int_{-\infty}^{\infty} \tilde{f}(v) \tilde{f}^*(v) dv = \int_{-\infty}^{\infty} |\tilde{f}(v)|^2 dv\end{aligned}$$

$$\boxed{\int_{-\infty}^{\infty} |f(x)|^2 dx = \int_{-\infty}^{\infty} |\tilde{f}(v)|^2 dv}$$

• Shift-phase

Consider the FT of a shifted function

$$\begin{aligned}\hat{f}_{x \rightarrow v} f(x-x_0) &= \int_{-\infty}^{\infty} \underbrace{f(x-x_0)}_{x'=x-x_0 \rightarrow x=x'+x_0, dx=dx'} e^{-i2\pi xv} dx \\ &= \int_{-\infty}^{\infty} f(x') e^{-i2\pi(x'+x_0)v} dx' = \int_{-\infty}^{\infty} f(x') e^{-i2\pi x'v} dx' e^{-i2\pi x_0 v} \\ &= \tilde{f}(v) e^{-i2\pi x_0 v}\end{aligned}$$

therefore

$$\hat{\mathcal{F}}_{x \rightarrow \nu} f(x-x_0) = \tilde{f}(\nu) e^{-i2\pi x_0 \nu} = \left[\hat{\mathcal{F}}_{x \rightarrow \nu} f(x) \right] e^{-i2\pi x_0 \nu}$$

which implies

$$\hat{\mathcal{F}}_{\nu \rightarrow x}^{-1} \left[\tilde{f}(\nu) e^{-i2\pi x_0 \nu} \right] = f(x-x_0)$$

Analogously, multiplying $f(x)$ by a linear phase function leads to the shift of the Fourier transform

$$\begin{aligned} \hat{\mathcal{F}}_{x \rightarrow \nu} \left[f(x) e^{i2\pi x \nu_0} \right] &= \int_{-\infty}^{\infty} f(x) e^{i2\pi x \nu_0} e^{-i2\pi x \nu} dx \\ &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi(\nu-\nu_0)x} dx = \tilde{f}(\nu-\nu_0) \end{aligned}$$

and therefore

$$\hat{\mathcal{F}}_{\nu \rightarrow x}^{-1} \tilde{f}(\nu-\nu_0) = f(x) e^{i2\pi \nu_0 x}$$

• Scaling

Consider the FT of $f\left(\frac{x}{a}\right)$

$$\begin{aligned} \hat{\mathcal{F}}_{x \rightarrow \nu} f\left(\frac{x}{a}\right) &= \int_{-\infty}^{\infty} f\left(\frac{x}{a}\right) e^{-i2\pi x \nu} dx \\ &= \begin{cases} a \int_{-\infty}^{\infty} f(x') e^{-i2\pi a x' \nu} dx', & a > 0 \\ a \int_{\infty}^{-\infty} f(x') e^{-i2\pi a x' \nu} dx', & a < 0 \end{cases} \end{aligned}$$

$$= \underbrace{\text{sgn}(a)}_{|a|} a \int_{-\infty}^{\infty} f(x') e^{-i2\pi x' (a\nu)} dx' = |a| \tilde{f}(a\nu)$$

• Derivative

$$\hat{\mathcal{F}}_{x \rightarrow \nu} f'(x) = \int_{-\infty}^{\infty} \underbrace{f'(x)}_u \underbrace{e^{-i2\pi x \nu}}_v dx = \int_{-\infty}^{\infty} u dv$$

Integrate by parts $dv = f' dx$ $u = e^{-i2\pi x \nu}$

$$= uv \Big|_{-\infty}^{\infty} - \int_{-\infty}^{\infty} v du = \underbrace{f(x) e^{-i2\pi x \nu}}_{v=f} \Big|_{-\infty}^{\infty} + i2\pi \nu \int_{-\infty}^{\infty} f(x) e^{-i2\pi x \nu} dx$$

$du = -i2\pi \nu e^{-i2\pi x \nu}$ $u = e^{-i2\pi x \nu}$

assume $f(\pm\infty) = 0$

$$= i2\pi \nu \tilde{f}(\nu)$$

More generally: $\hat{\mathcal{F}}_{x \rightarrow \nu} f^{(n)}(x) = (i2\pi \nu)^n \tilde{f}(\nu)$

Similarly

$$\hat{\mathcal{F}}_{x \rightarrow \nu} [x^n f(x)] = \int_{-\infty}^{\infty} f(x) \underbrace{x^n e^{-i2\pi x \nu}}_{\left(\frac{1}{i2\pi}\right)^n \frac{d^n}{d\nu^n} e^{-i2\pi x \nu}} dx$$

$$= \left(\frac{1}{i2\pi}\right)^n \frac{d^n}{d\nu^n} \int_{-\infty}^{\infty} f(x) e^{-i2\pi x \nu} dx = \frac{\tilde{f}^{(n)}(\nu)}{(i2\pi)^n}$$

• Convolution/product

$$\hat{\mathcal{F}}_{x \rightarrow \nu} [f * g] = \int_{-\infty}^{\infty} \left[\int_{-\infty}^{\infty} f(x) g(x-x') dx' \right] e^{-i2\pi x \nu} dx$$

$$= \int_{-\infty}^{\infty} f(x') \int_{-\infty}^{\infty} \underbrace{g(x-x') e^{-i2\pi x \nu}}_{\text{From shift/phase: } \tilde{g}(\nu) e^{-i2\pi x' \nu}} dx dx' = \tilde{g}(\nu) \int_{-\infty}^{\infty} f(x') e^{-i2\pi x' \nu} dx'$$

$$= \tilde{g}(\nu) \tilde{f}(\nu) = \tilde{f}(\nu) \tilde{g}(\nu)$$

Similarly

$$\begin{aligned} \hat{F}_{x \rightarrow v} [f(x)g(x)] &= \int_{-\infty}^{\infty} f(x)g(x) e^{-i2\pi xv} dx \\ &= \int_{-\infty}^{\infty} \tilde{g}(v') \int_{-\infty}^{\infty} f(x) e^{-i2\pi x(v-v')} dx = \int_{-\infty}^{\infty} \tilde{g}(v') \tilde{F}(v-v') dv' \\ &= \tilde{F} * \tilde{g} \end{aligned}$$

• Space-bandwidth product / uncertainty relation.

The average or "centroid" of $|f(x)|^2$ is defined as

$$\bar{x} = \frac{\int_{-\infty}^{\infty} x |f(x)|^2 dx}{\int_{-\infty}^{\infty} |f(x)|^2 dx}$$

and the rms spread is

$$\Delta x = \left[\frac{\int_{-\infty}^{\infty} (x - \bar{x})^2 |f(x)|^2 dx}{\int_{-\infty}^{\infty} |f(x)|^2 dx} \right]^{1/2}$$

Similarly

$$\bar{v} = \frac{\int_{-\infty}^{\infty} v |\tilde{F}(v)|^2 dv}{\int_{-\infty}^{\infty} |\tilde{F}(v)|^2 dv}, \quad \Delta v = \left[\frac{\int_{-\infty}^{\infty} (v - \bar{v})^2 |\tilde{F}(v)|^2 dv}{\int_{-\infty}^{\infty} |\tilde{F}(v)|^2 dv} \right]^{1/2}$$

It is now shown that

$$\Delta x \Delta v \geq \frac{1}{4\pi}$$

Proof.

Part a) Cauchy-Schwarz-Bunyakovski inequality

consider two functions g, h , then

$$\iint \underbrace{|g(x)h(y) - g(y)h(x)|^2}_{\text{this is always } \geq 0} dx dy \geq 0.$$

But we can write this as

$$\begin{aligned} & \iint \left[g^*(x)h^*(y) - g^*(y)h^*(x) \right] \left[g(x)h(y) - g(y)h(x) \right] dx dy \\ &= \iint \left[|g(x)|^2 |h(y)|^2 - g^*(x)h(x)h^*(y)g(y) \right. \\ & \quad \left. - g^*(y)h(y)h^*(x)g(x) + |g(y)|^2 |h(x)|^2 \right] dx dy \\ &= \int |g(x)|^2 dx \int |h(y)|^2 dy + \int |g(y)|^2 dy \int |h(x)|^2 dx \\ & \quad - \left[\int g^*(x)h(x) dx \int h^*(y)g(y) dy + \int g^*(y)h(y) dy \int h^*(x)g(x) dx \right] \end{aligned}$$

but x & y are now dummy variables, so we can write

$$= 2 \left[\int |g(x)|^2 dx \right] \left[\int |h(x)|^2 dx \right] - 2 \left| \int g^*(x)h(x) dx \right|^2.$$

and recall that all this ≥ 0 . Therefore

$$\int |g(x)|^2 dx \int |h(x)|^2 dx \geq \left| \int g^*(x)h(x) dx \right|^2$$

Part b)

Let $g(x) = \frac{(x-\bar{x})f(x)}{\Phi^{1/2}}$, where

$$\Phi = \int_{-\infty}^{\infty} |f(x)|^2 dx$$

then

$$\int_{-\infty}^{\infty} |g(x)|^2 dx = \frac{\int_{-\infty}^{\infty} (x-\bar{x})^2 |f(x)|^2 dx}{\int_{-\infty}^{\infty} |f(x)|^2 dx} = \frac{\Delta x^2}{\cancel{\Phi}}$$

Now, $\int_{-\infty}^{\infty} |\tilde{h}(x)|^2 dx = \int_{-\infty}^{\infty} |\tilde{h}(v)|^2 dv$ (Parseval-Plancherel)

Let $\tilde{h}(v) = \frac{(v - \bar{v}) \tilde{f}(v)}{\Phi^{1/2}}$, so $\int_{-\infty}^{\infty} |\tilde{h}(x)|^2 dx = \Delta v^2$

Notice

$$\tilde{h}(v) = \frac{1}{\Phi^{1/2}} [v \tilde{f}(v) - \bar{v} \tilde{f}(v)]$$

← constant.

therefore

$$h(x) = \hat{F}_{v \rightarrow x}^{-1} \tilde{h}(v) = \frac{1}{\Phi^{1/2}} \left[\frac{1}{i2\pi} f'(x) - \bar{v} f(x) \right]$$

Therefore

$$\int_{-\infty}^{\infty} g^*(x) h(x) dx = \frac{1}{\Phi} \int_{-\infty}^{\infty} (x - \bar{x}) f^*(x) \left[\frac{1}{i2\pi} f'(x) - \bar{v} f(x) \right] dx$$

$$= \frac{1}{i2\pi\Phi} \int_{-\infty}^{\infty} (x - \bar{x}) f^*(x) f'(x) dx - \frac{\bar{v}}{\Phi} \int_{-\infty}^{\infty} (x - \bar{x}) |f(x)|^2 dx \quad (i)$$

integrate by parts:

$u = (x - \bar{x}) f^*$, $dv = f' dx$, $v = f$, $du = [f^* + (x - \bar{x}) f'^*] dx$

$$= \frac{1}{i2\pi\Phi} \left[(x - \bar{x}) f^*(x) f(x) \right]_{-\infty}^{\infty} - \frac{1}{i2\pi\Phi} \int_{-\infty}^{\infty} [f(x) + (x - \bar{x}) f'(x)]^* f(x) dx$$

assume this vanishes.

$$= - \frac{\int_{-\infty}^{\infty} |f(x)|^2 dx}{i2\pi\Phi} - \frac{1}{i2\pi\Phi} \left[\int_{-\infty}^{\infty} f^*(x) (x - \bar{x}) f'(x) dx \right]^*$$

$$- \frac{\bar{v}}{\Phi} \int_{-\infty}^{\infty} (x - \bar{x}) |f(x)|^2 dx$$

$$\int_{-\infty}^{\infty} g^*(x) h(x) dx = +\frac{i}{2\pi} + \left[\frac{1}{i2\pi\Phi} \int_{-\infty}^{\infty} (x-\bar{x}) f^*(x) f'(x) dx - \frac{\bar{v}}{\Phi} \int_{-\infty}^{\infty} (x-\bar{x}) |f(x)|^2 dx \right]^* \quad (ii)$$

Note that $\int_{-\infty}^{\infty} g^*(x) h(x) dx$ is given by either the expression in (i) or the one in (ii), therefore

also by their average:

$$\int_{-\infty}^{\infty} g^*(x) h(x) dx = \frac{1}{2} \left[\frac{1}{\Phi} \int_{-\infty}^{\infty} (x-\bar{x}) f^*(x) \left(\frac{f'(x)}{i2\pi} - \bar{v} f(x) \right) dx \right]$$

$$+ \frac{1}{2} \left[\frac{i}{2\pi} + \frac{1}{\Phi} \left(\int_{-\infty}^{\infty} (x-\bar{x}) f^*(x) \left(\frac{f'(x)}{i2\pi} - \bar{v} f(x) \right) dx \right)^* \right]$$

$$= \underbrace{\operatorname{Re} \left\{ \frac{1}{\Phi} \int_{-\infty}^{\infty} (x-\bar{x}) f^*(x) \left(\frac{f'(x)}{i2\pi} - \bar{v} f(x) \right) dx \right\}}_{\text{call this } \Delta_{xv}} + \frac{i}{4\pi}$$

$$= \underbrace{\Delta_{xv}}_{\substack{\uparrow \\ \text{Real}}} + \frac{i}{4\pi}$$

Therefore:

$$\left| \int_{-\infty}^{\infty} g^*(x) h(x) dx \right|^2 = \left(\Delta_{xv} - \frac{i}{4\pi} \right) \left(\Delta_{xv} + \frac{i}{4\pi} \right) = \Delta_{xv}^2 + \frac{1}{(4\pi)^2}$$

so $\int_{-\infty}^{\infty} |g(x)|^2 dx \int_{-\infty}^{\infty} |h(x)|^2 dx \geq \left| \int_{-\infty}^{\infty} g^*(x) h(x) dx \right|^2$ gives

$$\Delta_x^2 \Delta_v^2 \geq \Delta_{xv}^2 + \frac{1}{(4\pi)^2} \geq \frac{1}{(4\pi)^2} \quad \text{so } \boxed{\Delta_x \Delta_v \geq \frac{1}{4\pi}}$$

• Complex conjugate

$$\begin{aligned}\hat{\mathcal{F}}_{x \rightarrow \nu} [f^*(x)] &= \int_{-\infty}^{\infty} f^*(x) e^{-i2\pi x \nu} dx \\ &= \left[\int_{-\infty}^{\infty} f(x) e^{-i2\pi(-\nu)x} dx \right]^* = \tilde{f}^*(-\nu)\end{aligned}$$

Note then that, if f is real

$$f(x) = f^*(x) \Rightarrow \hat{f}(\nu) = \tilde{f}^*(-\nu)$$

$$\underbrace{\operatorname{Re} \hat{f}(\nu) = \operatorname{Re} \hat{f}(-\nu)}$$

$$\underbrace{\operatorname{Im} \tilde{f}(\nu) = -\operatorname{Im} \tilde{f}(-\nu)}$$

The real part of \tilde{f} is even

The imaginary part of \tilde{f} is odd.

Exercise:

$$\hat{\mathcal{F}}_{x \rightarrow \nu} [|f(x)|^2] =$$

Summary

1D Fourier transform

$$\begin{aligned}\tilde{F}(\nu) &= \int_{-\infty}^{\infty} f(x) e^{-i2\pi x\nu} dx \\ f(x) &= \int_{-\infty}^{\infty} \tilde{F}(\nu) e^{i2\pi x\nu} d\nu\end{aligned}$$

Properties

- Parseval-Plancherel
$$\int_{-\infty}^{\infty} f^*(x) g(x) dx = \int_{-\infty}^{\infty} \tilde{F}^*(\nu) \tilde{g}(\nu) d\nu$$
$$\int_{-\infty}^{\infty} |f(x)|^2 dx = \int_{-\infty}^{\infty} |\tilde{F}(\nu)|^2 d\nu$$
- Shift-Phase
$$\hat{\mathcal{F}}_{x \rightarrow \nu} f(x-x_0) = \tilde{F}(\nu) e^{-i2\pi x_0 \nu}$$
$$\hat{\mathcal{F}}_{x \rightarrow \nu} [f(x) e^{i2\pi \nu_0 x}] = \tilde{F}(\nu - \nu_0)$$
- Scaling
$$\hat{\mathcal{F}}_{x \rightarrow \nu} f\left(\frac{x}{a}\right) = |a| \tilde{F}(a\nu) \quad (a \text{ real, } \neq 0)$$
- Derivative
$$\hat{\mathcal{F}}_{x \rightarrow \nu} f^{(n)}(x) = (i2\pi\nu)^n \tilde{F}(\nu)$$
$$\hat{\mathcal{F}}_{x \rightarrow \nu} [x^n f(x)] = \frac{\tilde{F}^{(n)}(\nu)}{(-i2\pi)^n}$$
- Convolution/product
$$\hat{\mathcal{F}}_{x \rightarrow \nu} [f * g] = \tilde{F}(\nu) \tilde{g}(\nu)$$
$$\hat{\mathcal{F}}_{x \rightarrow \nu} [f(x)g(x)] = \tilde{F} * \tilde{g}$$
- Space-bandwidth product / uncertainty $\Delta x \Delta \nu \geq \frac{1}{4\pi}$
- Complex conjugate
$$\hat{\mathcal{F}}_{x \rightarrow \nu} [f^*(x)] = \tilde{F}^*(-\nu).$$

Exercises. Calculate the FT of:

1) $\delta(x)$

2) $\delta(x-x_0)$

3) $\text{rect}(x)$

4) $\text{rect}(x) * \text{rect}(x)$

5) $c \text{rect}\left(\frac{x-a}{b}\right)$

6) $e^{-\pi x^2}$

7) $x e^{-\pi x^2}$

2 Dimensions

$$\underline{x} = (x, y), \quad \underline{v} = (v_x, v_y)$$

Convolution

$$f * g = \iint_{-\infty}^{\infty} f(\underline{x}') g(\underline{x} - \underline{x}') d\underline{x}'$$

Delta function $\delta(\underline{x})$

$$\iint_{-\infty}^{\infty} \delta(\underline{x}) d\underline{x} d\underline{y} = 1 \quad \text{units of } \frac{1}{x^2}$$

sifting: $\iint_{-\infty}^{\infty} f(\underline{x}) \delta(\underline{x} - \underline{x}_0) d\underline{x} d\underline{y} = f(\underline{x}_0)$

Fourier transform

$$\tilde{f}(\underline{v}) = \iint_{-\infty}^{\infty} f(\underline{x}) e^{-i2\pi \underline{x} \cdot \underline{v}} d\underline{x} d\underline{y}$$

$$f(\underline{x}) = \iint_{-\infty}^{\infty} \tilde{f}(\underline{v}) e^{i2\pi \underline{x} \cdot \underline{v}} d\underline{v}_x d\underline{v}_y$$

Properties

• Parseval-Plancherel $\iint_{-\infty}^{\infty} f^*(\underline{x}) g(\underline{x}) d\underline{x} d\underline{y} = \iint_{-\infty}^{\infty} \tilde{f}^*(\underline{v}) \tilde{g}(\underline{v}) d\underline{v}_x d\underline{v}_y$

• Shift-Phase $\hat{\mathcal{F}}_{\underline{x} \rightarrow \underline{v}} f(\underline{x} - \underline{x}_0) = \tilde{f}(\underline{v}) e^{-i2\pi \underline{x}_0 \cdot \underline{v}}$

$$\hat{\mathcal{F}}_{\underline{x} \rightarrow \underline{v}} [f(\underline{x}) e^{i2\pi \underline{v}_0 \cdot \underline{x}}] = \tilde{f}(\underline{v} - \underline{v}_0)$$

• Scaling $\hat{\mathcal{F}}_{\underline{x} \rightarrow \underline{v}} f(\underline{x}/a) = a^2 \tilde{f}(a \underline{v})$

• Derivative $\hat{\mathcal{F}}_{\underline{x} \rightarrow \underline{v}} [\nabla_{\underline{x}} f(\underline{x})] = i2\pi \underline{v} \tilde{f}(\underline{v})$

$$\hat{\mathcal{F}}_{\underline{x} \rightarrow \underline{v}} [\underline{x} f(\underline{x})] = \frac{1}{-i2\pi} \nabla_{\underline{v}} \tilde{f}(\underline{v})$$

• Convolution $\hat{\mathcal{F}}_{\underline{x} \rightarrow \underline{v}} [f * g] = \tilde{f}(\underline{v}) \tilde{g}(\underline{v}), \quad \hat{\mathcal{F}}_{\underline{x} \rightarrow \underline{v}} [f(\underline{x}) g(\underline{x})] = \tilde{f} * \tilde{g}$

• Uncertainty $\Delta p \Delta v \geq \frac{1}{2\pi}$

2D Fourier transform in polar coordinates:

$$\underline{x} = (\rho \cos \theta, \rho \sin \theta) \quad , \quad \underline{v} = (v \cos \phi, v \sin \phi)$$

$$\tilde{F}(\underline{v}) = \int_0^{\infty} \int_0^{2\pi} f(\underline{x}) e^{-i2\pi \rho v \cos(\theta - \phi)} \rho \, d\theta \, d\rho$$

If $f(\underline{x})$ depends only on ρ , i.e. has rotational symmetry: $f(\underline{x}) = f_{\rho}(\rho)$

$$\tilde{F}(\underline{v}) = \int_0^{\infty} f_{\rho}(\rho) \rho \underbrace{\int_0^{2\pi} e^{-i2\pi \rho v \cos(\theta - \phi)} \, d\theta}_{2\pi J_0(2\pi \rho v)} \, d\rho$$

$2\pi J_0(2\pi \rho v)$, independent of ϕ

so $\tilde{F}(\underline{v}) = \tilde{F}_v(v)$ also has rotational symmetry.

$$\text{Hankel Transf. } \tilde{F}_v(v) = 2\pi \int_0^{\infty} f_{\rho}(\rho) J_0(2\pi \rho v) \rho \, d\rho$$

$$\text{Inverse HT } f_{\rho}(\rho) = 2\pi \int_0^{\infty} \tilde{F}_v(v) J_0(2\pi \rho v) v \, dv$$

In this case

$$\Delta_{\rho} = \left[\frac{\int_0^{\infty} |f_{\rho}(\rho)|^2 \rho^2 \, d\rho}{\int_0^{\infty} |f_{\rho}(\rho)|^2 \rho \, d\rho} \right]^{1/2}$$

$$\Delta_v = \left[\frac{\int_0^{\infty} |\tilde{F}_v(v)|^2 v^2 \, dv}{\int_0^{\infty} |\tilde{F}_v(v)|^2 v \, dv} \right]^{1/2}$$

$$\Delta_{\rho} \Delta_v \geq \frac{1}{2\pi}$$

Exercises:

• Calculate the Hankel transform of

$$1) f_p(\rho) = \delta(\rho - a)$$

$$2) f_p(\rho) = \begin{cases} 1, & \rho \leq a \\ 0, & \rho > a \end{cases}$$

$$3) f_p(\rho) = \begin{cases} 1 - \frac{\rho^2}{a^2}, & \rho \leq a \\ 0, & \rho > a \end{cases}$$

Formulas you might need

$$\int_0^u u' J_0(u') du' = u J_1(u)$$

$$\int_0^u u'^3 J_0(u') du' = 2u^2 J_2(u) - u^3 J_3(u)$$

$$J_{n+1} + J_{n-1} = 2n \frac{J_n}{u}$$

• Calculate the convolution of 2) with itself.

What is its Fourier transform?

Discrete Fourier transform (DFT)


Instead of $f(x)$ we have $f_n, n=0, \dots, N-1$

$$F_m = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} f_n e^{-i2\pi mn/N}$$

Discrete Fourier transform

Inverse: try:

$$f_{n'} = \frac{1}{\sqrt{N}} \sum_{m=0}^{N-1} F_m e^{i2\pi mn'/N}$$

$$= \frac{1}{N} \sum_{n=0}^{N-1} f_n \underbrace{\sum_{m=0}^{N-1} e^{i2\pi(n'-n)m/N}}_{N \delta_{n'-n}}$$


So:

$$f_n = \frac{1}{\sqrt{N}} \sum_{m=0}^{N-1} F_m e^{i2\pi mn/N}$$

Inverse Discrete Fourier transform

Approximating FT with DFT

(Notice that the sums can be shifted,) if we define

$$f_{n-N} = f_n$$

$$F_m = \frac{1}{\sqrt{N}} \sum_{n=-\lfloor \frac{N-1}{2} \rfloor}^{\lfloor \frac{N-1}{2} \rfloor} f_n e^{-i2\pi mn/N}$$

Let f_n be a sampling of $f(x)$:

$$f_n = f(n \Delta x)$$

$$F_m = \frac{1}{\sqrt{N}} \sum_{n=-\lfloor \frac{N-1}{2} \rfloor}^{\lfloor \frac{N-1}{2} \rfloor} f(n \Delta x) e^{-i2\pi mn/N}$$

For very large N , and small Δx ,
 can approximate the sum as an integral

$$F_m \approx \frac{1}{\sqrt{N}} \int_{-X_1}^{X_2} f(x) e^{-i2\pi m x} \frac{dx}{\Delta x}$$

where $n\Delta x \rightarrow x$

$$X_1 = \lfloor \frac{N-1}{2} \rfloor \Delta x, \quad X_2 = \lfloor \frac{N}{2} \rfloor \Delta x$$

Assume $\overset{\uparrow}{N} \overset{\uparrow}{\Delta x} = \text{big} \gg \text{width of } f(x)$.
big small note $X_1 \approx X_2 \approx \frac{N\Delta x}{2} = \text{big}$.

Then

$$F_m \approx \frac{1}{\sqrt{N} \Delta x} \int_{-\infty}^{\infty} f(x) e^{-i2\pi x \left(\frac{m}{N\Delta x}\right)} dx$$


$$= \frac{\tilde{f}\left(\frac{m}{N\Delta x}\right)}{\sqrt{N} \Delta x}$$

So the sampling distance in ν is $\frac{1}{N\Delta x} \approx \frac{1}{2X_2}$

where $2X_2$ is the width over which
 we're sampling $f(x)$.

Therefore:

- To increase resolution in $\tilde{f}(\nu) \longrightarrow$ must increase range in $f(x)$




- To increase range in $\tilde{f}(\nu)$ and avoid aliasing \longrightarrow must decrease sampling spacing in $f(x)$




Shifting the functions.

Notice that, if we sample:




we get


f_n



so we get



Similarly, once we get F_m , it will look like



To reconstruct $\tilde{f}(\nu)$ we must cut the second half and place it before the first. we also need to multiply by $\sqrt{N\Delta x}$.

Fast Fourier transform (FFT)

Notice that the, for each m , the DFT involves the sum of N terms. Since m runs from 0 to $N-1$, then N^2 must be performed. The time of computation can therefore be expected to be proportional to N^2 .

The FFT is an algorithm for performing the DFT whose time of computation is proportional to $N \log N$. While it can work for any N , its simplest form can be understood if $N = 2^M$ (so that $M = \log_2 N$):

$$F_m = \frac{1}{\sqrt{N}} \sum_{n=0}^{N-1} f_n e^{-i2\pi nm/N} = \frac{1}{\sqrt{N}} \left[\underbrace{\sum_{n'=0}^{\frac{N}{2}-1} f_{2n'} e^{-i2\pi(2n')m/N}}_{\text{terms with even } n} + \underbrace{\sum_{n'=0}^{\frac{N}{2}-1} f_{(2n'+1)} e^{-i2\pi(2n'+1)m/N}}_{\text{terms with odd } n} \right]$$

writes as $\frac{N}{2}$

$$= \frac{1}{\sqrt{2}} \left[\underbrace{\frac{1}{\sqrt{N/2}} \sum_{n'=0}^{\frac{N}{2}-1} f_{2n'} e^{-i2\pi n'm/(N/2)}}_{\text{DFT of size } N/2} + e^{-i2\pi m/N} \underbrace{\frac{1}{\sqrt{N/2}} \sum_{n'=0}^{\frac{N}{2}-1} f_{(2n'+1)} e^{-i2\pi n'm/(N/2)}}_{\text{DFT of size } N/2} \right]$$

Each of these two sums is itself a DFT of size $\frac{N}{2}$.

They can be joined.

$$F_m = \frac{1}{\sqrt{N}} \sum_{n'=0}^{\frac{N}{2}-1} \left(f_{2n'} + e^{-i2\pi m/N} f_{(2n'+1)} \right) e^{-i2\pi n'm/(N/2)}$$

The same separation can be done M times.

2D DF

$$F_{m_1, m_2} = \frac{1}{N} \sum_{n_1=0}^{N-1} \sum_{n_2=0}^{N-1} f_{n_1, n_2} e^{-i 2\pi \frac{(m_1 n_1 + m_2 n_2)}{N}}$$


$$f_{n_1, n_2} = \frac{1}{N} \sum_{m_1=0}^{N-1} \sum_{m_2=0}^{N-1} F_{m_1, m_2} e^{i 2\pi \frac{(m_1 n_1 + m_2 n_2)}{N}}$$

Using 2D DFT to approximate 2D FT:

if $f_{n_1, n_2} = f(n_1 \Delta x, n_2 \Delta x)$,

and $N \Delta x$ is bigger than width of f , then:

$$F_{m_1, m_2} \approx \frac{1}{N \Delta x^2} \tilde{f}\left(\frac{m_1}{N \Delta x}, \frac{m_2}{N \Delta x}\right)$$



Fast Fourier transform: time $\propto N^2 \log N$