



the
abdus salam
international centre for theoretical physics

**COURSE ON CLIMATE VARIABILITY
STUDIES IN THE OCEAN**

"Tracing & Modelling the Ocean Variability"
16 - 27 June 2003

301/1507-9

**Ocean Surface Temperatures During the
Last 150,000 Years-I**

Julian Sachs
MIT
Cambridge, MA
USA


Please note: These are preliminary notes intended for internal distribution only.

Ocean Surface Temperatures During the Last 150,000 Years-I

Julian Sachs

**Dept. of Earth, Atmospheric & Planetary Sciences
Massachusetts Institute of Technology
Cambridge, Massachusetts, USA**

"Mostly Sunny with a 10% Chance of Snow"




Kump et al. (1999)

**4.5 x 10⁹ Yr
of Climate
on Earth**


*Mostly
Sunny with
a 10%
Chance of
Snow*

Climate of the last Glacial Cycle




Climate of
the Last
150,000
Years
*A Complete
Glacial
Cycle*

• Climate archives in ice & sediment




• Alkenone paleothermometry


• Part I: Glacial temperatures in southern mid-latitudes




• Part II: Southern Ocean Expression of Massive Ice Discharge Events in the N. Atlantic



Climate Archives in Ice & Sediment




Greenland Ice Cores



Accomodations

www.noaa.gov






Antarctic Ice Cores



Processing Ice Cores



19 cm long section of GISP 2 ice core from 1855 m showing annual layer structure illuminated from below by a fiber optic source. Section contains 11 annual layers with summer layers (arrowed) sandwiched between darker winter layers.


Transport

Storage
(NICL, Denver, USA)

Dating


Determining Paleotemperature from Isotopes of Ice (H_2O)

Modern mean annual values of $\delta^{18}\text{O}$ and snowpack temperature from the Greenland Ice Sheet show an extremely close correspondence.



After Johnsen et al. (1988).

Influence of Continental Ice Volume on Oxygen Isotope Ratio of the Ocean



H₂¹⁸O ~ 1% Lower Vapor Pressure than H₂¹⁶O

$$\delta^{18}\text{O} = \left[\frac{{}^{18}\text{O}/{}^{16}\text{O}_{\text{smpl}}}{{}^{18}\text{O}/{}^{16}\text{O}_{\text{stdl}}} - 1 \right] \times 1000\%$$



Std = Std. Mean Ocean Water




JULEN



Coring the Seafloor




Rapidly-Accumulating Sediments
are Hard to Find!




⇒ Need > 15-30 cm/kyr!

High
Temporal
Resolution
Requires
Rapidly-
Accumulating
Sediments...

Rapidly-Accumulating Sediments Analyzed in this Study




Alkenone Paleothermometry



Coccoliths


Emiliania huxleyi

*Gephyrocapsa
oceanica*




Alkenones

Alkenone Temperature Relationship



$$Uk'_{37} = \frac{C_{37:2}}{C_{37:2} + C_{37:3}}$$

Prahl et al. (1988)




Global Core-
Top
Calibration
of Alkenone
Thermometer

60°S - 60°N


Müller et al. (1998)

Alkenone Identification by GC-MS (TN057-21, 479.5 cm)



Part I: The (unexpected) temperature history of southern mid-latitudes during the last glacial period

Unexpected SST History in Glacial SE Atlantic



Unexpected SST History of Glacial SE Atlantic


Criticisms

- (1) Sediment transport compromises fidelity of SST signal.
SST record in nearby non-drift core
- (2) Cannot extrapolate beyond local scale.
SST records in SW Pacific
Antarctic deuterium excess

Fidelity of the SST Proxy

**Do down-core changes in SST
reflect sediment transport or
surface temperature?**

Cape Basin Sediment Cores




Core	Latitude	Longitude	Water Depth (m)	Sed. Rate (cm/kyr)
ODP 1089	40°56.18' S	9°53.64' E	4621	16.2
TN057-21-PC2	41°08' S	7°49' E	4981	12.5
TN057-6-PC4	42°54.8' S	8°54' E	3751	3.4

Tucholke & Embley (1984)

•Cores from
Sediment Drifts

•Non-drift core




Synchronize
Drift
(TN057-21)
& Non-Drift
(TN057-6)
Cores


Sachs & Anderson (in press)

$\delta^{18}\text{O}$ data:
Hodell et al. (2000) &
Ninnemann et al. (1999)

SSTs at a Drift & Non-Drift Site




Climatological SST & Position of Cores



Levitus (1994)

Sediment Focusing and SST




Circum-hemispheric vs Regional Climate

**Over what spatial scale is the
Cape Basin SST record
representative?**

Glacial Surface Temperatures of the Southeast Atlantic Ocean

Julian P. Sachs,^{1*} Robert F. Anderson,² Scott J. Lehman³



SCIENCE VOL 293 14 SEPTEMBER 2001

- Justification for extrapolation to hemisphere-scale:
precipitation source T
records in AA ice
 $d = \delta D - 8 * \delta^{18}O$
 $1\text{‰} \sim 1^\circ\text{C} (\Delta T_{\text{source}})$

(Vimeux et al. 1999; Jouzel et al. 1982)


- Vostok precip: mid-latitude Indian & Pacific

(Koster et al. 1992; Delaygue et al. 2000)

- Dome C precip: mid-latitude Indian


(Stenni et al. 2001)

Moisture Sources for Central East Antarctic Ice Cores




Adapted from Stenni et al. (2001); Koster et al. (1992); Delaygue et al. (2000)

Cape Basin SST & Vostok ΔT_{source}




Southwest Pacific Cores

MD97-2121 40°S, 178°E, 2314 m
MD97-2120 46°S, 175°E, 1210 m




Glacial Climate of Southern Mid-latitudes




Sachs et al. (2001, in press & unpub.)
Vimeux et al. (2002)
Stenni et al. (subm.)
Imbrie et al. (1984)

Climate Forcing by Changes in Earth's Tilt?




Sachs et al. (2001, in press & unpub.)
Vimeux et al. (2002)
Stenni et al. (subm.)
Berger (1978)

The Ocean's Thermohaline Circulation Likely Varied in Concert with Changes in Earth's Tilt...




Rahmstorf (2002)




Boyle (1984)

Co-variation of Thermohaline Circulation & Southern Mid-latitude SST?



Cooling: More NADW
Warming: Less NADW

Rio Grande Rise
32°S 2925 m
Boyle (1984)




Models indicate S. Mid-latitudes Warm in Response to Diminished N. Atlantic Deep Water Production

Coupled GCM Evidence for a “Bipolar Seesaw”

Stocker (2002)

Low Obliquity




: 22°
Insolation: 234 W/m^2

S Insol.: 161 W/m^2
ient

$\uparrow \text{SST}_x$

High Obliquity




24.5°
 238 W/m^2

155 W/m^2

$\downarrow \text{SST}_x$


Obliquity
may also
influence S.
Ocean SST
by altering
meridional T
gradient

Poleward Shift of Westerlies in Response to Increased Eq-Pole T Gradient



J. Marshall (unpub.)
MITgcm

Cape Basin SST & Vostok CO₂



Part I: Conclusions

- Climate of S mid-latitudes differed from that of globe & N hemisphere during last glacial period
- Cooling 60-40 ka followed by warming 40-25 ka
- May have been associated with changes in Atlantic Thermohaline Circulation
- Antarctic air T poor proxy for much of S Hemisphere climate
- Additional & longer SST records needed to evaluate forcing mechanisms

Ocean Surface Temperatures During the Last 150,000 Years-II


Julian Sachs

**Dept. of Earth, Atmospheric & Planetary Sciences
Massachusetts Institute of Technology
Cambridge, Massachusetts, USA**


Part II: Southern Ocean
Expression of Massive Ice
Discharge Events in the North
Atlantic

“Abrupt Climate Change”

A change in climate that occurs more rapidly than a known forcing.





Climate of the last Glacial Cycle




Climate of the Last Glacial Cycle

Global Character of Abrupt Climate Change During Glacial Period







- Much of N. Hem. changed in concert
- Few detailed paleoclimate records from vast Southern Hemisphere oceans

Voelker et al. (2002) in
Rahmstorf (2002)




Two Types of Abrupt Climate Events in the Glacial North Atlantic: Dansgaard-Oeschger Events Heinrich Events






Typical Sand-Size ($> 150 \mu\text{m}$)
Fraction in NW Atlantic Core
Foraminifera





Sand-Size Fraction in
Heinrich Layer-2 in NW
Atlantic (670-672 cm)
Ice-Rafted Debris

Sediment Transport and Deposition Associated with Heinrich Events




Icebergs


www.noaa.gov



Northern Hemisphere Ice Sheets During the Last Glacial Period



Southern Hemisphere Ice Sheets During the Last Glacial Period





- Little additional land-based ice in S. Hemisphere during glacial period

Heinrich *Layers* in the North Atlantic



www.noaa.gov



Thickness of Heinrich Layers H-1 and H-2
from North Atlantic Cores Demonstrate
Source Areas and Diffusion of Ice-Rafted Debris
from the Laurentide Ice Sheet






Sites
Exhibiting
Heinrich Events
or Events
Synchronous
with Heinrich
Events




Peaks in *Pinus* (Pine) Pollen Data from Lake Tulane, Florida Correlate Well with Sedimentological Data from the North Atlantic for Heinrich Events 1 through 5



Global (?) Impact of Heinrich Events





Cacho et al. (1999)



Wang et al. (2001)
Schulz et al. (1998)
Arz et al. (1998)
Cacho et al. (1999)
Grimm et al. (1997)

3 Modes of Ocean Circulation During the Last Glacial Period?


Rahmstorf (2002)



Expression of H-events SE of New Zealand

- High algal productivity
- Warm SSTs


MD97-2120
46°S, 175°E
1210 m




$\delta^{18}\text{O}$: K. Pahnke (in prep.). HE Ages: van Kreveld et al. (2000) & Rashid et al. (in press)

Cause of Subantarctic SW Pacific Warmth During Heinrich Events

Coupled GCM Evidence for a “Bipolar Seesaw”



Asynchrony of Greenland & Antarctic Temperature




MD97-2120 SST with GISP2 & Byrd T

Cause of High Alkenone Concentrations During Heinrich Events


3 Factors Influencing Alkenone Concentrations in Sediments

- Dilution by other sedimentary components
²³⁰Th-normalized alkenone fluxes argue against this.
- Alkenone preservation
Co-variation of [alkenone] at 1250 m in subantarctic SW Pacific & 5000 m in subtropical SE Atlantic argue against this.
- Coccolith production
Most likely cause of [alkenone] maxima during H-events.

Circum-Hemispheric Productivity Events?




Unlike Most of the Global Ocean,
Macronutrients do not Limit Algal Productivity in the S. Ocean



...Instead, iron does
(Martin, 1990)

Levitus (1994)



Southern
Ocean
Primary
Production


9/97-8/98

<http://seawifs.gsfc.nasa.gov>

**While S. Ocean [Alkenone] Co-varies with Antarctic Air
Temp., Not Likely Causal...**


Evidence for Iron-Limitation of Algal Growth in Southern Ocean Water

S. Ocean 1° Production Limited by Fe: SOIREE-1






Boyd et al. (2000) *Nature*, Vol. 407: 695-701

S. Ocean 1° Production Limited by Fe: SOIREE-2




Boyd et al. (2000) *Nature*, Vol. 407: 695-701

In vitro Fe-Addition Experiment October, 1997 Subantarctic Water (F1)



Fe-Limitation of 1° Production in Subantarctic water SE of New Zealand




A massive sandstorm blowing off the northwest African desert has blanketed hundreds of thousands of square miles of the eastern Atlantic Ocean with a dense cloud of Saharan sand. The massive nature of this particular storm was first seen in this SeaWiFS image acquired on Saturday, 26 February 2000 when it reached over 1000 miles into the Atlantic. These storms and the rising warm air can lift dust 15,000 feet or so above the African deserts and then out across the Atlantic, many times reaching as far as the Caribbean where they often require the local weather services to issue air pollution alerts as was recently the case in San Juan, Puerto Rico. Recent studies by the U.S.G.S. (http://catberler.usgs.gov/african_dust/) have linked the decline of the coral reefs in the Caribbean to the increasing frequency and intensity of Saharan Dust events. Additionally, other studies suggest that Sahelian Dust may play a role in determining the frequency and intensity of hurricanes formed in the eastern Atlantic Ocean (<http://www.thirdworld.org/role.html>)

Provided by the SeaWiFS Project, NASA/GSFC and ORBIMAGE

<http://seawifs.gsfc.nasa.gov>

N. Atlantic Deep Water Supplies Fe to the S.Ocean




Fe* = Fe deficit
(relative to PO_4^{3-})

$$\text{Fe}^* = \text{Fe} - R_{\text{Fe}} \text{PO}_4$$
$$R_{\text{fe}} = \text{Fe}/\text{PO}_4$$

$$\text{Fe/C} = 4 \mu\text{mol/mol}$$
$$\text{C/P} = 117 \text{ mol/mol}$$
$$\text{Fe/P} = 0.47 \text{ mmol/mol}$$


Payal Parekh (2003)
Ph.D. Thesis, MIT-
WHOI.

Possible LGM Deepwater Circulation



“A shallower variety of North Atlantic deep water will have a more difficult time reaching high latitudes of the Southern Ocean where it can be recycled into the bottom water. . . . Instead, any high-¹³C water emerging beyond the tip of Africa might simply flow eastward into the Indian and Pacific basins, bypassing the Antarctic entirely.” (Imbrie et al., 1992)

**Irrespective of the style of N. Atl. Deep /
Intermediate Water, S. Ocean [Fe] Likely
increased w/ global dust flux during cold periods**



Part II: Conclusions

Massive Iceberg Discharge Events in N Atlantic (Heinrich Events) Were Likely Associated With:

- Large productivity increases in Southern Ocean
 - May have been caused by increased [Fe]
- Warming of subantarctic SW Pacific
- Additional & longer SST records needed to determine spatial extent & evaluate causal links

Acknowledgements

Discussions

Ed Boyle (MIT)

Bob Anderson (LDEO)

Payal Parekh (MIT-WHOI Jt. Prog)

Samples:

MD97-2120: R. Zahn, K. Pahnke (U. Cardiff, Wales)

MD97-2121: L. Carter, B. Manighetti (NIWA, NZ)

ODP (Bremen)

Lab Assistance:

Ying Chang (MIT)

Maria Shriver (MIT)

Bridgette Therriault (MIT)

Zach Gazak (MIT)