



**The Abdus Salam
International Centre for Theoretical Physics**



1939-22

**Joint ICTP-IAEA Workshop on Nuclear Structure and Decay Data:
Theory and Evaluation**

28 April - 9 May, 2008

ENSDF - Adopted Levels and Gammas


Coral M. BAGLIN
*Lawrence Berkeley National Laboratory
Nuclear Science Division
Univ. California
CA-94707 Berkeley
U.S.A.*

ENSDF – Adopted Levels and Gammas

Coral Baglin

Workshop on *Nuclear Structure and Decay Data: Theory and Evaluation*
ICTP, Trieste, 28 April – 9 May 2008

ENSDF Database Structure



Adopted Levels, Gammas

This dataset is the heart of any nuclide evaluation !

- It is the condensation of all the information in all the other datasets and provides the **best values** known at the time of the evaluation.
- It provides the information that goes into the summary database NUDAT.
- It may be the **only** dataset that some readers will ever look at.
- The source of all data appearing here must be made transparent to the reader and easily traceable.

General Information

Q values:

- Usually rounded values from latest mass table (presently 2003Au03).
- Add new $S(p)$, $Q(\alpha)$, *etc.*, (with keynumber) if available; compare with 2003Au03 value.
- Optional: Comment on uncertainties in 'SY' values; note newly-measured masses if very different from Audi's prediction.

General Comments:

e.g., Production/Identification, keynumber lists for major shell model calculations or isotope shift/hfs references (all optional).

Other Reactions:

Give reaction and keynumber if wanted for completeness, even though no data have been used and no reaction dataset has been created; *e.g.*, a continuum gamma study (optional).

Define XREF Symbols:

Every DSID in nuclide must be listed here, even if it won't be associated with any specific level.

167IR ADOPTED LEVELS

167IR C **Production:** 92MO(78KR,p2n) E=357, 384 MEV (1997DA07).

167IR C **Identification:** 1981HO10 unambiguously assign a new α group to 167IR

167IR2C by relating it to known transitions through a multi-dimensional

167IR3C analysis correlating parent energies, daughter energies, and the

167IR4C timing of events. The production reactions involved 58NI on

167IR5C molybdenum-tin targets and 107AG on vanadium-nickel targets

167IR C **For calculation of proton decay widths for 167IR GS and isomer see**

167IR2C **2000DA11.**

167IR Q 11760 SY-1070 6 6507 5 1995AU04,1997DA07

167IR CQ **|DS(n)=300 (1995AU04).**

167IR CQ **QA\$from measured EA=6351 5 (1997DA07) for GS to GS transition; 1995AU04**

167IR2CQ **give QA=6495 50, reflecting lack of information concerning daughter**

167IR3CQ **state at that time.**

167IR CQ **SP From measured EP=1064 6 (1997DA07) for GS to GS transition;**

167IR2CQ **SP=-1110 10 in 1995AU04.**

167IR **XA171AU A DECAY (1.02 MS)**

167IR **XB78KR(92MO,2NPG)**

167IR L 0 (1/2+) 35.2 MS 20

167IR2 L %A=48 6 (1997DA07)\$%P=32 4 (1997DA07)\$%EC+%B+=?

167IRX L **XREF=B**

167IR CL J comparison of calculated and measured partial lifetimes for

167IR 2CL p decay rule out $d\{-3/2\}$ and $h\{-11/2\}$ transitions, so 1997DA07 conclude

167IR 3CL that an L=0 p is emitted to the 0+ GS of 166OS.

167IR CL %A,%P From relative intensities of α and p decay from level,

Level & Gamma Properties - General

- Assignments are definite (no parens.) if based on ‘strong’ arguments but indefinite (in parens.) if justification includes a ‘weak’ argument; see Nuclear Data Sheets (NDS) introductory material for specific rules.
- Every nuclide must have at least 1 level.
- Document sources of all data (**dataset name**, not just keynumber).
- Comment on serious discrepancies.
- Specify whether ‘average’ is weighted or unweighted (use larger of internal & external uncertainties for weighted averages).
- Remember to round off so uncertainty <26.
- Remember that ‘level’ and ‘gamma’ data appear in different tables in NDS; unhelpful to say “Jpi for levels with γ to 8+ isomer are based on ...” (in level table) or “mult for γ ’s observed in low spin reactions is from ...” (in γ table).
- Do not include:
 - continuation G records that give CC, KC, *etc.*;
 - primary γ rays from n capture;
 - neutron capture state(s);
 - coincidence ‘C’ from col. 78 of G records.
 - unplaced γ rays listed in source datasets.

Level Properties

Level Energy:

- Use GTOL to calculate from adopted E_γ (in most cases).
- Include all discrete levels and giant resonances; identify analog resonances.
- Adopt minimum number of levels consistent with source datasets.

T1/2 (or Γ):

- Specify source, *e.g.*, “from $B(E2)^\uparrow$ in Coulomb excitation”, *etc.*
- Give bare-atom half-lives in comment (*e.g.*, “ $T_{1/2}(52\text{Fe}26^+) = \dots$ ”).
- Remember $\Gamma = \Gamma_\gamma + \Gamma_p + \dots$ for resonance, so note any assumptions made, such as ‘ $\Gamma = \Gamma_{\gamma_0} + \Gamma_{\gamma_1}$ ’ or ‘ $\Gamma = \Gamma_p$ ’.

Band Flag: (if relevant)

Give rotational band parameters in comment (if meaningful) from:

$$E_K(J) = E_0 + A(J(J+1) - K^2) + B(J(J+1) - K^2)^2 + (-)^{(J+K)}(J+K)!/(J-K)!(A_{2K} + B_{2K}(J(J+1) - K^2)).$$

Isospin: very important for low A nuclides !

Level Decay Branches: for g.s. and $T_{1/2} \geq 0.1$ s levels, include all modes that might reasonably be expected, even if not yet observed.


```

92RB  Q  8096          6  5098      10 10750      60 -6460      40      2003AU03
92RB  L  0.0              0-              4.492 S      20
92RB2 L  %B-=100 $ %B-N=0.0107 5 (1993RU01)$
92RBX L  XREF=AB
-----
192PO  Q              11.0E3 SY 2.2E3      SY 7320      7      1995AU04
192PO  CQ              |DS (n)=360, |DS (p)=450 (1995AU04) .
192PO  L  0.0              0+              33.2 MS      14
192POX L  XREF=AB
192PO2 L  %A AP 100$ %EC+%B+=?$
192PO  CL              %A: only A DECAY observed. %(EC+B+) AP 0.4 can be
192PO2CL estimated from gross B decay theory (partial T AP 8 S)
192PO3CL (1973TA30), or AP 0.54 from partial BETA T of 6.1 S
192PO4CL calculated by 1997MO25.
-----
168RE  Q -5800          SY8960          SY830          SY5063      13      1995AU04
168RE  CQ              |DQ(|b)=400, |DS (n)=420, |DS (p)=510 (1995Au04) .
168RE  L  0.0              (5+, 6+, 7+)          4.4 S      1
168RE2 L  %EC+%B+=100$ %A AP 5E-3 (1992Me10)$
168REX L  XREF=AB
168RE  CL              %A: deduced from IA/RI(199.3G in 168W) and EC decay
168RE2CL scheme for 168RE (1992Me10) .

```

Example 2: decay branches

XREF Flags:

- Use 'X(*)' if level from dataset X cannot be uniquely identified with level in question.
- Use 'X(energy)' to resolve any ambiguity due to poor energy match between adopted level and dataset X level.

Example 3: XREF's

59NI L 5821 10

59NIX L XREF=BN(*5830)

59NI CL JPI=3/2+ FROM (POL P,D) AND L(P,D)=2 FOR 5821 AND/OR

59NI2CL 5844 LEVEL(S).

59NI L 5844 10 (3/2+,5/2+)

59NIX L XREF=BN(*5830)

59NI CL J L(D,P)=(2). JPI=3/2+ FROM (POL P,D) AND L(P,D)=2 FOR 5821

59NI2CL AND/OR 5844 LEVEL(S).

- Watch out for systematic energy scale deviations between various reaction studies.

- Avoid associating a transfer reaction level with an adopted level whose configuration it would not excite.

Example 4

$^{169}\text{Tm}(d,p)$ Target: $\pi 1/2[411]$ g.s.

n stripped from d

^{170}Tm states populated must be $\pi 1/2[411] \otimes \nu \Omega[xxx]$

Populated:

$$\pi 1/2[411] \pm \nu 1/2[521]$$

$$\pi 1/2[411] \pm \nu 5/2[512]$$

$$\pi 1/2[411] \pm \nu 7/2[633]$$

$$\pi 1/2[411] \pm \nu 3/2[521]$$

Not populated:

$$\pi 7/2[404] \pm \nu 7/2[633]$$

$$\pi 1/2[541] \pm \nu 5/2[512]$$

$$\pi 1/2[541] \pm \nu 7/2[633]$$

B(L λ) \uparrow :

Include with level information **only** when value measured, but photon branching or $T_{1/2}$ unknown (e.g., E3 Coulomb excitation measured, but no E3 transition observed).

Moments (μ , Q): static, model-independent values.

- Summarized in 1989Ra17 (evaluation) and 2005St24 (compilation); add any new measurements.
- Specify method used.
- Mention standards used, corrections applied (e.g., Sternheimer).
- Signs matter.
- Convert g-factor data to μ .

$\Delta\langle r^2 \rangle$ (DAVRSQ): include data in comment on g.s. (or isomer) if available.


Example 5: μ , $\Delta\langle r^2 \rangle$, etc.

```
167LU L 0.0+X 1/2(+) 1 M GE CM
167LUX L XREF=B
167LU2 L %EC+%B+=?$%IT=?
167LU3 L MOMM1=-0.0999 13 (1998GE13)$
167LU CL DAVRSQ(170LU,167LU)=-0.291 (1998GE13); 10%
167LU2CL systematic uncertainty.
167LU CL J,MOMM1: from collinear fast beam laser spectroscopy
167LU2CL (1998GE13). PI based on proximity of MOMM1 to value expected for
167LU3CL 1/2[411] orbital (-0.05) cf. that for the only other nearby J=1/2
167LU4CL orbital (viz. 1/2[541], |m AP +0.7).
167LU CL T estimated by 1998GE13; based on known rare-earth diffusion ...
```

Spin and Parity:

- An argument must be provided for every $J\pi$ that is given.
- Use fewest and best strong arguments for definite $J\pi$; the more arguments the better if J or π is uncertain. Try to convince reader; enable a quick check on the impact of any new data that may become available later.
- Use flagged comments for long, repetitive arguments (e.g., “ $J\pi$ based on presence of primary γ from $1/2^+$ capture state in (n, γ) $E=\text{thermal}$ and $\log f^{1ut} < 8.5$ from $1/2^-$ in ... EC decay”).
- If J is directly measured (e.g., atomic beam), state the method.
- Note that μ no longer provides a strong $J\pi$ argument (it used to).
- Avoid using multiply-placed γ 's in “ γ to $J\pi=\dots$ ” type arguments.
- Note that “ γ 's to $3/2^+$ and $5/2^-$ ” (2 levels) differs from “ γ 's to $3/2^+$, $5/2^-$ ” (1 level) – avoid ambiguities.
- “ γ to $J\pi$ ” is a weak argument.
- In “ γ to ...” arguments, the level $J\pi$ is what matters, not $E(\text{level})$.
- Use “ $\log ft=\dots$ from $J\pi=1/2^-$ ” and $L(d,p)=2$ for $9/2^+$ target” type arguments; the parent/target $J\pi$ is part of the argument.

Sample $J\pi$ Arguments:



Argument(s)	$J\pi$
E2 737γ to $7/2+$ g.s.; $\log ft < 5.9$ from $1/2+$.	$3/2+$
Primary γ from $1/2+$ in (n,γ) E=thermal; E1 438γ from $7/2-$ 832 level.	$5/2+$
From (pol d,p) and $L(d,p)=2$ for $0+$ target.	$5/2+$
$\text{Log} f^{1ut} < 8.5$, $\log ft = 7.0$ from $2-$; M1 558γ from $4+$ 1038 level.	$3+$
M1+E2 78γ to $1/2-$ 132 level.	$3/2-$
E1 122γ to $2-$ g.s.; 72γ to $4+$ 50 level.	$(2,3)+$
Probable analog of $3/2-$ 358 level in ^{AA}ZZ .	$(3/2-)$
Unhindered ($HF < 4$) α decay from $(10-)$ parent.	$(10-)$
γ to $2-$ and γ to $4+$.	$(2+,3,4-)$

Gamma-Ray Properties

Energy:

If E_γ came from level-energy difference, say so and recalculate after GTOL has been run (without that E_γ included, of course).

Relative Branching:

- Scale I_γ so strongest branch is 100;

Exceptions:

Strongest line is multiply placed (& in col. 77) (give as $<(I+\Delta I)$).

Strongest line is given as a limit.

Transition is within a superdeformed band.

- Omit uncertainty if only 1 branch.
- Give TI for E0 or fully converted transitions (if known).

Multipolarity:

- [mult] means 'deduced solely from level scheme'; use [E2], *etc.*, only if needed to calculate transition probability or CC for a transition with no measured multipolarity.
- Convert 'D' or 'Q' to '(E1)', '(E2)', *etc.*, if desired or if needed for calculation or $J\pi$ argument; specify how $\Delta\pi$ was deduced.

- Remember that ‘M1,E2’ and ‘M1+E2’ are not equivalent.

Mixing Ratio:

- Include sign, if known. Absence of sign indicates modulus δ .
- If 2 solutions, give both in comment, none in MR field.
- Watch for cases where experiment gives higher limit than RUL allows; modify adopted δ if appropriate.

Total Conversion Coefficient (CC):

Give whenever significant.

E0 Transitions:

Quote $\rho^2(E0)$ from 2005Ki02 or 1999Wo07 (or from authors of later papers who provide it).

Reduced Transition Probabilities:

- Give whenever calculable.
- If δ overlaps 0 or ∞ , calculate for pure D or pure Q, respectively.
- Calc. for [E1], [E2], [$\Delta J > 2$].
- Watch out for δ , $T_{1/2}$ or I_γ data given as a limit.

Reduced Transition Probability Calculations (Special Cases)

I: Data given as limit:

$\delta(M1, E2) < 0.3$:

$B(E2)_W$: give as upper limit.

$B(M1)_W$: give av. of $B(M1)_W(\delta=0)$ and $B(M1)_W(\delta=0.3)$.

$TI < i$ for non-dominant branch:

Assign $1/2i \pm 1/2i$ to this transition to enable calculation of $B(L\lambda)_W$'s for other branches.

$T_{1/2} < t$:

Give resulting lower limits on $B(L\lambda)_W$'s.

$T_{1/2} > t$:

Typically, forget it !

However, $B(E2)_W < 0.005$ or $B(E1)_W < 2 \times 10^{-10}$ might, e.g., be worth mentioning.

II: When $T_{1/2}$ was calculated directly from $B(L\lambda)$:

Calculate $B(L\lambda)_W \downarrow$ from measured $B(L\lambda) \uparrow$ and single-particle value (available from RULER).

Checking Your File

- Make sure that all data sets satisfy current ENSDF policies/practice.
- Run FMTCHK and make the necessary corrections.
- Read through file (ENSDAT output may be helpful); it is amazing what the eye can catch this way!
- Check band drawings – a typographical error in $J\pi$ or an incorrect band flag may be extremely easy to see there.
- Run PANDORA.
 - Use PANDORA.ERR file to identify physics errors.
 - Use PANDORA.GLE file to check for:
 - (i) Inconsistencies in $J\pi$, MULT, δ between adopted and decay datasets.
 - (ii) Adopted photon branching that has not been renormalised so the strongest photon branch is 100.
 - (iii) Levels or transitions in decay or reaction datasets which were accidentally omitted from *Adopted Levels, Gammas* (or conversely).