


2440-7

**16th International Workshop on Computational Physics and Materials Science:
Total Energy and Force Methods**

10 - 12 January 2013

**A new generation of density-functional methods based on the adiabatic-
connection dissipation-fluctuation theorem**


Andreas Goerling
*University of Erlangen-Nürnberg
Germany*



Department
Chemie und Pharmazie



ICP
Interdisziplinäres Zentrum für
"Interface Controlled Processes"



Friedrich-Alexander-Universität
Erlangen-Nürnberg



A new generation of density-functional methods based on the adiabatic-connection fluctuation-dissipation theorem





Andreas Görling, Patrick Bleiziffer, Daniel Schmidtel, and Andreas
Heßelmann

Lehrstuhl für Theoretische Chemie
Universität Erlangen–Nürnberg

J. Gebhardt, G. Gebhardt, T. Gimon, W. Hieringer, C. Neiß, I. Nikiforidis, H. Schulz,
E. Trushin, K.-G. Warnick, T. Wölfle

- 1 Introduction
 - Orbital-dependent functionals
- 2 Exact-exchange (EXX) Kohn-Sham methods
- 3 Direct RPA and EXXRPA correlation energy
 - Adiabatic-connection fluctuation-dissipation theorem for KS correlation energy
 - EXXRPA methods
 - Direct RPA methods
 - Selfconsistent direct RPA
 - Correlation potentials
 - Total energies, reaction energies
 - Bond dissociation, static correlation
 - Weak interactions, Van-der-Waals bonds
- 4 Concluding Remarks
- 5 Literature

Orbital- and eigenvalue-dependent exchange-correlation functionals in Kohn-Sham methods in order to

-  improve accuracy
-  get rid of Coulomb self-interactions and self-interactions in general
 (qualitatively correct KS spectra, improved input data for TDDFT)
-  enable description of static correlation (bond formation and breaking)
-  enable description of Van der Waals interactions

Conventional Kohn-Sham methods

$$E_0 = T_s + U[\rho_0] + E_x[\rho_0] + E_c[\rho_0] + \int d\mathbf{r} v_{\text{nuc}}(\mathbf{r}) \rho_0(\mathbf{r})$$

$$\left[\hat{T} + \hat{v}_{\text{nuc}} + \hat{v}_H[\rho_0] + \hat{v}_x[\rho_0] + \hat{v}_c[\rho_0] \right] \phi_i = \varepsilon_i \phi_i$$

KS methods with orbital- and eigenvalue-dependent xc-functionals

$$E_0 = T_s[\{\phi_i\}] + U[\rho_0] + E_x[\{\phi_i\}] + E_c[\{\varepsilon_s\}, \{\phi_s\}] + \int d\mathbf{r} v_{\text{nuc}}(\mathbf{r}) \rho(\mathbf{r})$$

$$\left[\hat{T} + \hat{v}_{\text{nuc}} + \hat{v}_H[\rho_0] + \hat{v}_x[\{\phi_i\}] + \hat{v}_c[\{\varepsilon_s\}, \{\phi_s\}] \right] \phi_i = \varepsilon_i \phi_i$$

Exchange energy

$$E_x = -\frac{1}{2} \sum_{i,j}^{\text{occ.}} \int d\mathbf{r} d\mathbf{r}' \frac{\phi_i(\mathbf{r}')\phi_j(\mathbf{r}')\phi_j(\mathbf{r})\phi_i(\mathbf{r})}{|\mathbf{r}' - \mathbf{r}|}$$

$$\text{Exchange potential } v_x(\vec{r}) = \frac{\delta E_x[\{\phi_i\}]}{\delta \rho(\vec{r})}$$

$$\int d\mathbf{r}' \chi_0(\mathbf{r}, \mathbf{r}') v_x(\mathbf{r}') = t_x(\mathbf{r})$$

$$\text{KS response function } \chi_0(\mathbf{r}, \mathbf{r}') = 4 \sum_i^{\text{occ.}} \sum_a^{\text{unocc.}} \frac{\phi_i(\mathbf{r})\phi_a(\mathbf{r})\phi_a(\mathbf{r}')\phi_i(\mathbf{r}')}{\epsilon_i - \epsilon_a}$$

$$t_x(\mathbf{r}) = 4 \sum_i^{\text{occ.}} \sum_a^{\text{unocc.}} \frac{\phi_i(\mathbf{r})\phi_a(\mathbf{r}) \langle a | \hat{v}_x^{\text{NL}} | i \rangle}{\epsilon_i - \epsilon_a}$$

Plane wave methods for solid numerically stable
Gaussian basis set methods for molecules numerically demanding

❁ Auxiliary basis set: Electrostatic potential of Gaussian functions

$$v_x(\mathbf{r}) = \sum_k v_{x,k} f_k(\mathbf{r}) \quad \text{with} \quad f_k(\mathbf{r}) = \int d\mathbf{r}' g_k(\mathbf{r}') / |\mathbf{r} - \mathbf{r}'|$$

❁ Incorporation of exact conditions to treat asymptotic of $v_x(\mathbf{r})$


$$\int dr \rho_x(\mathbf{r}) = -1 \quad \text{with} \quad \rho_x(\mathbf{r}) = \sum_k v_{x,k} g_k(\mathbf{r})$$

$$\langle \phi_{HOMO} | v_x | \phi_{HOMO} \rangle = \langle \phi_{HOMO} | \hat{v}_x^{NL} | \phi_{HOMO} \rangle$$


❁ Balancing of auxiliary and orbital basis sets (**uncontracted** orbital basis sets required)

JCP **127**, 054102 (2007)


By preprocessing of auxiliary basis set and regularization techniques, EXX calculations with standard **contracted** orbital basis sets possible (aug-cc-pCVxZ, x = 3, 4, 5)



$2 t_{2u} \quad -2.934 \text{ eV}$




$3 a_{1g} \quad -4.438 \text{ eV}$



$1 t_{2u} \quad -14.724 \text{ eV}$




$2 a_{1g} \quad -22.437 \text{ eV}$



$2 t_{2u} \quad +0.533 \text{ eV}$



$3 a_{1g} \quad -0.396 \text{ eV}$




$1 t_{2u} \quad -9.448 \text{ eV}$




$2 a_{1g} \quad -17.054 \text{ eV}$


contour value 0.032



$2 t_{2u}$ -2.934 eV




$3 a_{1g}$ -4.438 eV




$1 t_{2u}$ -14.724 eV




$2 a_{1g}$ -22.437 eV



$2 t_{2u}$ $+0.533 \text{ eV}$



$3 a_{1g}$ -0.396 eV




$1 t_{2u}$ -9.448 eV



$2 a_{1g}$ -17.054 eV

contour value 0.013

FLAPW vs. PP EXX band gaps



		EXX+VWNc		Exp.
		FLAPW ^a	PP ^b	
Si	$\Gamma \rightarrow \Gamma$	3.21	3.26	3.4
	$\Gamma \rightarrow L$	2.28	2.35	2.4
	$\Gamma \rightarrow X$	1.44	1.50	
SiC	$\Gamma \rightarrow \Gamma$	7.24	7.37	
	$\Gamma \rightarrow L$	6.21	6.30	
	$\Gamma \rightarrow X$	2.44	2.52	2.42
Ge	$\Gamma \rightarrow \Gamma$	1.21	1.28	1.0
	$\Gamma \rightarrow L$	0.94	1.01	0.7
	$\Gamma \rightarrow X$	1.28	1.34	1.3
GeAs	$\Gamma \rightarrow \Gamma$	1.74	1.82	1.63
	$\Gamma \rightarrow L$	1.86	1.93	
	$\Gamma \rightarrow X$	2.12	2.15	2.18
C	$\Gamma \rightarrow \Gamma$	6.26	6.28	7.3
	$\Gamma \rightarrow L$	9.16	9.18	
	$\Gamma \rightarrow X$	5.33	5.43	

^aPRB **83**, 045105 (2011)

^bPRB **59**, 10031 (1997)

$$E_c = \frac{-1}{2\pi} \int_0^1 d\alpha \int d\mathbf{r} d\mathbf{r}' \frac{1}{|\mathbf{r} - \mathbf{r}'|} \int_0^\infty d\omega \left[\chi_\alpha(\mathbf{r}, \mathbf{r}', i\omega) - \chi_0(\mathbf{r}, \mathbf{r}', i\omega) \right]$$

Integration of response functions along complex frequencies

$$\chi_\alpha(\mathbf{r}, \mathbf{r}', i\omega) = -2 \sum_{n \neq 0} \frac{E_n - E_0}{(E_n - E_0)^2 + \omega^2} \langle \Psi_0^\alpha | \hat{\rho}(\mathbf{r}) | \Psi_n^\alpha \rangle \langle \Psi_n^\alpha | \hat{\rho}(\mathbf{r}') | \Psi_0^\alpha \rangle$$

$$\int_0^\infty d\omega \frac{a}{a^2 + \omega^2} = \frac{\pi}{2}$$

later on $g(\mathbf{r}, \mathbf{r}') = \frac{1}{|\mathbf{r} - \mathbf{r}'|}$

$$\frac{-1}{2\pi} \int_0^\infty d\omega \int d\mathbf{r} d\mathbf{r}' g(\mathbf{r}, \mathbf{r}') \chi_\alpha(\mathbf{r}, \mathbf{r}', i\omega) = \int d\mathbf{r} d\mathbf{r}' g(\mathbf{r}, \mathbf{r}') \left[\rho_2^\alpha(\mathbf{r}, \mathbf{r}') - \frac{1}{2} \rho(\mathbf{r}) \rho(\mathbf{r}') + \rho(\mathbf{r}) \delta(\mathbf{r} - \mathbf{r}') \right]$$

Integration along adiabatic connection

$$E_c = \int_0^1 d\alpha V_c(\alpha) \quad \text{with} \quad V_c(\alpha) = \langle \Psi_0(\alpha) | \hat{V}_{ee} | \Psi_0(\alpha) \rangle - \langle \Phi_0 | \hat{V}_{ee} | \Phi_0 \rangle$$

$$E_c = \frac{-1}{2\pi} \int_0^1 d\alpha \int d\mathbf{r} d\mathbf{r}' \frac{1}{|\mathbf{r} - \mathbf{r}'|} \int_0^\infty d\omega \left[\chi_\alpha(\mathbf{r}, \mathbf{r}', i\omega) - \chi_0(\mathbf{r}, \mathbf{r}', i\omega) \right]$$

Required input quantities are $\chi_0(\mathbf{r}, \mathbf{r}', i\omega)$ and $\chi_\alpha(\mathbf{r}, \mathbf{r}', i\omega)$

KS response function $\chi_0(\mathbf{r}, \mathbf{r}', i\omega)$

$$\chi_0(\mathbf{r}, \mathbf{r}', i\omega) = -4 \sum_i^{\text{occ}} \sum_a^{\text{unocc}} \frac{\epsilon_{ai}}{\epsilon_{ai}^2 + \omega^2} \varphi_i(\mathbf{r}) \varphi_a(\mathbf{r}) \varphi_a(\mathbf{r}') \varphi_i(\mathbf{r}')$$

Response functions $\chi_\alpha(\mathbf{r}, \mathbf{r}', i\omega)$ from EXX-TDDFT

$$\mathbf{X}_\alpha = \mathbf{X}_0 + \alpha \mathbf{X}_0 \mathbf{F}_{\text{Hx}} \mathbf{X}_\alpha \quad \mathbf{X}_\alpha = [\mathbf{1} - \alpha \mathbf{X}_0 \mathbf{F}_{\text{Hx}}]^{-1} \mathbf{X}_0$$

$$\mathbf{X}_\alpha = \mathbf{X}_0 [\mathbf{X}_0 - \alpha \mathbf{H}_{\text{Hx}}]^{-1} \mathbf{X}_0 \quad \text{with } \mathbf{X}_0 \mathbf{F}_{\text{Hx}} \mathbf{X}_0 = \mathbf{H}_{\text{Hx}}$$

OEP-like equation for sum $f_{\text{HX}}(\omega, \mathbf{r}, \mathbf{r}')$ of Coulomb and EXX kernel

$$\int d\mathbf{r}'' \int d\mathbf{r}''' \mathbf{X}_0(\mathbf{r}, \mathbf{r}'', \omega) f_{\text{HX}}(\omega, \mathbf{r}'', \mathbf{r}''') \mathbf{X}_0(\mathbf{r}''', \mathbf{r}', \omega) = h_{\text{HX}}(\omega, \mathbf{r}, \mathbf{r}')$$

with

$$\begin{aligned} h_{\text{HX}}(\omega, \mathbf{r}, \mathbf{r}') &= \frac{1}{4} \mathbf{Y}^\top(\mathbf{r}) \boldsymbol{\lambda}(\omega) [\mathbf{A} + \mathbf{B} + \boldsymbol{\Delta}] \boldsymbol{\lambda}(\omega) \mathbf{Y}(\mathbf{r}') \\ &+ \omega^2 \frac{1}{4} \mathbf{Y}^\top(\mathbf{r}) \boldsymbol{\lambda}(\omega) \boldsymbol{\epsilon}^{-1} [\mathbf{A} + \mathbf{B} + \boldsymbol{\Delta}] \boldsymbol{\epsilon}^{-1} \boldsymbol{\lambda}(\omega) \mathbf{Y}(\mathbf{r}) \\ &+ \sum_i \sum_j \sum_a Y_{ia}(\mathbf{r}) \lambda_{ia}(\omega) \frac{\langle a | \hat{v}_x^{\text{NL}} - \hat{v}_x | j \rangle}{\epsilon_a - \epsilon_j} \phi_i(\mathbf{r}) \phi_j(\mathbf{r}) + \dots \\ &+ \sum_a \sum_b \sum_i Y_{ia}(\mathbf{r}) \lambda_{ia}(\omega) \frac{\langle b | \hat{v}_x^{\text{NL}} - \hat{v}_x | i \rangle}{\epsilon_b - \epsilon_i} \phi_a(\mathbf{r}) \phi_b(\mathbf{r}) + \dots \end{aligned}$$

$$A_{ia,jb} = 2(ai|jb) - (ab|ji) \quad B_{ia,jb} = 2(ai|bj) - (aj|bi)$$

$$\Delta_{ia,jb} = \delta_{ij} \langle \varphi_a | \hat{v}_x^{\text{NL}} - \hat{v}_x | \varphi_b \rangle - \delta_{ab} \langle \varphi_i | \hat{v}_x^{\text{NL}} - \hat{v}_x | \varphi_j \rangle$$

$$\lambda_{ia,jb} = \delta_{ia,jb} \frac{-4\epsilon_{ia}}{\epsilon_{ia}^2 + \omega^2} \quad \epsilon_{ia,jb} = \delta_{ia,jb} \epsilon_{ia} = \delta_{ia,jb} (\epsilon_a - \epsilon_i)$$

$$Y_{ia}(\mathbf{r}) = \phi_i(\mathbf{r}) \phi_a(\mathbf{a})$$

$$h_{\text{H}}(\mathbf{r}, \mathbf{r}', \omega) = \mathbf{Y}^\top(\mathbf{r}) \boldsymbol{\lambda}(\omega) \mathbf{C} \boldsymbol{\lambda}(\omega) \mathbf{Y}(\mathbf{r}) \quad \text{with} \quad C_{ia,jb} = (ia|jb)$$

$$E_c = \frac{-1}{2\pi} \int_0^\infty d\omega \int_0^1 d\alpha \int d\mathbf{r} d\mathbf{r}' \frac{1}{|\mathbf{r} - \mathbf{r}'|} \left[\chi_\alpha(\mathbf{r}, \mathbf{r}', i\omega) - \chi_0(\mathbf{r}, \mathbf{r}', i\omega) \right]$$

Representation of $\chi_\alpha(i\omega)$, $h_x(i\omega)$ in RI basis set with respect to Coulomb norm

$$\mathbf{X}_\alpha = [\mathbf{1} - \alpha \mathbf{X}_0 \mathbf{F}_{H_x}]^{-1} \mathbf{X}_0 \quad \Rightarrow \quad \mathbf{X}_\alpha = \mathbf{X}_0 [\mathbf{X}_0 - \alpha \mathbf{H}_{H_x}]^{-1} \mathbf{X}_0$$

(using $\mathbf{F}_{H_x} = \mathbf{X}_0^{-1} \mathbf{H}_{H_x} \mathbf{X}_0^{-1}$)

$$E_c = \frac{-1}{2\pi} \int_0^\infty d\omega \int_0^1 d\alpha \text{Tr} \left[\mathbf{S}^{-1} \mathbf{X}_0 [\mathbf{X}_0 - \alpha \mathbf{H}_{H_x}]^{-1} \mathbf{X}_0 - \mathbf{S}^{-1} \mathbf{X}_0 \right]$$

with

$$\mathbf{X}_0(i\omega) = \mathbf{D}^T \boldsymbol{\lambda}(i\omega) \mathbf{D} \quad \text{with} \quad D_{ia,h} = (\varphi_i \varphi_a | f_h)_{\text{Coul}} \quad \text{and} \quad \lambda_{ia,jb} = \delta_{ia,jb} \frac{-4\epsilon_{ia}}{\epsilon_{ia}^2 + \omega^2}$$

and

$$\mathbf{H}(i\omega) = \frac{1}{4} \mathbf{D}^T \boldsymbol{\lambda}(i\omega) [\mathbf{A} + \mathbf{B} + \Delta] \boldsymbol{\lambda}(i\omega) \mathbf{D} + (i\omega)^2 \frac{1}{4} \mathbf{D}^T \boldsymbol{\lambda}(i\omega) \boldsymbol{\epsilon}^{-1} [\mathbf{A} + \mathbf{B} + \Delta] \boldsymbol{\epsilon}^{-1} \boldsymbol{\lambda}(i\omega) \mathbf{D}$$

$$+ \mathbf{W}_1(i\omega) + \mathbf{W}_1^T(i\omega) + \mathbf{W}_2(i\omega) + \mathbf{W}_2^T(i\omega)$$

$$E_c = \frac{-1}{2\pi} \int_0^\infty d\omega \int_0^1 d\alpha \operatorname{Tr} \left[\mathbf{S}^{-1} \mathbf{X}_0 [\mathbf{X}_0 - \alpha \mathbf{H}_{\text{Hx}}]^{-1} \mathbf{X}_0 - \mathbf{S}^{-1} \mathbf{X}_0 \right]$$

Orthonormalize RI basis set, i.e. make $\mathbf{S} = \mathbf{E}$, and use $\mathbf{X}_0 = -(-\mathbf{X}_0)^{\frac{1}{2}} (-\mathbf{X}_0)^{\frac{1}{2}}$ and carry out analytic integration over coupling constant

$$E_c = \frac{-1}{2\pi} \int_0^\infty d\omega \operatorname{Tr} \left[(-\mathbf{X}_0(i\omega))^{\frac{1}{2}} \mathbf{U}(i\omega) \left(-\tau^{-1}(i\omega) \ln[|\mathbf{1} + \boldsymbol{\tau}(i\omega)|] + \mathbf{1} \right) \mathbf{U}(i\omega)^T (-\mathbf{X}_0(i\omega))^{\frac{1}{2}} \right]$$

$$\text{with } (-\mathbf{X}_0(i\omega))^{-\frac{1}{2}} \mathbf{H}(i\omega) (-\mathbf{X}_0(i\omega))^{-\frac{1}{2}} = \mathbf{U}(i\omega) \boldsymbol{\tau}(i\omega) \mathbf{U}^T(i\omega)$$

- Complete exchange kernel can be treated (RI-EXXRP+)
- N^5 scaling

RI2-dRPA

$$E_c = \frac{-1}{2\pi} \int_0^\infty d\omega \int_0^1 d\alpha \operatorname{Tr} \left[\mathbf{S}^{-1} \mathbf{X}_0 [\mathbf{X}_0 - \alpha \mathbf{H}_H]^{-1} \mathbf{X}_0 - \mathbf{S}^{-1} \mathbf{X}_0 \right]$$

For dRPA, with second RI approximation and $\mathbf{S} = \mathbf{E}$, \mathbf{H}_H simplifies to

$$\mathbf{H}_H = \mathbf{X}_0 \mathbf{X}_0$$

With spectral representation $\mathbf{X}_0(i\omega) = -\mathbf{V}(i\omega) \boldsymbol{\sigma}(i\omega) \mathbf{V}^\top(i\omega)$

$$E_c = \frac{1}{2\pi} \int_0^\infty d\omega \operatorname{Tr} [\ln[\mathbf{1} + \boldsymbol{\sigma}(i\omega)] - \boldsymbol{\sigma}(i\omega)]$$

- Only KS response function $\mathbf{X}_0(i\omega)$ required
- N^4 scaling

Optimized-effective-potential equation for v_c^{dRPA}

$$\int d\mathbf{r}' \chi_0(\mathbf{r}, \mathbf{r}') v_c^{\text{dRPA}}(\mathbf{r}') = t_c^{\text{dRPA}}(\mathbf{r})$$


with

$$t_c^{\text{dRPA}}(\mathbf{r}) = \sum_s \int d\mathbf{r}' \frac{\delta E_c^{\text{dRPA}}}{\delta \phi_s(\mathbf{r}')} \frac{\delta \phi_s(\mathbf{r}')}{\delta v_s(\mathbf{r})} + \sum_s \frac{\delta E_c^{\text{dRPA}}}{\delta \epsilon_s} \frac{\delta \epsilon_s}{\delta v_s(\mathbf{r})}$$

v_c^{dRPA} is represented in auxiliary basis set


$$\mathbf{X}_0 \mathbf{v}_c^{\text{dRPA}} = \mathbf{t}_c^{\text{dRPA}}$$

v_c^{dRPA} for neon atoms





dRPA correlation potential qualitatively correct in contrast to LDA and GGA correlation potential

v_c^{dRPA} for C_2H_2





Deviations of total energies from CCSD(T) energies, $\Delta E = E^{\text{Method}} - E^{\text{CCSD(T)}}$

Orbital basis: aug-cc-pCVTZ RI basis (for E_c and v_{xc}): aug-cc-pVTZ/MP2-fit



Deviations of total energies from CCSD(T) energies, $\Delta E = E^{\text{Method}} - E^{\text{CCSD(T)}}$


Orbital basis: aug-cc-pCVQZ RI basis (for E_c and v_{xc}): aug-cc-pVQZ/MP2-fit



Deviations of reaction energies from CCSD(T) reaction energies

Orbital basis: aug-cc-pCVTZ


RI basis (for E_x and v_{xc}): aug-cc-pVTZ/MP2-fit




Deviations of reaction energies from CCSD(T) reaction energies

Orbital basis: aug-cc-pCVQZ

RI basis (for E_x and v_{xc}): aug-cc-pVQZ/MP2-fit




Orbital basis: aug-cc-pCVQZ RI basis (for E_c and v_{xc}): aug-cc-pVQZ/MP2-fit




Dissociation limit of other molecules (CO , N_2 , etc.) is also treated correctly

Coupling constant integrand


$$E_c = \int_0^1 d\alpha V_c(\alpha)$$



$V_c(\alpha)$




$V_c^{HL}(\alpha)$



Special treatment of singularity in ω -Integrand ($-\tau^{-1} \ln [|\mathbf{1} + \tau(i\omega)|] + 1$)

Twisting of ethene

Orbital basis: avqz RI basis: avqz



RI-EXXRPAs for noncovalent bonds

Interaction energies in kcal/mol

Complex	Basis	CCSD(T)	dRPA @EXX	dRPA @DRPA	EXXRPAs @EXX	EXXRPAs @dRPA	MP2	CCSD ^a
(NH ₃) ₂	3	3.00	2.36	2.41	2.84	2.89	3.00	2.75
	4		2.52	2.57	2.83	2.99	3.11	
	CBS	3.15	2.62	2.64	2.81	3.02	3.18	2.88
(H ₂ O) ₂	3	4.83	3.95	4.03	4.81	4.92	4.72	4.60
	4		4.25	4.33	4.91	5.07	5.02	
	CBS	5.07	4.40	4.46	4.92	5.09	5.21	4.74
(HCONH ₂) ₂	3	15.37	13.65	13.95	15.56	15.90	15.08	14.65
	4		14.32	14.67	16.01	16.42	15.57	
	CBS	16.11	14.75	15.06	16.28	16.67	15.86	15.28
(HCOOH) ₂	3	17.86	15.51	16.03	18.28	18.73	17.61	17.08
	4		16.41	16.98	18.79	19.34	18.23	
	CBS	18.81	16.97	17.49	19.06	19.60	18.61	17.92
(C ₂ H ₄) ₂	3	1.38	0.99	0.99	1.44	1.33	1.47	1.04
	4		1.08	1.05	1.55	1.42	1.55	
	CBS	1.48	1.16	1.05	1.64	1.44	1.60	1.14
C ₂ H ₄ ...C ₂ H ₂	3	1.44	1.23	1.16	1.34	1.50	1.59	1.26
	4		1.31	1.21	1.42	1.56	1.64	
	CBS	1.50	1.37	1.22	1.49	1.58	1.67	1.31
(CH ₄) ₂	3	0.50	0.31	0.32	0.41	0.41	0.46	0.38
	4		0.35	0.36	0.44	0.44	0.48	
	CBS	0.53	0.37	0.35	0.44	0.44	0.53	0.41
RMS	3		1.19	0.97	0.19	0.39	0.13	0.45
RMS	CBS		0.93	0.73	0.19	0.37	0.15	0.51

^a CBS obtained via F12-CCSD

- ❁ EXXRPAs correlation functional combines accuracy at equilibrium geometries with a correct description of dissociation (static correlation) and a highly accurate treatment of VdW interactions
- ❁ EXXRPAs correlation functional is self-interaction free
- ❁ Self-consistent dRPA yields qualitatively reasonable correlation potentials in contrast to conventional KS methods
- ❁ Promising starting point for further developments, e.g., completely self-consistent EXXRPAs method or inclusion of correlation in kernel

Orbital-dependent functionals open up fascinating new possibilities in DFT

EXX for solids

Phys. Rev. Lett. **79**, 2089 (1997)

Phys. Rev. B **83**, 045105 (2011)

Phys. Rev. B **81**, 155119 (2010)

EXX for molecules

Phys. Rev. Lett. **83**, 5459 (1999)

J. Chem. Phys. **128**, 104104 (2008)

TDEXX

Phys. Rev. A **80**, 012507 (2009)

Int. J. Quantum. Chem. **110**, 2202 (2010)

Phys. Rev. Lett. **102**, 233003 (2009)

J. Chem. Phys. **134**, 034120 (2011)

EXX-RPA

Mol. Phys. **108**, 359 (2010)

Phys. Rev. Lett. **106**, 093001 (2011)

Mol. Phys. **109**, 2473 (2011) Review

J. Chem. Phys. **136**, 134102 (2012)

Ground state energy of an electronic system

$$E_0 = T_s + U + E_x + E_c + \int d\mathbf{r} v_{nuc}(\mathbf{r})\rho(\mathbf{r})$$

Thomas-Fermi-Dirac

$$E_0 = T_s[\rho] + U[\rho] + E_x[\rho] + E_c[\rho] + \int d\mathbf{r} v_{nuc}(\mathbf{r})\rho(\mathbf{r})$$

$$\delta E / \delta \rho(\mathbf{r}) = \mu$$

Conventional Kohn-Sham

$$E_0 = T_s[\{\phi_i\}] + U[\rho] + E_x[\rho] + E_c[\rho] + \int d\mathbf{r} v_{nuc}(\mathbf{r})\rho(\mathbf{r})$$

$$[\hat{T} + \hat{v}_{nuc} + \hat{v}_H + \hat{v}_x + \hat{v}_c]\phi_i = \varepsilon_i \phi_i$$

Kohn-Sham with orbital dependent functionals

$$E_0 = T_s[\{\phi_i\}] + U[\rho] + E_x[\{\phi_i\}] + E_c[\{\phi_i\}] + \int d\mathbf{r} v_{nuc}(\mathbf{r})\rho(\mathbf{r})$$

$$[\hat{T} + \hat{v}_{nuc} + \hat{v}_H + \hat{v}_x + \hat{v}_c]\phi_i = \varepsilon_i \phi_i$$