Excitations of the antiferromagnetic XXZ spin-1/2 spin chain - spinons and Bethe strings

A. K. Bera,^{1,2} J. Wu,³ W. Yang,⁴ R. Bewley,⁵ M. Boehm,⁶ J. Xu,^{1,7} M. Bartkowiak,^{1,5} O. Prokhnenko,¹ B. Klemke,¹ A. T. M. N. Islam,¹ J. M. Law,⁸ Z. Wang,⁹ and B. Lake^{1,10}

¹Helmholtz-Zentrum Berlin fur Materialien und Energie, Berlin, Germany ²Technical Solid State Physics Division,

Bhabha Atomic Research Centre, Mumbai, India

³Tsung-Dao Lee Institute & School of Physics and Astronomy,

Shanghai Jiao Tong University, Shanghai, China

⁴Stewart Blusson Quantum Matter Institute,

Vancouver, British Columbia, Canada

⁵ISIS Facility, STFC Rutherford Appleton Laboratory, Harwell Oxford, Didcot, UK ⁶Institut Laue–Langevin, Grenoble, France

⁷Heinz Maier-Leibnitz Zentrum at Lichtenbergstr. 1, 85748, Garching, Germany

⁸Hochfeld-Magnetlabor Dresden (HLD-EMFL),

 $Helmholtz\hbox{-}Zentrum \ Dresden\hbox{-}Rossendorf, \ Dresden, \ Germany$

⁹Institute of Physics II, University of Cologne, Cologne, Germany

¹⁰Institut für Festkorperphysik, Technische Universitat Berlin, Berlin, Germany

The antiferromagnetic spin-1/2 spin chain with Heisenberg-Ising (XXZ) anisotropy is a rich source of novel phenomena. Good physical realizations are the compounds $SrCo_2V_2O_8$ and $BaCo_2V_2O_8$ where the Co^{2+} ions have effective spin-1/2 and are coupled by antiferromagnetic interactions into chains while long-range magnetic order occurs at $T_N = 5$ K due to weak interchain coupling. The excitations are spinons which become bound into pairs to form magnons below T_N . In a longitudinal magnetic field applied along the easy axis, the magnetic order is suppressed and using inelastic neutron scattering and optical spectroscopy we find the first evidence for complex bound states of magnetic excitations, known as Bethe strings [1,2]. Furthermore, the characteristic energy, scattering intensity and linewidth of the observed string states exhibit excellent agreement with precise Bethe ansatz calculations. Our results confirm the existence of the long-sought Bethe string excitations which were predicted almost a century ago [3], establish their role in the quantum spin dynamics of onedimensional systems and reveal the effectiveness of new measurement and calculation techniques.

References:

- Z. Wang et al., Nature 554 219 (2018).
- [2] A. K. Bera et al., Nature Physics 16 625 (2020).
- [3] H. Bethe. Z. Phys. 71, 205 226 (1931).