## **Introduction to Wannierization**



## **Javier Junquera**



### Most important references followed in this lecture

## Berry Phases in Electronic Structure Theory

Electric Polarization, Orbital Magnetization and Topological Insulators

DAVID VANDERBILT

ISBN: 978-1-107-15765-1

### Important bibliography:

#### For a review on Maximally Localized Wannier functions:

#### REVIEWS OF MODERN PHYSICS, VOLUME 84, OCTOBER-DECEMBER 2012

#### Maximally localized Wannier functions: Theory and applications

#### Nicola Marzari

Theory and Simulation of Materials (THEOS), École Polytechnique Fédérale de Lausanne, Station 12, 1015 Lausanne, Switzerland

#### Arash A. Mostofi

Departments of Materials and Physics, and the Thomas Young Centre for Theory and Simulation of Materials, Imperial College London, London SW7 2AZ, United Kingdom

#### Jonathan R. Yates

Department of Materials, University of Oxford, Parks Road, Oxford OX1 3PH, United Kingdom

#### Ivo Souza

Centro de Física de Materiales (CSIC) and DIPC, Universidad del País Vasco, 20018 San Sebastián, Spain and Ikerbasque Foundation, 48011 Bilbao, Spain

#### **David Vanderbilt**

Department of Physics and Astronomy, Rutgers University, Piscataway, New Jersey 08854-8019, USA

## Outline

Definition

**Basic properties** 

**Potential uses** 

**Maximally Localized Wannier Functions** 

Wannierization in SIESTA

## Outline

#### Definition

**Basic properties** 

**Potential uses** 

**Maximally Localized Wannier Functions** 

Wannierization in SIESTA

Let us assume that we have isolated bands  $E_n(\vec{k})$  meaning that they never touch the bands below or above them

We can assume that  $E_n(\vec{k})$  are smooth, continuous and periodic functions in  $\vec{k}$  in the 3D reciprocal space



Let us assume that we have isolated bands  $E_n(\vec{k})$  meaning that they never touch the bands below or above them

We can assume that  $E_n(\vec{k})$  are smooth, continuous and periodic functions in  $\vec{k}$  in the 3D reciprocal space

Then, it is natural to consider its Fourier transform in real space

$$E_{n\vec{R}} = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} E_{n\vec{k}} \ d^3k \iff E_{n\vec{k}} = \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} E_{n\vec{R}}$$
**FFT**

If  $E_n(\vec{k})$  is smooth in  $\vec{k}$  space, we can expect to be large only for a few lattice vectors  $\vec{R}$  near the origin, and decay rapidly with increasing  $|\vec{R}|$ 



We can choose a smooth and periodic gauge for the Bloch functions  $|\psi_{n\vec{k}}\rangle$  associated with each band

The Fourier-transform partners of the Bloch functions are known as Wannier functions associated with band *n* 

$$|w_{n\vec{R}}\rangle = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} |\psi_{n\vec{k}}\rangle d^3k \quad \Longleftrightarrow \quad |\psi_{n\vec{k}}\rangle = \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} |w_{n\vec{R}}\rangle$$
**FFT**

As long as  $\psi_{n\vec{k}}(\vec{r})$  is a smooth function of  $\vec{k}$ , then  $w_{n\vec{R}}(\vec{r})$  decays rapidly with  $|\vec{R}|$  for a given  $\vec{r}$ .



We can choose a smooth and periodic gauge for the Bloch functions  $|\psi_{n\vec{k}}\rangle$  associated with each band

The Fourier-transform partners of the Bloch functions are known as Wannier functions associated with band *n* 

$$|w_{n\vec{R}}\rangle = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} |\psi_{n\vec{k}}\rangle d^3k \quad \Longleftrightarrow \quad |\psi_{n\vec{k}}\rangle = \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} |w_{n\vec{R}}\rangle$$

This is an special case of an unitary transformation

We can view the Bloch and Wannier functions as providing two different basis sets describing the same manifold of states associated with the electron band in question

The Fourier-transform partners of the Bloch functions are known as Wannier functions



$$\begin{split} |w_{2\vec{R}}\rangle &= \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} |\psi_{2\vec{k}}\rangle d^3k \\ |w_{1\vec{R}}\rangle &= \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} |\psi_{1\vec{k}}\rangle d^3k \end{split}$$

## Outline

Definition

**Basic properties** 

**Potential uses** 

**Maximally Localized Wannier Functions** 

Wannierization in SIESTA

**1.** They are localized functions in real space

$$|w_{nec{R}}(ec{r})| o 0$$
 as  $|ec{r}-ec{R}|$  gets large

Each Wannier function  $w_{n\vec{R}}(\vec{r})$  is a localized function centered near  $\vec{R}$ , so it is more natural to describe the situaton by saying that  $w_{n\vec{R}}(\vec{r})$  decays rapidly with  $|\vec{r} - \vec{R}|$ 

2. The Wannier functions are translational images of one another

$$w_{n\vec{R}}(\vec{r}) = w_{n\vec{0}}(\vec{r} - \vec{R})$$

Or more formally

$$|n\vec{R}\rangle = T_{\vec{R}}|n\vec{0}\rangle$$

 $T_{ec{R}}$  is the operator that translates the system by a lattice vector  $ec{R}$  .

#### 3. The Wannier functions form an orthonormal set

Single band case

$$\langle w_{n\vec{R}} | w_{n\vec{R}'} \rangle = \delta_{\vec{R}\vec{R}'}$$

4. The Wannier functions span the same subspace of the Hilbert space as is spanned by the Bloch functions from which they are constructed

#### Single band case

Let  $\mathcal{P}_n$  be the projector operator on band n then this can be expressed as

$$\mathcal{P}_n = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} |\psi_{n\vec{k}}\rangle \langle \psi_{n\vec{k}} | d^3k = \sum_{\vec{R}} |w_{n\vec{R}}\rangle \langle w_{n\vec{R}} |$$

From this it follows that the total charge density  $\rho_n(\vec{r})$  in band *n* is the same when computed in every representation

$$\rho_n(\vec{r}) = -e\langle \vec{r} | \mathcal{P}_n | \vec{r} \rangle = -e \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} |\psi_{n\vec{k}}(\vec{r})|^2 d^3k = -e \sum_{\vec{R}} |w_{n\vec{R}}(\vec{r})|^2$$



Example in 1D

- Exponentially localized
- Normalized
- Neighboring Wannier functions are periodic images of one another
- They are shown as having a negative lobe so that  $\langle w_{n0}|w_{na}\rangle$  vanish as a result of the cancellation between contributions of opposite sign in the integral over  $\mathcal{X}$

# The Hamiltonian matrix elements between Wannier functions are band-diagonal

$$\langle w_{m\vec{R}}|H|w_{n\vec{R}'}
angle=0 \qquad {\rm if} \qquad n
eq m$$

In the previous example

 $\langle w_{1\vec{R}}|H|w_{2\vec{R}'}\rangle=0$ 



$$|w_{2\vec{R}}\rangle = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} |\psi_{2\vec{k}}\rangle d^3k$$

$$w_{1\vec{R}}\rangle = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} |\psi_{1\vec{k}}\rangle d^3k$$

The diagonal elements of the Hamiltonian matrix elements between Wannier functions are the coefficients in the Fourier expansion of the band energy

$$\langle w_{n\vec{0}}|H|w_{n\vec{R}}\rangle = E_{n\vec{R}}$$

where

$$E_{n\vec{R}} = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} E_{n\vec{k}} \ d^3k$$

From the knowledge of:

- the band structure
- The k-point mesh
- The unit cell volumen

We get the values of the Hamiltonian matrix written in a basis of Wannier functions

## Outline

Definition

**Basic properties** 

**Potential uses** 

**Maximally Localized Wannier Functions** 

Wannierization in SIESTA

# The Wannier functions provide an exact tight-binding representation of the dispersion $E_{n\vec{k}}$ of band n

The Fourier components of the energy can be considered as the on-site and hopping elements of a tight-binding Hamiltonian

Here we show three periodic replicas and their corresponding Wannier centers



The hoppings correspond to  $t_1 \rightarrow E_{n,\vec{a}_1}$  $t_2 \rightarrow E_{n,\vec{a}_2}$  $t_{12} \rightarrow E_{n,\vec{a}_1+\vec{a}_2}$ 

 $E_{nec{0}}$  is the on-site energy

The tight-binding equations are nothing else that the expression of the inverse Fourier transform for the energy terms

$$\tilde{H}_{ij}(\vec{k}) = \langle \tilde{\chi}_i(\vec{k}) | H | \tilde{\chi}_j(\vec{k}) \rangle = \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} H_{ij}(\vec{R})$$

$$\sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} E_{n\vec{R}} = E_{n\vec{k}}$$

And, by construction, it reproduces exactly the band dispersion

# Wannier interpolation: approach of keeping only a few hoppings

Since the Wannier functions are localized, the hopping matrix elements decay rapidly with distances, so that a few hoppings are tipically retained

$$E_{n\vec{k}} \approx \sum_{\substack{\vec{R} \\ |\vec{R}| < R_{\rm c}}} e^{i\vec{k}\cdot\vec{R}} E_{n\vec{R}}$$

If these are the dominant hoppings, this simple model may be expected to provide a good approximation to the full band structure.

But the model is guaranteed to become more and more accurate as farther-neighbor hoppings are added

## Relationship between Wannier centers and Berry phase polarization (isolated bands)

The electronic contribution to the polarization is determined entirely by the Wannier centers of the occupied bands

$$\vec{P} = \frac{-e}{V_{\text{cell}}} \sum_{n}^{\text{occ}} \vec{\vec{r}}_{n}$$

 $\overline{\vec{r}_n} = \langle w_{n\vec{0}} | \vec{r} | w_{n\vec{0}} \rangle$ 



## Outline

Definition

**Basic properties** 

**Potential uses** 

**Maximally Localized Wannier Functions** 

Wannierization in SIESTA

### Wannier functions are not unique

$$|w_{n\vec{R}}\rangle = \frac{V_{\text{cell}}}{(2\pi)^3} \int_{\text{BZ}} e^{-i\vec{k}\cdot\vec{R}} |\psi_{n\vec{k}}\rangle d^3k \quad \Longleftrightarrow \quad |\psi_{n\vec{k}}\rangle = \sum_{\vec{R}} e^{i\vec{k}\cdot\vec{R}} |w_{n\vec{R}}\rangle$$

- Many different gauge transformations to chose the phases of the Bloch functions are possible

As a result, the Wannier functions may change shape, becoming somewhat more or less localized

#### **Maximally localized Wannier function**

How to chose a sensible gauge for the computation of the Wannier functions

**Criterion of maximal localization:** 

minimize the sum of spreads of the Wannier functions in one unit cell

$$\Omega_{\text{spread}} = \sum_{n=1}^{J} \left[ \langle w_{n\vec{0}} | r^2 | w_{n\vec{0}} \rangle - |\bar{\vec{r}}_n|^2 \right]$$

Mean square variation of the Wannier electron density away from its mean position

N. Marzari and D. Vanderbilt, Phys. Rev. B 56, 12847 (1997)

This is the criterion followed in WANNIER90

## Outline

Definition

**Basic properties** 

**Potential uses** 

**Maximally Localized Wannier Functions** 

Wannierization in SIESTA

### Interface between SIESTA and WANNIER90

#### http://www.wannier.org

OPEN ACCESS OP Publishing

J. Phys.: Condens. Matter 32 (2020) 165902 (25pp)

https://doi.org/10.1088/1361-648X/ab51ff

# Wannier90 as a community code: new features and applications

Giovanni Pizzi<sup>1,29,30</sup>, Valerio Vitale<sup>2,3,29</sup>, Ryotaro Arita<sup>4,5</sup>, Stefan Blügel<sup>6</sup>, Frank Freimuth<sup>6</sup>, Guillaume Géranton<sup>6</sup>, Marco Gibertini<sup>1,7</sup>, Dominik Gresch<sup>8</sup>, Charles Johnson<sup>9</sup>, Takashi Koretsune<sup>10,11</sup>, Julen Ibañez-Azpiroz<sup>12</sup>, Hyungjun Lee<sup>13,14</sup>, Jae-Mo Lihm<sup>15</sup>, Daniel Marchand<sup>16</sup>, Antimo Marrazzo<sup>10</sup>, Yuriy Mokrousov<sup>6,17</sup>, Jamal I Mustafa<sup>18</sup>, Yoshiro Nohara<sup>19</sup>, Yusuke Nomura<sup>4</sup>, Lorenzo Paulatto<sup>20</sup>, Samuel Poncé<sup>21</sup>, Thomas Ponweiser<sup>22</sup>, Junfeng Qiao<sup>23</sup>, Florian Thöle<sup>24</sup>, Stepan S Tsirkin<sup>12,25</sup>, Małgorzata Wierzbowska<sup>26</sup>, Nicola Marzari<sup>1,29</sup>, David Vanderbilt<sup>27,29</sup>, Ivo Souza<sup>12,28,29</sup>, Arash A Mostofi<sup>3,29</sup>, and Jonathan R Yates<sup>21,29</sup>

#### Can be run as:

- Postprocessing tool
- As a library within SIESTA (coming son).

#### **Requirements to feed WANNIER90 from SIESTA**

SIESTA must compute:

## The overlap of the periodic part of the wave functions at neighbour k-points

 $M_{mn}^{(\vec{k},\vec{b})} = \langle u_{m\vec{k}} | u_{n\vec{k}+\vec{b}} \rangle$ 

 $\vec{b}$  is the vector which connect a given  $\vec{k}$  with its neighbours

The overlap of the Bloch states  $|\psi_{m\vec{k}}
angle$  onto trial localised orbitals  $|g_n
angle$  $A_{mn}^{(\vec{k})}=\langle\psi_{m\vec{k}}|g_n
angle$ 

The eigenvalues of the Hamiltonian in grid of k-points

# Exercise 1: Wannierize the top of the valence band and the bottom of the conduction band of SrTiO<sub>3</sub>

#### Top of valence band



#### O 2p in character

#### **Bottom of conduction band**



#### Ti t<sub>2g</sub> in character

# Exercise 2: Wannierize the bands of Graphene (disentanglement)





#### $\boldsymbol{\pi}$ bonding

#### $\sigma$ bonding

# Exercise 3: Plotting the Fermi Surface of doped SrTiO<sub>3</sub>





#### **Doping with electrons**

#### **Doping with holes**