



## Joint ICTP-IAEA School on Atomic and Molecular Processes in Plasmas | (SMR 4214)

04 May 2026 - 08 May 2026  
ICTP, Trieste, Italy

---

**P01 - ABDELAZIM Ahmed Abdelazim Ahmed**

THERMAL AND ELECTROSTATIC CONTROL OF ATOMIC EMISSION, SPECTRAL CONTRAST, AND SELF-ABSORPTION IN ULTRAFAST PICOSECOND LIBS OF COPPER

**P02 - ALARCON Miguel Angel**

Characterization of states of Ra with relativistic variational R-matrix method

**P03 - ANDREI Osipov**

Development of a Framework for Modelling Reaction Systems in Fusion Plasmas

**P04 - ARYA Sudhanshu**

Electron impact total and partial ionization cross sections for  $W(CO)_6$  and  $Fe(CO)_5$

**P05 - BANERJEE Soumitra**

A Novel Diagnostic Probe for the Direct Detection of Runaway Electrons in ADITYA-U Tokamak

**P06 - BANKS Joel**

Electron Scattering on Gallium

**P07 - BREWSTER Bowie**

Spectral Narrowing in Highly Charged Tin Ions

**P08 - CHEBYKIN Evgenii**

Synchrotron calibration of a transmission grating spectrometer for broadband EUV measurements.

**P09 - CHILDERS Ryan**

A Python-based Hybrid Structure Collisional-Radiative Model for Spectroscopic Plasma Diagnostics and Equation-of-state Computations

**P10 - DOUGAN David John**

A New Approach for Ground and Excited State Electron-Impact Ionization of C I within Magnetically Confined Plasmas

**P11 - EL MESBAHI Lamya**

Modeling Atomic Processes in Fusion Plasma

**P12 - FALTA Falta**

Atomic Structure Calculations and identification of EUV and FUV spectral lines of  $W_{10+}$

**P13 - FERGUSON Niamh**

The Challenge of Accurate Recombination Data for Lanthanides and Actinides

**P14 - FORDE Aaron**

Atomic and Molecular Opacity in Low-Temperature Plasmas: Quantifying Uncertainty in Atomic and Plasma Physics Models

**P15 - FORESTIER COLLEONI Pierre**

Temporal and Spatial evolution of the ion temperature in the WEST tokamak

**P16 - GARCIA-MUÑOZ Alejandro Luis**

Theoretical study of electron-molecule collisions for plasma simulations

**P17 - HENRY John**

Dirac R-matrix electron-impact excitation of Ni I for the interpretation of magnetically-confined/astrophysical plasmas

**P18 - HOQUE Sk Injamul**

Effect of pulsed neutral gas injection on heat flux scrape-off layer (SOL) width in ADITYA-U tokamak

**P19 - HVIZDOS David**

Theoretical Advances in Electron Molecular-ion Scattering, Method Benchmarking and Connecting Collisional and Photofragmentation Resonances

**P20 - JAFAR Nurlana**

Mechanisms of Plasma–Matrix Interactions and Their Compensation Strategies in ICP-MS

**P21 - KHETO Soumyo**

Irradiation-driven fragmentation of  $W(CO)_6$  in gas phase: Analysis of scaling behaviour

**P22 - KLEIN Merlin**

Spectroscopic Time-of-Flight Spatial Distribution Measurements of Mono- and Polycrystalline W with Varying Surface Properties in Ne and Ar Plasma at the Linear Plasma Device PSI-2

**P23 - LEE Changhoo**

Observation of Non-Conventional Electron Dynamics and Temperature Scaling Driven by Relativistic Brunel Heating

**P24 - LEE Kyungbae**

Frustrated Brunel Heating by Relativistic Gyromagnetic effects in Ultra-intense Laser-Matter Interaction

**P25 - LEWIS Sina Genevieve**

First principles optical constants of water ice: geometry effects and vibrational contributions

**P26 - LÓPEZ-MIRANDA Belén**

Analysis of carbon ion temperatures using an upgraded multichannel Doppler spectroscopic diagnostic in NBI-heated TJ-II plasmas

**P27 - MADA Gurubhavannarayana**

Development of machine learning augmented CR models using the cross-sections calculated by relativistic distorted wave method.

**P28 - MEIER Vanni**

Laser-collision induced fluorescence for measurement of collision rate coefficients in helium plasmas

**P29 - MER Pinalben Chandrakantbhai**

Electron and positron driven molecular processes for W and WH for application in plasma systems

**P30 - MEZLINI Rawand**

ELECTRON-INDUCED VIBRATIONAL TRANSITIONS AND DISSOCIATIVE RECOMBINATION OF  $H_2^+$  AND  $BH^+$

**P31 - MORI Nicolas Antonio**

Construction of atomic collisional radiative models for use in solar atmospheric modelling

**P32 - MUKHERJEE Nilanjan**

Electron collision studies on boron hydride species in fusion edge plasmas

**P33 - NAM Jeongwoo**

Steady-State-Informed Latent Dynamics Surrogate Model for TD-NLTE Atomic Kinetics

**P34 - NGUELEO BALDAGUI Joseph**

Effective cross sections and rates coefficients derived from collision-induced rotational excitation of  $\text{HCl}^{\{+\}}$  ( $X^{\{2\}}\Pi$ ) with  $\text{He}^{\{1\}}S$ : Isotopic effects

**P35 - PARKER Payton**

Absolute calibration of various crystal types and geometries for National Ignition Facility x-ray spectrometers

**P36 - POPOVA Maria**

Near-Surface Hydrogen Inventory Response to Picosecond Laser Pulses in Tungsten

**P37 - RENGGLI Amélie**

Optical Emission Spectroscopy in Helium plasmas for validation of collisional-radiative model on the Resonant Antenna Ion Device

**P38 - RYNDYK Daniil**

EMC3-EIRENE predictions of radiative detachment scenarios in W7-X equipped with a tungsten based divertor

**P39 - SCHRICK Kyna**

A general approach to calculating ion-atom cross sections, applied to proton collisions with oxygen

**P40 - ŠENK Jan**

Interparticle Coulombic electron capture beyond the virtual photon approximation

**P41 - SHI Lei**

Full-dimensional quantum dynamic study for the H atom scattering from graphene surface

**P42 - SINGHAL Akansha**

Relativistic Calculations Of Transition Parameters Of Doubly Ionized Argon

**P43 - SINGOR Adam Julianus Chant**

Photoionization, Rayleigh, and Raman scattering cross sections for the hydrogen molecule

**P44 - STANGL Sarah Marie**

GEOSH: Chemical Equilibrium Calculations of Ideal Gas Mixtures

**P45 - STEFANIKOVA Radka**

1D Imaging X-ray Spectroscopy for Inferring Time-Integrated, Spatially-Resolved Volumetric Temperature Distributions in Relativistic Laser-Solid Interactions

**P46 - TATSCH Maria**

Vibrationally resolved core excitation of the  $\text{CN}^+$  molecular ion

**P47 - TIMKOVSKII Georgii**

Initial predictive modeling of plasma-wall interactions using ERO2.0 for W7-X with tungsten wall divertor

**P48 - UNİYAL Akash**

Machine Learning-Assisted Diagnostics of Laser-Induced Aluminium Plasma

**P49 - VAN HOOMISSEN Timothy Russell**

Streaked Visible Spectroscopy Analysis for Laser-Driven Collisionless Shocks on Z

**P50 - VARSHNEY Ruchi**

Characterisation of Neutral Plasma Interactions for Gas Puff Imaging Using the DEGAS-2 Code

**P51 - WHITE Jackson Richard**

Self-Energy Line Broadening in Dense Plasmas

**P52 - XU Yanlan**

Dielectronic Recombination Study on Sr<sup>4+</sup>, Se<sup>4+</sup>, Ga<sup>4+</sup> and In<sup>4+</sup> Ions

# THERMAL AND ELECTROSTATIC CONTROL OF ATOMIC EMISSION, SPECTRAL CONTRAST, AND SELF-ABSORPTION IN ULTRAFAST PICOSECOND LIBS OF COPPER

**Ahmed Abdelazim<sup>1</sup>, Mohamed Fikry<sup>1</sup>, Omar K. Okasha<sup>1</sup>, Abdelnaser Aboulfotouh<sup>1</sup>, Magdy Omar<sup>1</sup>**

<sup>1</sup> *Ultrafast Picosecond Laser Lab, Physics Department, Faculty of Science, Cairo University, Giza 12613, Egypt*

Laser-induced breakdown spectroscopy (LIBS) is a versatile technique for elemental analysis; however, its analytical performance is often constrained by plasma–radiation coupling effects such as continuum background and spectral line self-absorption. These challenges are particularly pronounced in ultrafast (picosecond) LIBS, where dense, transient plasmas are formed, and the emitted radiation is highly sensitive to charge transport and collisional–radiative balance.

This work presents an integrated plasma-control strategy that combines **target preheating** with a **static external electric field** to modulate atomic emission and spectral contrast in air. Building on recent advances in enhancement methods for LIBS, we focus on how thermal and electrostatic boundary conditions reshape the near-surface sheath and influence the redistribution of charged species within the expanding plume. Preheating increases the ablated mass flux and plume density, leading to stronger line emission while often enhancing the continuum contribution. In contrast, an external static bias primarily governs charge separation and ion–electron drift through sheath-driven transport and ambipolar fields, enabling plume reshaping without inducing substantial changes in electron temperature.

A key outcome of this approach is the polarity-dependent decoupling between net line intensity and signal-to-noise ratio behaviour. By tuning the electric-field polarity, the electron density and radiative transfer pathways can be selectively controlled, allowing for the independent optimization of emission strength (sensitivity) and spectral contrast (measurement robustness). This provides a practical route to mitigate self-absorption while preserving strong analytical signals—an essential requirement for quantitative LIBS of conductive targets and trace-level detection.

The proposed framework links field-assisted plasma dynamics with atomic emission formation and radiative processes, and is broadly applicable to ultrafast LIBS diagnostics, including multi-element analysis in complex matrices and real-time monitoring scenarios. A related experimental implementation in picosecond LIBS of copper is currently under review.

## REFERENCES

- [1] A. Abdelazim, A. Aboulfotouh, M. Omar, M. Fikry, *Enhanced Laser-Induced Breakdown Spectroscopy by Pre-heated Target and Electric Field: Comprehensive Review*, **Journal of Optics** (2025). <https://doi.org/10.1007/s12596-025-02651-5>
- [2] D. A. Cremers, L. J. Radziemski, *Handbook of Laser-Induced Breakdown Spectroscopy*, Wiley (2006).
- [3] J. P. Singh, S. N. Thakur (eds.), *Laser-Induced Breakdown Spectroscopy*, Elsevier (2007).

## Characterization of states of Ra with relativistic variational R-matrix method

**Miguel A. Alarcón<sup>1</sup>, and Chris H. Greene<sup>2</sup>**

<sup>1</sup>*Univeristy of Arizona, Tucson, Arizona, 85719, USA*

<sup>2</sup>*Purdue University, West Lafayette, Indiana, 47907, USA*

On the recent experiment of Fan, et al [1] radium atoms were photoionized in order to trap and cool radium ions. This development opens the door to high accuracy measurements of spectroscopic properties of the atom that for which there is no available data.

In this study, we have used the recently implemented relativistic variational R-matrix method [2] to study the radium atom in different energy ranges that complement the existing theory using the relativistic random phase approximation [3] and known experimental data.

We start by studying the position of bound states, and more specifically the highly excited Rydberg series that the variational R-matrix method is well suited to predict. We have calculated and characterized series for both parities and different values of total angular momentum. Our results are in good agreement with the only reported series for radium: the  $7snp\ ^1P_o$  [4].

The second energy range we focus on is the low-lying autoionizing resonances just below the 6d threshold. In this case we obtain the ground-state and first odd parity excited-state one-photon photoionization cross sections. Using techniques from multichannel quantum defect theory [5] we label the found resonances and calculate their quantum defects and widths. In the future, we plan to incorporate the Breit interaction into these codes to improve the accuracy of the calculations of spectroscopic properties.

### References:

- [1] Fan, M., et al (2023). Laser cooling and trapping of  $^{224}\text{Ra}^+$ . *Phys. Rev. Research* 5, 043201
- [2] Alarcón, M., et al (2024). Application of the variational R-matrix method for the Dirac equation to the Be atom. *Phys. Rev. A* 110, 042825
- [3] Deshmukh, P. *et al*, (1992). Photoionization of the outer shells of radon and radium: Relativistic random-phase approximation for high-Z atoms. *Phys. Rev. A* 45, 6339.
- [4] Dammalapati, U. et al (2016). Compilation of Spectroscopic data of radium (RaI and RaII). *J. Phys. Chem. Ref. Data* 45, 013101 (2016)
- [5] Aymar, M. et al (1996). Multichannel Rydberg spectroscopy of complex atoms. *Rev. Mod. Phys* 68, 1015

## Development of a Framework for Modelling Reaction Systems in Fusion Plasmas

**Andrei Ludvig-Osipov, Stefan Mijin, Matthew Bluteau, Kristian Zarebski, Sophie Frankel, and Nathan Cummings**

*UKAEA (United Kingdom Atomic Energy Authority), Culham Campus, Abingdon, Oxfordshire, OX14 3DB, UK*

The atomic and molecular processes play crucial role in the edge region of tokamak plasmas, as experimentally evidenced at magnetic confinement fusion devices, e.g., MAST-U [1], JET, AUG [2] and Alcator C-Mod [3]. A robust simulation capability with accounting for such processes is essential for analysis and planning of experiments and for design of the future fusion devices. As the number of the involved levels of atomic and molecular species may reach hundreds and even thousands (see, e.g., a system with rovibrationally resolved molecular hydrogen [4]), a brute-force inclusion in fluid codes for plasma edge simulations is extremely computationally prohibitive. To account for the atomic and molecular effects while attaining reasonable computational costs, the reduced (effective) models can be used instead.

A new framework for modelling reaction systems in fusion plasmas is currently under development at UKAEA, with a goal of providing a comprehensive toolset for analysis, modelling and reduction of plasma chemical kinetics to the scientific community. The framework is being developed with an emphasis on the best modern software development practices, and with an outlook for provenance and metadata tracking. The present capabilities include import of ADAS adf04 files [5], reduction methods for linear problems (quasi-steady-state assumption and a method by Greenland [6], i.e., classic collisional-radiative modelling), automatic identification of reduced models [6], solution of full and reduced systems, a solver for non-linear system, and synthetic spectrum generation. The reaction systems can be parametrized with scanning and automatic differentiation (using JAX library [7] for Python) capabilities allowing for tabulation of effective rates/cross-coupling coefficients. The planned future work includes implementation of uncertainty quantification and of reduction techniques for non-linear problems, such as Intrinsic Low-Dimensional Manifold [8] and Computational Singular Perturbation [9].

- [1] K. Verhaegh et al., Nucl. Fusion **63**, 016014 (2023).
- [2] M. Bernert et al., Nucl. Mater. Energy **12**, 111-118 (2017).
- [3] A. Mathews et al., Rev. Sci. Instrum. **93**, 063504 (2022).
- [4] K. Sawada, M. Goto, Atoms **4(4)**, 29 (2016).
- [5] H. P. Summers, The ADAS User Manual, version 2.6 <http://www.adas.ac.uk> (2004)
- [6] P. T. Greenland, Proc. R. Soc. A: Math. Phys. Eng. Sci., **457**, 1821-1839 (2001).
- [7] R. Frostig, M. J. Johnson, C. Leary, SysML Conf. 2018, hal-05188750, (2019).
- [8] U. Maas, S.B. Pope, Proc. Combust. Inst. **24**, 103-112 (1992).
- [9] S. H. Lam, D. A. Goussis, Int. J. Chem. Kinet. **26(4)**, 461-486 (1994).

# Electron impact total and partial ionization cross sections for $\text{W}(\text{CO})_6$ and $\text{Fe}(\text{CO})_5$

Sudhanshu Arya<sup>1</sup>, and Bobby Antony<sup>1</sup>

<sup>1</sup>*Atomic and Molecular Physics Laboratory, Department of Physics, Indian Institute of Technology (ISM), Dhanbad, Jharkhand 826004, India*

Tungsten hexacarbonyl,  $\text{W}(\text{CO})_6$ , and iron pentacarbonyl,  $\text{Fe}(\text{CO})_5$ , are widely used volatile precursors in electron/ion driven and plasma-assisted processes such as focused electron beam induced deposition (FEBID), chemical vapor deposition, and plasma-enhanced thin-film growth. In these environments, low-to-intermediate energy electrons initiate ionization and dissociative ionization, controlling precursor depletion, fragment formation (e.g., sequential CO loss), and ultimately the composition and purity of deposited W- and Fe-containing films. Despite this practical importance, comprehensive electron-impact ionization cross sections, particularly partial (channel-resolved) ionization data, remain scarce over the broad energy range needed for plasma kinetics and electron-transport modeling [1]. In this work, we report total ionization cross sections (TICS) and partial ionization cross sections (PICS) for  $\text{W}(\text{CO})_6$  and  $\text{Fe}(\text{CO})_5$  from the ionization threshold up to 10 keV using the Binary-Encounter-Bethe (BEB) framework [2]. Orbital-resolved BEB inputs, such as binding energies, kinetic energies, and occupation numbers, are obtained from quantum-chemistry calculations, and the total cross section is constructed by summing contributions from all occupied orbitals. Where experimental ionization thresholds are available, they are used to anchor the low-energy onset; otherwise, the threshold is estimated from the HOMO energy (Koopmans-like approximation). Partial ionization cross sections are provided in two complementary forms: (i) subshell/orbital-resolved ionization and (ii) dissociative ionization channel-resolved yields derived from mass-spectrometric branching information; when complete branching data are unavailable, we employ physically motivated, energy-dependent branching schemes to generate continuous PICS across the full energy range. The resulting cross-section set is intended to support plasma chemistry, FEBID modeling, and Monte Carlo electron-transport simulations involving metal-carbonyl precursors.

- [1] K. Wnorowski, M. Stano, W. Barszczewska, A. Jówko, Š. Matejčík, *Int. J. Mass Spectrom.* **314**, 42 (2012).
- [2] Y. K. Kim, M. E. Rudd, *Phys. Rev. A* **50**, 3954 (1994).

## A Novel Diagnostic Probe for the Direct Detection of Runaway Electrons in ADITYA-U Tokamak

**Soumitra Banerjee<sup>1,2</sup>, Santosh P. Pandya<sup>1</sup>, Bhargav Soni<sup>1</sup>, Harshita Raj<sup>1,2</sup>, Injamul Hoque<sup>1,2</sup>, Rakesh L. Tanna<sup>1</sup>, Joydeep Ghosh<sup>1,2</sup>**

<sup>1</sup>*Institute for Plasma Research, Gandhinagar, India*

<sup>2</sup>*Homi Bhabha National Institute, Mumbai, India*

Runaway electrons (REs), with energies ranging from a few keV to several MeV, are generated in ADITYA-U tokamak discharges under high loop voltage and low-density conditions. These energetic particles can degrade plasma performance, may damage plasma-facing components (PFCs), or cause structural harm [1], while also serve as diagnostic probes of particle magnetic confinement and instability dynamics. To explore this dual role, a new RE flux probe has been designed, developed, and calibrated for ADITYA-U tokamak [2]. It directly comes into contact with the confined REs and measures their flux deposited on the probe area and functions as an energy analyzer, enabling quantitative evaluation of RE effects on PFCs and detailed studies of their behavior during instabilities. The probe head shape made up from Graphite material is specifically designed to distribute heat flux impinging from plasma over a large area and to optimize heat load, it is insertable into the edge plasma up to ~3 cm through scrap-off layer using the fast-reversed drive system, it employs three Ceriumdoped Lutetium Yttrium Silicate (LYSO:Ce), a high-performance scintillator crystals, with Stainless Steel (SS-304L) filters inside a Graphite shield for detection and subtraction of background Hard Xray signal from RE flux signal. The probe materials and their thickness are carefully designed using a Monte-carlo code PHITS to optimize for its functionality during the measurement namely, the Graphite layer reduces backscattering less than 6% and the Graphite housing restricts detection to REs below 1 MeV. The first crystal records REs above this threshold, the second crystal selects REs exceeding 3 MeV via additional SS filtering, and the third measures only Hard X-Rays background, blocked from the direct incidence of REs by optimized thickness of SS shielding. Experimental performance assessment and calibration are performed in the energy range from 1–10 MeV monochromatic electron beam and compared with PHITS simulations.

[1] B. N. Breizman, P. Aleynikov, E. M. Hollmann, and M. Lehnen, Nucl. Fusion, vol. 59, no. 8, p. 083001 (Jun. 2019).

[2] R. L. Tanna et al., Nucl. Fusion, vol. 63, no. 8, p. 086011 (Aug. 2023).

## Electron Scattering on Gallium

Joel D. Banks, Igor Bray and Dmitry V. Fursa

*Department of Physics and Astronomy, Curtin University, Perth, WA 6102, Australia*

The relativistic convergent close-coupling (RCCC) method [1] is applied to calculate cross sections for electron scattering on atomic and singly ionised gallium [2, 3]. Gallium is important for high-performance semiconductors such as gallium nitride and gallium oxide. Recent research found that semiconductor nanoparticles can be successfully synthesised using gas and plasma phase reactants [4]. Due to gallium's thermal properties, it has also been proposed for use as a liquid plasma-facing surface in future fusion reactor chambers [5]. Both applications result in plasma-gallium interactions that must be accounted for in modelling and design, which would be enabled by a reliable cross section dataset for electron scattering from neutral and ionic gallium.

The Ga atom is modelled as a quasi one-electron atom, while  $\text{Ga}^+$  is modelled as quasi two-electron, each with an inert Dirac-Fock core. Integrated excitation and estimated single ionisation cross sections (ISICS) are presented for the ground state of both targets, at incident energies between 0.03 eV and 1000 eV for Ga, and between 5.9 eV and 500 eV for  $\text{Ga}^+$ . The estimated ISICS include direct ionisation from the  $4p$ ,  $4s$  and  $3d$  electrons and excitation-autoionisation contributions from  $4s$  and  $3d$  electrons, where appropriate. The estimated ISICS for neutral Ga was found to agree very well with existing experimental data [6, 7]. By contrast, the estimated ISICS for  $\text{Ga}^+$  has varying agreement between the various existing experimental results, which are inconsistent among themselves [8, 9, 10].

- [1] D. V. Fursa, I. Bray, *Phys. Rev. Lett.* **35**, 015017 (2026).
- [2] J. Banks, I. Bray, D. V. Fursa, *Plasma Sources Sci. Technol.* **33**, 105021 (2024).
- [3] J. Banks, I. Bray, D. V. Fursa, *Plasma Sources Sci. Technol.* **100**, 113201 (2008).
- [4] K. H. You *et al*, *Curr. Appl. Phys.* **18**, 1553-1557 (2018).
- [5] R. B. Gomes *et al*, *Fusion Eng. Des.* **80**, 102-111 (2008).
- [6] C. J. Patton *et al*, *J. Phys. B* **29**, 1409 (1996).
- [7] R. S. Freund *et al*, *Phys. Rev. A* **41**, 3575-3595 (1990).
- [8] W. T. Rogers *et al*, *Phys. Rev. A* **25**, 737-748 (1982).
- [9] B. Peart, J. R. A. Underwood, *J. Phys. B* **23**, 2343 (1990).
- [10] J. A. Shaw *et al*, *Phys. Rev. A* **63**, 032709 (2001).

## Spectral Narrowing in Highly Charged Tin Ions

**Bowie Brewster<sup>1</sup> and John Sheil<sup>1</sup>**

<sup>1</sup>*Advanced Research Center for Nanolithography (ARCNL), The Netherlands  
b.brewster@arcnl.nl*

The unique emission spectrum of tin ions in charge states  $\text{Sn}^{8+}$ – $\text{Sn}^{14+}$  produces radiation in a narrow region around 13.5 nm which is used for print patterns silicon wafers to make computer chips. To efficiently print narrower features, future shorter-wavelength printing techniques would benefit from a similarly narrow emission spectrum. The experimentally observed narrow spectrum is only reproduced theoretically when configuration interaction is included. To predict other element–charge-state combinations that may exhibit this effect, we investigate the physical origin of spectral narrowing induced by configuration interaction.

The Hamiltonian matrix elements  $H_{bb'}$  are sums of products of radial integrals and angular factors. The radial integrals include the electrostatic (Coulomb) direct integrals  $F^k$ , the exchange integrals  $G^k$ , and the spin–orbit parameter  $\zeta$ , all obtained by numerical integration. The angular coefficients arise due to the coupling between subshells and depend on the quantum numbers  $l$  and  $m_l$  of the interacting basis states  $b$  and  $b'$ . The Hamiltonian matrix elements are constructed following Cowan [1], Eq. (12.47).

$$H_{bb'} = \delta_{bb'} E_{av} + \sum_{l_i, l_j, k} f_k^{bb'}(l_i l_j) F^k(l_i l_j) + d_j^{bb'} \zeta_j + g_k^{bb'}(l_i l_j) G^k(l_i l_j) \quad (1)$$

The calculation of the energy levels contributing to the emission spectrum requires assembling and diagonalising a large sparse Hamiltonian matrix. Inclusion of triply, quadruply, and quintuply excited configurations is necessary [2, 3] but can increase the matrix dimension substantially (exceeding  $10^6$  is within the scope of this project). To address this computational challenge, we aim to perform the diagonalization on graphical processing units (GPUs) to achieve a speed-up of up to a factor fifty [4]. Ultimately, this large-scale atomic-structure modelling is aimed at isolating the physical mechanism by which configuration interaction leads to spectral narrowing.

By resolving how extensive configuration mixing redistributes line strength in highly charged tin ions, we seek general criteria that explain why narrowing occurs in  $\text{Sn}^{8+}$ – $\text{Sn}^{14+}$  and to identify other element–charge-state combinations that may exhibit similarly favourable emission characteristics for future shorter-wavelength lithography.

- [1] R. D. Cowan, *The Theory of Atomic Structure and Spectra* (University of California Press, Berkeley, 1981).
- [2] J. Sheil, O. O. Versolato, A. J. Neukirch, and J. Colgan, *J. Phys. B: At. Mol. Opt. Phys.* **54**, 035002 (2021).
- [3] F. Torretti, J. Sheil, R. Schupp *et al.*, *Nat. Commun.* **11**, 2334 (2020).
- [4] Inductiva Research Team, “CPU vs GPU Eigensolver Benchmark,” Inductiva Blog (2023).

## Synchrotron calibration of a transmission grating spectrometer for broadband EUV measurements

**E.O. Chebykin<sup>1</sup> A.E.H. Gaballah<sup>1</sup>, R. van der Horst<sup>2</sup>, M.D. Ackermann<sup>1</sup>, M. van de Kerkhof<sup>2</sup> and M. Bayraktar<sup>1</sup>**

<sup>1</sup> *XUV Optics Group, University of Twente, Enschede, The Netherlands*

<sup>2</sup> *ASML Research, Veldhoven, The Netherlands*

Laser-produced plasma (LPP) of tin is used as the light source in the latest generation extreme ultraviolet (EUV) nanolithography applications. The plasma dominantly emits light in the spectral range around 13.5 nm [1]. In addition to 13.5 nm band, emission in other spectral ranges, including vacuum and deep ultraviolet (DUV) ranges are also present in the spectra. The longer wavelength emission may cause creation of secondary hydrogen plasma, hence interacting with the surfaces of the scanner [2]. A broadband characterization of the EUV emitting LPP is therefore necessary. Using a transmission grating spectrometer (TGS) [3] is found to be useful in such broadband characterization due to ease of alignment in using different gratings for covering different spectral ranges. Besides the gratings, the TGS consists of filters, slits, a photodiode and a charge-coupled device. These components have different spectral efficiencies needing individual calibrations for interpreting the plasma emission spectra.

Here, we present the calibration of the TGS that has been performed at Elettra synchrotron facility in Trieste, Italy. The calibration covers the 5-400 nm range such that absolute intensity measurements of EUV light sources can be done in the needed EUV to DUV range.

[1] S.K. Moore, EUV lithography finally ready for fabs, IEEE Spectr. 55(1), 46(2018)

[2] Beckers, et al, Applied Sciences, 9(14), 2827 (2019).

[3] M. Bayraktar et al., NEVAC blad 54(1), 14 (2016)

# A Python-based Hybrid Atomic Structure Collisional-Radiative Model for Spectroscopic Plasma Diagnostics and Equation-of-state Computations

**Ryan R. Childers**

*U.S. Naval Research Laboratory, Washington, D.C. USA*

*Joint ICTP-IAEA School on Atomic and Molecular Processes in Plasmas*

*May 4 – 8, 2026, Trieste, Italy*

Collisional-radiative modelling is a powerful predictive utility for diagnosing the physical parameters of plasmas. The accuracy of a collisional-radiative model relies on the *detail* (of the energy levels and their level refinement) and *completeness* (of the excited state levels and the processes coupling them) of the atomic data underpinning the collisional-radiative processes. These components are inherently competitive; achieving statistical completeness with highly detailed atomic data requires tracking an exceedingly large number of energy levels that rapidly become computationally intractable in, for example, mid-to-high  $Z$  elements. Efforts to balance the detail, completeness, and tractability of an atomic model have shown promise when employing “hybrid” level averaging techniques [1-3], which selectively blend highly detailed levels with ensemble-averaged level arrays that can comprise hundreds of detailed states in a single array. A hybrid structure collisional-radiative atomic model [4] has been developed for diagnostic application to plasma experiments and equation-of-state tabulation at the U.S. Naval Research Laboratory. The model blends sets of atomic data calculated with varying degree of detail using the Flexible Atomic Code [5] to achieve high-fidelity predictions of plasma ionization and x-ray radiation, while maintaining computational tractability through selective implementation of statistical averaging techniques to the energy level data. The coupled collisional-radiative rates problem is solved using a mixture of detailed and ensemble-averaged energy levels in collisional-radiative (steady-state) equilibrium to generate level population densities. Radiative emissivities and synthetic spectra are computed from the population solution, while key equation-of-state parameters are derived from the combined populations and emissivities, including charge state distribution, K- and L-shell radiative powers, and line opacities – determined from an iterative probability-of-escape formalism [6]. Theoretical validation is carried out against the state-of-the-art spectroscopic collisional-radiative model SCRAM [2], which also employs a hybrid structure technique for the collisional-radiative problem. Comparison between the two hybrid models is performed for argon and krypton elements, looking at predictions of ionization balance, average ion charge, fractional ion abundance, L- and K-shell synthetic spectra, and probability-of-escape factors.

This work was supported by the NRL Karle Fellowship and in part by the DOE NNSA.

---

[1] O. Peyrusse, *J. Phys. B* **33**, 4303 (2000)

[2] S. B. Hansen *et al.*, *High En. Dens. Phys.* **3**, 109 (2007)

[3] Q. Porcherot *et al.*, *High En. Dens. Phys.* **7**, 234 (2011)

[4] R. Childers DTIC Ref TBD

[5] M.F. Gu, *Can. J. Phys.* **86**, 675 (2008)

[6] J. P. Apruzese, *J. Quant. Spectrosc. Radiat. Transfer* **25**, 419 (1981)

## A New Approach for Ground and Excited State Electron-Impact Ionization of C I within Magnetically Confined Plasmas

David J. Dougan<sup>1</sup>, Connor P. Ballance<sup>1</sup>, Catherine A. Ramsbottom<sup>1</sup> and Stuart D. Loch<sup>2</sup>

<sup>1</sup>*Astrophysics Research Centre, Queen's University Belfast, Belfast, BT7 1NN, Northern Ireland, United Kingdom*

<sup>2</sup>*Department of Physics, Auburn University, Auburn, 36832, AL, United States of America*

In magnetically-confined plasmas the temperatures and densities are such that the effective ionization rate may be dominated by the contributions from the metastable ionization rates rather than the ground state [1]. Sophisticated approaches, such as Convergent Close-Coupling (CC) and Time-Dependent Close-Coupling (TDCC), have often focused on the comparison of ground state ionization with experimental data [2, 3], in contrast to comprehensive ionization from all excited states required for collisional-radiative modelling. We update the electron-impact ionization and excitation data available for neutral carbon (C I). Carbon embodies many favourable characteristics required by Plasma Facing Components within tokamaks, in terms of erosion, redeposition, H retention and neutron damages [4]. This is due to its simple electron structure and excellent temperature characteristics relative to its low atomic number [5].

We expand upon the  $R$ -matrix with Pseudostates (RMPS) methodology and incorporate it more generally into the Dirac Atomic  $R$ -matrix Codes (DARC) [6], to allow both ionization and excitation rates to be calculated from the same initial target structure. This is a generalization of the work of Badnell (2008) [7] which considered only a neutral hydrogen system. The cross sections from two distinct C I targets show good agreement with experimental data and other theoretical calculations for the C I ground level. However, this is not our primary focus, given the agreement of previous calculations with existing experimental data [8]. Rather, the DARC  $R$ -matrix with Pseudostates (DRMPS) method allows for the first level-resolved excited state ionization cross sections to be calculated. In addition, our level-resolved excitation collision strengths are also in alignment with other theoretical calculations.

The availability of both electron-impact excitation and ionization rates allows for the determination of effective ionization rates through collisional radiative calculations. We show that for the temperatures and densities applicable in magnetically confined plasmas, the ionization contributions from metastable levels have a non-trivial effect on the effective ionization rate coefficients of C I. These results provide confidence in our new DRMPS approach, which will now make ionization cross sections, both ground and excited, more readily available for non-hydrogenic systems.

- [1] J. P. Allain, D. G. Whyte & J. N. Brooks, *Nucl. Fusion* **44**, 655 (2004)
- [2] M. S. Pindzola, J. Colgan, F. Robicheaux & D. C. Griffin, *Phys. Rev. A* **62**, 042705 (2000)
- [3] Sh. A. Abdel-Naby, C. P. Ballance, T. G. Lee et al., *Phys. Rev. A* **87**, 022708 (2013)
- [4] T. Tanabe, *Plasma Phys. Rep.* **45**, 300 (2019)
- [5] A. Miyahara & T. Tanabe, *J. Nucl. Mater.* **155-157**, 49 (1988)
- [6] C. P. Ballance, DARC  $R$ -matrix codes, <https://connorb.freeshell.org/> (2025)
- [7] N. R. Badnell, *J. Phys. B: At. Mol. Opt. Phys.* **41**, 175202 (2008)
- [8] N. Bachi, S. Otranto & K. Tórkési, *Atoms* **11**, 2 (2023)

# Modeling Atomic Processes in Fusion Plasmas

Lamya El Mesbahi

University of Debrecen, Hungary

**Abstract:** Understanding atomic processes is essential for accurately describing the behavior of magnetically confined fusion plasmas. In this work, we investigate key atomic interactions—including excitation, ionization, recombination, and radiation emission—under conditions relevant to fusion devices. Using state-of-the-art theoretical models and available atomic databases, we aim to improve the reliability of collisional-radiative descriptions and spectral diagnostics for high-temperature plasmas. The study focuses on the role of atomic rate coefficients, population modelling, and spectral line formation, with particular emphasis on their impact on plasma parameters such as electron density, temperature, and charge-state distributions. The results contribute to developing more accurate modelling tools that support the interpretation of spectroscopic measurements in fusion experiments. This work forms an essential part of my PhD research on the modelling of atomic processes in fusion plasma and aims to advance our understanding of atomic-level mechanisms that govern plasma behavior in fusion-relevant environments.

# Atomic Structure Calculations and identification of EUV and FUV spectral lines of $W^{10+}$

Falga Yadav<sup>1</sup>, Arun Goyal<sup>2</sup>, Narendra Singh<sup>2</sup>

<sup>1</sup> *Department of Physics & Astrophysics, Delhi University, Delhi-110007, India*

<sup>2</sup> *Zakir Husain Delhi College, Delhi University, Delhi-110002, India*

In the present work, a comprehensive spectroscopic investigation of the tungsten  $W^{10+}$  ion is carried out to determine its atomic structure and radiative properties of relevance to fusion and astrophysical plasmas. Relativistic atomic structure calculations are performed using the Flexible Atomic Code (FAC) and the General-purpose Relativistic Atomic Structure Package (GRASP), incorporating extensive configuration interaction effects. Energy levels of the lowest excited configurations are calculated and systematically analyzed. Radiative transition parameters are evaluated for electric dipole (E1), electric quadrupole (E2), magnetic dipole (M1), and magnetic quadrupole (M2) transitions, primarily in the extreme ultraviolet (EUV) and Far ultraviolet (FUV) spectral region. The calculated data are critically compared with available theoretical and experimental results, showing good agreement and improved accuracy. The present results provide a reliable and self-consistent set of atomic data for  $W^{10+}$ , which are essential for plasma diagnostics, opacity modeling, and radiation transport studies in magnetic confinement fusion devices.

## The Challenge of Accurate Recombination Data for Lanthanides and Actinides

N.Ferguson<sup>1</sup>, and M. G. O'Mullane<sup>1</sup>

<sup>1</sup> *Department of Physics, University of Strathclyde, Glasgow G4 0NG, United Kingdom.*

Interpreting kilonova spectra from neutron-star mergers requires accurate atomic data for the heavy r-process elements synthesized in these events. From a plasma-physics perspective, this problem is closely connected to population modelling and collisional–radiative (CR) modelling in complex, low-density, transient plasmas, where departures from local thermodynamic equilibrium (LTE) are significant. Reliable modelling therefore demands self-consistent non-LTE level populations, ionization balance, and emissivities derived from detailed atomic data rather than equilibrium assumptions. These requirements pose a major challenge for lanthanides and actinides, whose open f-shell configurations produce extremely dense level structures and strong sensitivity to atomic structure approximations. Such ions represent some of the most complex species encountered in CR modelling, with atomic data uncertainties propagating directly into predicted charge-state distributions and spectral modelling. I present recent progress in the production and benchmarking of non-LTE atomic data for key heavy f-shell elements, with a focus on neodymium (Nd), uranium (U), and related species. New atomic-structure calculations are used to compute radiative and dielectronic recombination rates, as well as direct excitation data, suitable for population and CR modelling under kilonova ejecta conditions. For open f-shell ions, the resulting atomic structure is found to be highly sensitive to the choice of orbital scaling parameters, with even minor variations leading to significant changes in level ordering and transition probabilities. This sensitivity is particularly evident in the determination of ground configurations and terms. As a case study, Nd VI exhibits extreme sensitivity: two competing ground-state configurations reported in the literature [1] can both be reproduced within AUTOSTRUCTURE [2] by adjusting scaling parameters at the level of the third decimal place. Small structural differences translate into substantial spreads in predicted recombination rates, with direct consequences for CR population modelling and non-LTE spectral synthesis. These results highlight both the challenges and the necessity of robust atomic data for heavy f-shell elements, not only for interpreting kilonova spectra but also for advancing plasma population modelling in regimes where atomic complexity and non-equilibrium effects dominate.

[1] F. Goryaev, I. Tupitsyn, *Atomic Data and Nuclear Data Tables* **160**, 101673 (2024).

[2] N.R. Badnell, *Computer Physics Communications* **182**, 1528 (2011).

## Atomic and Molecular Opacity in Low-Temperature Plasmas: Quantifying Uncertainty in Atomic and Plasma Physics Models

Aaron Forde<sup>1</sup>, Isuru Ariyaratna<sup>1</sup>, Jeffery A. Leiding<sup>1</sup>, Chris Fontes<sup>2</sup>, Amanda J. Neukirch<sup>1</sup>, James Colgan<sup>1</sup>, Eddy Timmermans<sup>2</sup>, Mark Zammit<sup>1</sup>

<sup>1</sup>Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, United States

<sup>2</sup>Computational Physics Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, United States

### ABSTRACT

Radiative properties of heated air species are vital ingredients in detecting, modeling, and characterizing transport. This information can be utilized to describe the radiative energy transfer, collisional-radiative, radiation-hydrodynamic, and plasma kinetic models which are utilized to model spectra, energy transfer, and bulk-plasma transport in media such as the atmosphere. Ambient air at sea-level is composed primarily of N<sub>2</sub> and O<sub>2</sub>, at 78% and 21%, respectively, with contributions from H<sub>2</sub>O, Ar, CO<sub>2</sub>, Ne, He, CH<sub>4</sub>, Kr, and N<sub>2</sub>O. The composition of air can change when heated into a mixture of molecular and atomic species. Historically, the ATOMIC opacity software package developed at Los Alamos National Laboratory has implemented efficient approximations to the atomic physics (intermediate-coupling), plasma physics, and atomic-plasma interactions appropriate for high-temperature and high-density regimes used to generate the OPLIB tables.<sup>1</sup> These approximations breakdown for low-temperature plasmas containing neutral and near-neutral species. To address this we use improved atomic electronic structure with a selective-CI approach<sup>2</sup> and a semi-empirical ‘sum-over-states’ approach for the atom-electron scattering for improved line-broadening.<sup>3</sup> The error in gray-mean opacities are reported and the improved opacities are tested on a benchmark 1-D radiation hydrodynamics simulation.

(1) Colgan, J.; Kilcrease, D. P.; Magee, N. H.; Sherrill, M. E.; Abdallah Jr, J.; Hakel, P.; Fontes, C. J.; Guzik, J. A.; Mussack, K. A. A NEW GENERATION OF LOS ALAMOS OPACITY TABLES. *The Astrophysical Journal* **2016**, 817 (2), 116. DOI: 10.3847/0004-637X/817/2/116.

(2) Colgan, J.; Kilcrease, D. P.; Abdallah, J.; Sherrill, M. E.; Fontes, C. J.; Hakel, P.; Armstrong, G. S. J. Atomic structure considerations for the low-temperature opacity of Sn. *High Energy Density Physics* **2017**, 23, 133-137.

(3) Dimitrijević, M. S.; Konjević, N. Stark widths of doubly- and triply-ionized atom lines. *Journal of Quantitative Spectroscopy and Radiative Transfer* **1980**, 24 (6), 451-459.



# Temporal and Spatial evolution of the ion temperature in the WEST tokamak

P. Forestier-Colleoni<sup>1\*</sup>, O. Peyrusse<sup>2</sup>, G. Colledani<sup>1</sup>, C. Desgranges<sup>1</sup>, R. Guirlet<sup>1</sup>, D. Guibert<sup>1</sup>, P. Maget<sup>1</sup>, P. Manas<sup>1</sup>, G. Miglionico<sup>1</sup>, R. Dumont<sup>1</sup>, on behalf of the WEST team<sup>1,3</sup>

<sup>1</sup> CEA, IRFM, F-13108 St-Paul-Lez-Durance, France  
<sup>2</sup> Aix-Marseille Université, CNRS, Laboratoire LP3, UMR7341, F-13288 Marseille, France  
<sup>3</sup> http://west.cea.fr/WESTTeam  
 \* pierre.forestier-colleoni@cea.fr

## Ion temperature a crucial parameter for achieving fusion

The WEST tokamak is a device in which D-D plasmas are being produced to investigate various aspects related to the operation of future devices[1]. The typical plasma density and temperature reachable on this device is  $n_e = 10^{19}\text{-}10^{20} \text{ m}^{-3}$  and  $T_e = 3 \text{ to } 8 \text{ KeV}$ .

The ability to measure spatial and temporal variations in the ion temperature is crucial for:


- improving plasma confinement
- optimizing fusion reactions.

Photo of the WEST tokamak

### How to measure the temporal and spatial evolution of the plasma ion temperature ?


## He-like argon impurity emission suitable for determining ion temperature

- Spectral range from 3100 to 3150 eV is convenient to clearly distinguish the He-like lines of argon.



Argon He-like emission spectrum from collisional radiative model for  $n_e = 10^{20} \text{ m}^{-3}$  and  $T_e = 1500 \text{ eV}$  [3].

- Line FWHM is directly related to ion temperature.



$$\frac{\Delta E_{FWHM}}{E_0} \sim \sqrt{\frac{8 \ln 2 k T_i}{m_i c^2}}$$

with :


- $kT_i$  the ion temperature.
- $\Delta E_{FWHM}$  the Full Width at Half Maximum of the spectral line
- $E_0$  the central wavelength of the spectral line
- $m_i$  the ion mass

## The X-ray Imaging Crystal Spectrometer designed for retrieving the ion temperature

- A Spherical Bragg crystal in Johann configuration to obtain the spectral resolution as well as spatial resolution [4].


Johann configuration imaging the collected photon on poloidal plan (top) and diffract them on the toroidal plan by Bragg law (bottom).

- 2d map of the collected photon emitted by the plasma reveal  $K\alpha$  spectral lines (shot #61530)



Intensity map of a WEST shot (#61530) (left). Spectra obtained at 3 relative Z positions (left).

- Temporal and spatial evolution of the ion temperature give valuable information on the plasma evolution i.e. increase of  $T_i$  while N2 injection (X-Point Radiator).



- Comparison with Ti-Neutron flux estimation for #60359.

References

1 J. Bucalossi, et al. WEST full tungsten operation with an ITER grade divertor. *Nucl. Fusion* **64**, 112022 (2024).  
 2 J. Wesson and D.J. Campbell. Tokamaks, volume 149. Oxford University Press, (2011).  
 3 O. Peyrusse, A model for the simulation of nonequilibrium line transfer in laboratory plasmas *Physics of Fluids B: Plasma Physics* (1989-1993) **4**, 2007 (1992).  
 4 A. DaRos et al. Electron and ion temperature measurement with a new x-ray imaging crystal spectrometer on WEST *Rev. Sci. Instrum.* **95**, 043505 (2024).

## Theoretical study of electron-molecule collisions for plasma simulations

**Alejandro L. García-Muñoz**<sup>1,2</sup>, Nicolas Sisourat<sup>1</sup>, Zdeněk Mašín<sup>2</sup>

<sup>1</sup> Sorbonne Université, CNRS, Laboratoire de Chimie Physique Matière et Rayonnement, UMR 7614, F-75005 Paris, France

<sup>2</sup> Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 180 00 Praha 8, Czech Republic

The development of low-power electric propulsion thrusters [1] is a crucial step to respond to the growing demand for small satellite technology. In this context, iodine is a promising candidate to replace the currently used propellants (e.g. xenon), for both technological and economic reasons [2]. Indeed, in 2019 the first satellite propelled by iodine was launched into space [3].

However, the physics and the chemistry of iodine low-temperature plasma generated in the thruster are not well understood, due to the lack of data on the elementary electronic processes occurring. Moreover, electron impact collisional processes, and their associated reaction cross sections, are critical inputs required for plasma modeling. In this context, we have studied the electronic processes in electron-iodine molecule collisions up to 100 eV using the R-matrix method [4] combined with Effective Core Potentials [5] for the description of the core electrons.

### REFERENCES

- [1] Goebel, Dan M., Ira Katz, and Ioannis G. Mikellides. Fundamentals of electric propulsion. John Wiley & Sons, 2023.
- [2] Jia-Richards, Oliver, and Trevor Lafleur. "Iodine electric propulsion system thrust validation: from numerical modeling to in-space testing." *Journal of Propulsion and Power* 39.6 (2023): 896-904.
- [3] Rafalskyi, Dmytro, et al. "In-orbit demonstration of an iodine electric propulsion system." *Nature* 599.7885 (2021): 411-415.
- [4] Tennyson, Jonathan. "Electron–molecule collision calculations using the R-matrix method." *Physics Reports* 491.2-3 (2010): 29-76.
- [5] Dolg, Michael, and Xiaoyan Cao. "Relativistic pseudopotentials: their development and scope of applications." *Chemical reviews* 112.1 (2012): 403-480.

# Dirac R-matrix electron-impact excitation of Ni I for the interpretation of magnetically-confined/astrophysical plasmas

John J. Henry<sup>1</sup> and Connor P. Ballance<sup>1</sup>

<sup>1</sup>*Astrophysics Research Centre, Queen's University of Belfast, Belfast, BT7 1NN,  
Northern Ireland, United Kingdom*

Interpreting the spectra from magnetically-confined/astrophysical plasmas involves generating synthetic spectra from collisional-radiative models at required densities and temperatures. These models are underpinned by accurate atomic and collisional rates, ideally employing the same atomic structure. For example, within an astrophysical context, near-neutral ion stages of Nickel (Ni) are required for non-local thermodynamic equilibrium of the nebular phase of supernovae [1].

Collisional rates for near neutral targets, benefit from non-perturbative approaches such as the Dirac R-matrix (DARC) or Dirac R-Matrix with Pseudo-states approach (DRMPS). Initially, relativistic atomic orbitals are calculated within a Multi-Configuration-Dirac-Fock (MCDF) approximation employing a modified version of the General Relativistic Atomic Structure Package (Grasp0). Where available, Energy levels and A-values were compared with experimental values and in general found good agreement. Subsequently, a relativistic electron-impact excitation was then performed (DARC) [2] producing collision strengths that have been Maxwellian convolved for implementation within collisional-radiative codes. Some of our results shall be presented.

## References :

- [1] F. P. Callan, A. Holas, J. Morán-Fraile & S. A. Sim. *A&A* **702** A29 (2025)
- [2] C. P. Ballance, DARC R-matrix codes, <https://connorb.freeshell.org/> (2025)

## Effect of pulsed neutral gas injection on heat flux scrape-off layer (SOL) width in ADITYA-U tokamak

**SK Injamul Hoque<sup>1,2</sup>, Harshita Raj<sup>1,2</sup>, Ritu Dey<sup>3</sup>, Soumitra Banerjee<sup>1,2</sup>, Ruchi Varshney<sup>1,2</sup>, Komal<sup>1,2</sup>, Kaushlender Singh<sup>1,2</sup>, Suman Dolui<sup>1,2</sup>, Ankit Kumar<sup>1,2</sup>, Ashok Kumawat<sup>1,2</sup>, Bharat Hegde<sup>1,2</sup>, Sharvil Patel<sup>4</sup>, Kiran Patel<sup>1</sup>, Rohit Kumar<sup>1</sup>, Suman Aich<sup>1,2</sup>, Pramila Gautam<sup>1</sup>, Umesh Nagora<sup>1,2</sup>, Asha N Adhiya<sup>1</sup>, K. A. Jadeja<sup>1</sup>, K.M. Patel<sup>1</sup>, Ankit Patel<sup>1</sup>, R.L. Tanna<sup>1</sup>, and Joydeep Ghosh<sup>1,2</sup>**

<sup>1</sup>*Institute for Plasma Research, Gandhinagar, India*

<sup>2</sup>*Homi Bhabha National Institute, Mumbai, India*

<sup>3</sup>*Department of Physics, Indian Institute of Technology Tirupati, Yerpedu, India*

<sup>4</sup>*University of Virginia, Department of Physics, Charlottesville, USA*

Handling the extremely high heat fluxes, often reaching  $\sim 10 \text{ MW m}^{-2}$ , on plasma-facing components is a major challenge for future tokamak reactors. The heat-flux scrape-off layer width,  $\lambda_q$ , represents how rapidly the parallel heat flux decays radially in the SOL and is a key parameter that determines divertor and limiter heat loads. When  $\lambda_q$  is very narrow, the power is deposited over a small area, leading to severe localized thermal stress. Plasma detachment produced by impurity seeding is a widely used method to reduce divertor heat flux, but it often comes at the cost of degraded core confinement [1]. Achieving a detached edge while simultaneously maintaining or improving core performance is therefore essential for reactor-relevant operation. Experiments on devices such as TCV, ASDEX Upgrade, and DIII-D have previously investigated this challenge using advanced divertor configurations [2] and real-time impurity-seeding control techniques [3,4].

In this study, we present results showing that short neutral gas-pulse injection can simultaneously broaden the heat-flux SOL width, reduce edge heat loads, and improve core confinement. A comparative analysis further reveals that pulsed fuelling is significantly more effective than continuous gas fuelling in increasing  $\lambda_q$ , reducing edge power deposition, and enhancing core density and temperature. An analytical model is developed to explain how ionization and recombination of neutrals in the ADITYA-U edge region influence the heat-flux width  $\lambda_q$ . Further, simulations with UEDGE code reproduce the key transport features as predicted through analytical model. These findings suggest that short neutral gas-pulse injection provides an effective approach to reducing and broadening heat loads on the limiter or divertor, while simultaneously maintaining or even improving core confinement in fusion plasmas.

[1] A. Kallenbach et al. Nuclear Fusion, vol. 55, no. 5, May 2015,

[2] H. Raj et al. Nuclear Fusion, vol. 62, no. 12, Dec. 2022

[3] O. Gruber et al., 1995. PHYSICAL REVIEW LETTERS vol. 74, no 21

[4] H. Q. Wang et al. Phys Plasmas, vol. 28, no. 5, May 2021

# Theoretical Advances in Electron Molecular-ion Scattering, Method Benchmarking and Connecting Collisional and Photofragmentation Resonances

**Dávid Hvizdoš<sup>1</sup>, Chris H. Greene<sup>2</sup>, Roman Čurík<sup>3</sup>, and Ioan F. Schneider<sup>1</sup>**

<sup>1</sup>*Laboratoire Ondes et Milieux Complexes, UMR-6294 CNRS and Université Le Havre Normandie, 25 rue Philippe Lebon, BP 540, 76058, Le Havre, France*

<sup>2</sup>*Department of Physics and Astronomy and Purdue Quantum Science and Engineering Institute, Purdue University, West Lafayette, Indiana 47907 USA*

<sup>3</sup>*J. Heyrovský Institute of Physical Chemistry, ASCR, Dolejškova 3, 18223 Prague, Czech Republic*

Electron scattering off of molecular ions is an important and complicated topic for plasma chemistry studies, especially when talking about plasma neutralization which is dominated by dissociative recombination both in astrochemical (interstellar medium) and plasma reactor (fusion edge plasma) scenarios. In the past few years we've seen a substantial evolution in the theoretical methodology describing these processes.

We present here mainly the methodological improvements of the energy dependent frame transformation (EDFT) method [1, 2] which can now handle the branching ratios of individual dissociation channels as well as rotational degrees of freedom (neglected in most previous EDFT studies). We show several benchmarks, using a simplified model of Hydrogen and comparing with both more “brute force” solutions [3] and other contemporary frame transformation methods such as the step-wise multichannel quantum defect theory [4].

We also show our latest developments [5] coupling the scattering to photofragmentation and how the resonant features in the cross sections compare between the processes.

[1] H. Gao, C. H. Greene, *Phys. Rev. A* **42**, 6946 (1990).

[2] D. Hvizdoš, R. Čurík, C. H. Greene, *Phys. Rev. A* **111**, 012805 (2025).

[3] R. Čurík, D. Hvizdoš, C. H. Greene, *Phys. Rev. A* **98**, 062706 (2018).

[4] O. Motapon, N. Pop, F. Argoubi, J. Zs Mezei, M. D. Epee Epee, A. Faure, M. Telmini, J. Tennyson, and I. F. Schneider, *Phys. Rev. A* **90**, 012706 (2014).

[5] D. Hvizdoš, R. Čurík, C. H. Greene, *Mol. Phys.*, *soon to be published*.

**T.K.Nurubeyli<sup>1,2</sup> N.Sh.Jafar<sup>1</sup>**

<sup>1</sup>*Institute of Physics of the Ministry of Science and Education of the Republic of Azerbaijan,  
H.Javid avenue 131, Baku AZ-1073, Azerbaijan, Baku [t.nurubeyli@physcis.science.az](mailto:t.nurubeyli@physcis.science.az),  
[nurlanajafarr@gmail.com](mailto:nurlanajafarr@gmail.com)*

<sup>2</sup>*Azerbaijan State Oil and Industry University,  
Azadlig avenue 34, Baku AZ-1010, Azerbaijan, Baku [omartarana@gmail.com](mailto:omartarana@gmail.com)*

In this work, plasma–matrix interaction mechanisms in radio-frequency inductively coupled plasma (ICP) were investigated with emphasis on the physical processes governing ion formation, ionization equilibrium, and ion transport in ICP-MS under the presence of complex biological matrices. The study focuses on how matrix-induced perturbations of plasma energetics affect ionization efficiency and mass-spectrometric signal formation.

Spectral interferences were shown to originate predominantly from plasma-driven formation of polyatomic, molecular, and isobaric ions, reflecting ion–molecule and charge-transfer reactions occurring in high-temperature RF plasma. Non-spectral matrix effects arise from the interaction of organic components, salt residues, and acid-derived species with the plasma energy balance, leading to modifications of electron heating efficiency, ionization kinetics, and ion extraction conditions.

Isotopes of Mg, Ni, Cu, Zn, Ga, As, Se, Sb, Pt, and Pb were examined to evaluate plasma-induced distortions of ion populations. Pronounced spectral interference was observed for specific isotopes, such as <sup>63</sup>Cu and <sup>69</sup>Ga, due to enhanced formation of overlapping plasma-generated species. Selection of alternative isotopes with reduced sensitivity to plasma-driven interference significantly stabilized ion signals under optimized plasma conditions.

Non-spectral matrix effects were found to correlate strongly with elemental ionization potentials and shifts in ionization equilibrium within the plasma. Variation of RF generator power demonstrated its central role in controlling electron heating and ionization efficiency: reduced RF power suppressed analyte ionization, whereas excessive power enhanced the formation of doubly charged ions, leading to altered mass-to-charge distributions. An optimal RF power of 1430 W provided a stable compromise between ionization efficiency and suppression of multi-charged ion formation.

Optimization of nebulizer gas flow (1.00 L min<sup>-1</sup>) and sampling depth (8 mm) revealed the influence of the axial zonal structure of the ICP on ion formation and transport. Additional optimization of sample uptake rate and ion-optical extraction potentials resulted in enhanced ion transmission efficiency.

The results demonstrate that effective compensation of biological matrix effects in ICP-MS requires plasma-physics-guided optimization of RF plasma parameters, enabling controlled ionization conditions and stable ion transport for reliable mass-spectrometric detection.

- [1]. Nurubeyli T.K., Jafar N.Sh, Ahmadova, Kh.N. *Methods and Objects of Chemical Analysis* 20(1), 39–45 2025
- [2]. Nurubeyli, T.K., Jafar, N.S., Mammadova, G.N. *International Journal of Mass Spectrometry* 507, 117355, 2025
- [3]. T.K. Nurubeyli, A.M. Hashimov, N.S. Jafar, “Optimization of analytical accuracy in analysis of metals in biological samples by ICP-MS using an artificial matrix” *International Journal on Technical and Physical Problems of Engineering*, vol. 17, № 1, pp. 59–68, 2025

## Irradiation-driven fragmentation of $W(CO)_6$ in gas phase: Analysis of scaling behaviour

Soumyo Kheto<sup>1</sup>, Alexey Verkhovtsev<sup>2</sup>, Bobby Antony<sup>1</sup>, Andrey solov'yov<sup>2</sup>

<sup>1</sup>*Department of Physics, IIT(ISM) Dhanbad, 826004 Dhanbad, Jharkhand, India*

<sup>2</sup>*MBN Research Center, Altenhöferallee 3, 60438 Frankfurt am Main, Germany*

Irradiation-Driven Molecular Dynamics (IDMD)[1] is an emerging multiscale simulation methodology that bridges quantum chemistry, molecular dynamics, and radiation physics to study how energetic particles interact with complex molecular systems. Unlike conventional MD approaches, IDMD incorporates stochastic irradiation events—such as bond cleavage, ionization, and secondary electron generation—directly into the dynamical evolution of molecules and materials. And one of the promising applications of this methodology is focused electron beam induced deposition (FEBID)[2], a direct-write nanofabrication technique under different irradiation conditions. This makes it uniquely suited for investigating processes where radiation plays a decisive role, including the fragmentation of organometallic precursors, radiation damage in biomolecules, and the formation of clusters or nanoparticles. Implemented in the advanced computational platform, MBN Explorer[3], IDMD enables the integration of experimental cross-section data with large-scale simulations to offer mechanistic perspectives on both atomistic and mesoscopic levels. In this upcoming school, the scaling dependencies on the fragmentation process of  $W(CO)_6$  molecules in the gas phase under different irradiation conditions will be presented. The outcome of the work is expected to deepen our understanding of radiation-driven chemistry and open avenues for predictive modelling of nanostructure growth, offering direct relevance to nanofabrication, materials science, and radiation biology.

- [1] A.V. Verkhovtsev, I.A. Solov'yov, and A.V. Solov'yov, Irradiation-driven molecular dynamics: a review, *Eur. Phys. J. D*, vol. 75, 213 (2021)
- [2] Huth, M., et al. (2012). “Focused electron beam induced deposition: A perspective.” *Beilstein Journal of Nanotechnology*, 3, 597–619.
- [3] I. A. Solov'yov, A. V. Yakubovich, P. V. Nikolaev, I. Volkovets, and A. V. Solov'yov, MBN Explorer: A Universal Program for Multiscale Computer Simulations of Complex Molecular Structure and Dynamics. *J. Comput. Chem.*33, 2412–2439 (2012).

## Spectroscopic Time-of-Flight Spatial Distribution Measurements of Mono- and Polycrystalline W with Varying Surface Properties in Ne and Ar Plasma at the Linear Plasma Device PSI-2

M. Klein<sup>1</sup>, O. Marchuk<sup>1</sup>, M. Sackers<sup>1</sup>, A. Kreter<sup>1</sup> and S. Brezinsek<sup>1,2</sup>

<sup>1</sup>*Forschungszentrum Jülich GmbH, Institute of Fusion Energy and Nuclear Waste Management, Plasmaphysik (IFN-1), Germany*

<sup>2</sup>*HHU Düsseldorf, Faculty of Mathematics and Natural Sciences*

Tungsten (W) is the plasma-facing material (PFM) for future fusion reactors, such as ITER. Its erosion by the plasma and subsequent lifetime is of great interest. Additionally, this sputtered W can enter the plasma and cool it through radiation losses. The quantity and spatial distribution of sputtered W from PFMs therefore has a direct impact on reactor performance: The erosion can be modelled by spectroscopic investigation of W and W<sup>+</sup> or re-deposition processes [1]. Therefore, monitoring the W flux is necessary for operating the reactor, and utilising optical spectroscopy is an important tool to do so. For energy impacts in the keV range, the sputtered W atoms follow a cosine angular distribution [2]. For lower energies, as typically exist in the divertor region, experimental data shows a discrepancy to existing models [3]. This can be investigated at the linear plasma device PSI-2. Low-temperature plasmas in the eV range can be studied with densities up to  $n_e \approx 10^{18} \text{ m}^{-3}$ .

In this experiment, polished poly- and monocrystalline (100, 111) W is exposed to a linear Neon and Argon plasma at kinetic energies of about 100 eV. Additionally, W fuzz on a polycrystalline substrate is examined. The W atoms sputtered into the Ne/Ar plasma column allow for the acquisition of line shapes in a spectroscopic time-of-flight (ToF) measurement; multiple lines can be acquired due to further excitations in the plasma beam. The total sputtering yield can be deduced from multiple acquisitions. In this ToF measurement it is possible to differentiate between angular distributions of sputtered W flux, while all other experimental parameters stay fixed. Multiple spectral lines are investigated for cross-reference in the 400-500 nm region.

These new experimental results show deviations in angular distributions between lattice structures and can be used to support modelling of low temperature plasma-wall interactions. In addition, the surface nanostructure impacts photon reflectivity of the PFM and angular distribution of sputtered atoms.

[1] S. Brezinsek et al, Nucl. Fusion 59 096035 (2019)

[2] W.O. Hofer, "Sputtering by Particle Bombardment III", ed. by R. Behrisch, K. Wittmaack (Springer, Berlin, Heidelberg, 1991), Topics in Applied Physics, vol. 64, pp. 24–25

[3] M. Stepanova et al, J. Vac. Sci. Technol. A 19 2805 (2001)

## Observation of Non-Conventional Electron Dynamics and Temperature Scaling Driven by Relativistic Brunel Heating

Changhoo Lee<sup>1,2</sup>, Kyungbae Lee<sup>1,2</sup>, Gysang Lee<sup>1,2</sup>, Hyeonjin Lee<sup>1,2</sup>, Jeongwoo Nam<sup>1,2</sup>,  
and Byoung-Ick Cho<sup>1,2</sup>

<sup>1</sup> *Gwangju Institute of Science and Technology, Korea*

<sup>2</sup> *Center for Relativistic Laser Science, Institute of Basic Science, Gwangju, Korea*

We investigate hot-electron temperature scaling in relativistic laser–plasma interactions using two-dimensional particle-in-cell (PIC) simulations. Electron angular distributions extracted from the simulations are integrated to generate energy–electron-number spectra, from which hot-electron temperatures [1] are obtained. A systematic parameter scan is performed over a broad range of laser amplitudes ( $a_0$ ) and plasma density scale lengths.

The simulation results reveal electron dynamics that differ markedly from classical Brunel heating [2] and the conventional  $\mathbf{j} \times \mathbf{B}$  mechanism. Under ultra-intense laser irradiation combined with steep plasma density gradients, two clearly separated electron bunches appear. In this regime, relativistic gyromotion confines electrons near the target surface, which in turn restricts their ability to gain kinetic energy. Such electron dynamics are not captured by classical heating models. Analysis of the corresponding angular and energy distributions shows that, at high intensities and sharp gradients, the resulting hot-electron temperature falls to less than half of the value predicted by the widely used Wilks scaling. This pronounced reduction in temperature indicates the onset of a relativistic Brunel–dominated heating regime.

The distinct temperature evolution exhibited by the two electron populations provides valuable insight into how dominant energy-absorption mechanisms change with laser intensity and plasma scale length. These findings refine the current understanding of hot-electron generation in relativistic laser–matter interactions and help clarify the physical processes governing energy absorption at steep plasma interfaces.

[1] Wilks, S. C., Krueer, W. L., et al., *Phys. Rev. Lett.* 69, 1383–1386 (1992).

[2] Brunel, F., et al., *Appl. Phys. Lett.* 96, 071109 (2010)

## Frustrated Brunel Heating by Relativistic Gyromagnetic effects in Ultra-intense Laser-Matter Interactions

**Kyungbae Lee<sup>1,2</sup>, Changhoo Lee<sup>1,2</sup>, Jeongwoo Nam<sup>1,2</sup>, Hyeonjin Lee<sup>1,2</sup>, Gyusang Lee<sup>1,2</sup>, Jang Hyeob Sohn<sup>5</sup>, Elgizawi Eman<sup>1,2,4</sup>, Eunchan Lee<sup>1</sup>, Leejin Bae<sup>3</sup>, Chul Min Kim<sup>1,2,3</sup>, Byoung-Ick Cho<sup>1,2</sup>**

<sup>1</sup>*Department of Physics and Photon Science, Gwangju Institute of Science and Technology (GIST), Gwangju 61005, Republic of Korea*

<sup>2</sup>*Center for Relativistic Laser Science, Institute for Basic Science (IBS), GIST, Gwangju 61005, Republic of Korea*

<sup>3</sup>*Advanced Photonics Research Institute, Gwangju Institute of Science and Technology (GIST), Gwangju 61005, Republic of Korea*

<sup>4</sup>*Physics Department, Faculty of Woman for Arts, Science and Education, Ain Shams University, Cairo 11577, Egypt*

<sup>5</sup>*ASML Korea, Hwaseong 18488, Republic of Korea*

We report experimental and numerical investigation of a novel laser absorption mechanism, termed frustrated Brunel heating, arising from relativistic gyromagnetic effects in ultraintense laser-plasma interactions. Hot electron bunches were generated at a step-like plasma boundary irradiated by a high-contrast 150 TW, 25 fs laser pulse and propagated into a foil target. These electrons were diagnosed via coherent transition radiations (CTR) [1], revealing a unique feature: two spatially separated CTR spots, at both fundamental and second-harmonic frequencies, observed simultaneously. To elucidate the underlying mechanism, we conducted a series of 1-D and 2-D particle-in-cell (PIC) simulations. The 1-D boosted frame simulation revealed that Brunel heating [2] is frustrated when the relativistic electron gyrofrequency exceeds the laser frequency, leading to magnetic trapping and suppression of vacuum heating process. Complementary 2-D simulations reproduced the angular distribution of hot electrons, consistent with experimental observation. These results uncover a previously underappreciated role of laser-induced magnetization in modifying electron trajectory and energy absorption at steep plasma boundaries. Our findings provide deeper insight into relativistic laser energy coupling mechanisms and the generation of directional hot electron sources.

[1] J. Zheng et al., Phys. Plasmas 10, 2994 (2003).

[2] F. Brunel, Phys. Rev. Lett. 59, 52 (1987).

# First principles optical constants of water ice: geometry effects and vibrational contributions

Sina G. Lewis<sup>1</sup>, Mark C. Zammit<sup>1,2</sup>, and Amanda J. Neukirch<sup>1,2</sup>

<sup>1</sup>(*Presenting author underlined*) *Los Alamos National Laboratory, T-4: Quantum and Condensed Matter Physics*

In plasmas, atmospheres, and interstellar environments, the relevant material is often inaccessible to direct in situ measurements. Instead, spectroscopy and scattering measurements provide our main insight into the composition, density, and thermodynamic state. Radiative transfer calculations that model these observations reduce to knowing, as a function of frequency, how strongly electromagnetic radiation is attenuated and re-distributed by the medium. In practice, this requires specifying the radiative properties for each constituent in the medium, from atomic and molecular opacities to the optical constants of condensed phases.

A substantial fraction of the potential required inputs is now supported by community databases and curated tables. For gas-phase atmospheres, spectroscopic databases such as HITRAN and GEISA provide line and cross-section data that underpin predictive calculations of transmission and emission spectra [1, 2]. At higher temperatures, dominated by ionized particles, opacity tables computed from atomic structure and radiation models provide an analogous foundation; the LANL ATOMIC framework, and the associated OPLIB tables, are a representative example used widely in radiation-transport and astrophysical modeling [3, 4]. These resources are powerful, but they do not address larger particles and condensed phases such as dust, aerosols, and ice, which require knowledge of their complex optical constants.

Hexagonal ice (ice Ih) is a particularly consequential condensed phase because it appears in atmospheric clouds and snow/ice surfaces, and as mantles or solid phases in cold astrophysical environments. Widely used optical-constant datasets for ice—notably the Warren and Brandt compilation [5]—aggregate and interpolate heterogeneous measurements across preparation methods, temperatures, and morphologies. Calculating optical constants from first-principles would allow for internal consistency and yield insight into temperature, interface, and size dependence.

Here we use density functional theory calculations to compute the complex dielectric response of ice Ih and the corresponding complex refractive index. We treat electronic and ionic (vibrational) contributions separately and complement our bulk calculations with surface (slab) calculations to assess how the interface modifies absorption. We benchmark the results against the Warren and Brandt compilation [5] of heterogeneous “hexagonal” ice measurements. From this comparison, we identify regions where more complex computational techniques are required, establishing a workflow for other phases of ice and other large particles and condensed phases essential for atmospheric and astrophysical modeling.

- [1] I. E. Gordon, L. S. Rothman, et al., *J. Quant. Spectrosc. Radiat. Transf.* **277** 107949 (2022)
- [2] T. Delahaye, R. Armante, et al., *J. Mol. Spectrosc.* **380** 111510 (2021)
- [3] J. Colgan, D. P. Kilcrease, et al., *AIP Conf. Proc.* **1811**, 160002 (2017)
- [4] J. Colgan, D. P. Kilcrease, et al., *Astrophys. J.* **817** 116 (2016)
- [5] S. G. Warren, R. E. Brandt, *J. Geophys. Res.* **113**, D14220 (2008).

## Analysis of carbon ion temperatures using an upgraded multichannel Doppler spectroscopic diagnostic in NBI-heated TJ-II plasmas

**López-Miranda, B.<sup>1</sup>, Amador, D.<sup>2</sup>, Vega, J.<sup>1</sup>, Dormido-Canto, S.<sup>3</sup>, Baciero, A.<sup>1</sup>,  
McCarthy, K. J.<sup>1</sup>, and de la Riva, J.<sup>1</sup>, and TJ-II team<sup>1</sup>**

<sup>1</sup>*Laboratorio Nacional de Fusión, CIEMAT, Spain*

<sup>2</sup>*Dpto. Arquitectura de Computadores y Automática, UCM, Spain.*

<sup>3</sup>*Dpto. Informática y Automática, UNED, Spain.*

Understanding the dynamics of impurity ions in magnetically confined fusion (MCF) plasmas is essential to optimize the performance of future fusion reactors. Impurities affect plasma radiation and stability, requiring precise monitoring. Spectroscopic techniques, particularly the Doppler Effect analysis of spectral line broadening, are used to determine ion temperature ( $T_i$ ), related to ion thermal motion.

This work reports Doppler spectroscopic measurements of  $C^{+4}$  ion temperatures using an enhanced system: a Spec10 CCD with a  $400 \times 1340$ -pixel matrix coupled to a 1 m focal length spectrometer. Unlike previous TJ-II setups with 9 lenses and fibers, the new system orients 400 pixels along the wavelength dispersion and uses 1340 pixels across the plasma diameter, with binning during readout, achieving 72 points per profile (up from 9) and improving signal by ~25%. Calibration with Hg pen lamps ensures alignment precision and reliable spectroscopic parameter extraction.

Experiments with 150 neutral beam injection (NBI) heated discharges across various plasma conditions, including H isotopes, reveal  $T_i$  dependence on density and absorbed NBI power. Notably,  $T_i$  is higher during counter-injection than co-injection. These findings will be compared with neoclassical transport simulations and radial electric field measurements to analyze  $T_i$  profiles. The results contribute to understanding impurity transport in stellarators and provide a basis for studying fast NBI-driven ion interactions with impurities in MCF plasmas.

[1] Carmona, J. M. Rev. Sci. Instrum. **77** (2006) 10F107.

[2] Baciero, A. Rev. Sci. Instrum. **72** (2001) 971.

## Development of machine learning augmented CR models using the cross-sections calculated by relativistic distorted wave method

M Guru Bhavan Narayana<sup>1</sup>, Indhu Suresh<sup>1</sup>, Reetesh Kumar Gangwar<sup>1</sup>

<sup>1</sup>Department of Physics, Indian Institute of Technology Tirupati, Tirupati, Andhra Pradesh, India, 517619.

E-mail: [ph24d503@iittp.ac.in](mailto:ph24d503@iittp.ac.in)

### Abstract

Accurate modelling and diagnostics of plasma parameters are fundamental to advancing applications ranging from industrial laser processing and welding to understanding stellar atmospheres and nuclear fusion. Effective plasma diagnostics allow for the extraction of key plasma parameters, such as electron density ( $n_e$ ) and electron temperature ( $T_e$ ), which are essential for process optimization and control. In most of the applications, the plasma exhibits non-equilibrium nature and therefore to extract the physics (mostly electron density  $n_e$  and electron temperature  $T_e$ ) from the optical emission measurements, the measurements are needed to couple with adequate population-kinetic model. The plasma model requires a large set of consistent fine-structure-resolved excitation cross-sections and rate-coefficients in wide range of electron-impact energies. One such population-kinetic model is Collisional-Radiative Model (CR model) in which it considers the various collisional and radiative processes between different species in plasma to deduce the population densities of various excited levels. However, the extraction of plasma parameters using OES-cum-CR model approach is time consuming and not suitable for real time diagnostics which is crucial for many plasmas assisted processing applications. This work addresses this limitation by exploring advanced the data-driven Machine Learning approaches such as Random Forest (RF) and Deep Learning techniques such as Neural Networks and their variants <sup>[1,2]</sup> could be useful in accelerating the optical diagnostic of plasmas.

The objective of the work is to explore the suitable ML methods for the purpose of real-time diagnostics and using this approach to study Laser Produced Al plasmas. In the first phase of the work the electron impact excitation (EIE) cross sections have been calculated for fine-structure-resolved transitions of neutral aluminum (Al I) using the Relativistic Distorted Wave (RDW) approximation. Electron Impact Excitation (EIE) cross sections of Aluminum are important for spectral modeling of plasmas in or in contact with Al target. The atomic structure, including wavefunctions, energy levels, oscillator strengths ( $f_{ij}$ ), and  $A_{ij}$  values is obtained from multi-configurational Dirac-Fock calculations. The study considers transitions from the  $3s^2 3p$  manifold ( $^2P_{1/2}$ ,  $^2P_{3/2}$ ) to the higher excited manifolds  $3s^2 ns$  ( $^2S_{1/2}$ ),  $3s^2 np$  ( $^2P_{1/2}$ ,  $^2P_{3/2}$ ) with  $n = 4-6$ ,  $3s^2 nd$  ( $^2D_{3/2}$ ,  $^2D_{5/2}$ ) with  $n = 3-4$ , and  $3s^2 4f$  ( $^2F_{5/2}$ ,  $^2F_{7/2}$ ), over a projectile electron energy range of 0-100 eV. Computed energies,  $f_{ij}$  and  $A_{ij}$  values have been compared with available literature <sup>[3,4]</sup>, validating the reliability of the wavefunctions subsequently employed in the cross-section calculations. The calculated EIE cross sections are compared with previous calculations and measurements<sup>[3]</sup>. Further, Maxwellian-averaged rate coefficients have also been obtained from the calculated cross sections, providing input for collisional-radiative models to estimate plasma parameters such as electron density ( $n_e$ ) and electron temperature ( $T_e$ ). The other part of the work is ongoing, would be presented in the Joint ICTP - IAEA School.

### Keywords:

Real-time Diagnostics, Laser Produced Al Plasma, Collisional-Radiative Model, Machine Learning, Random Forest, Deep Learning, Neural Networks, Relativistic Distorted Wave approach, Electron Impact Excitation cross sections

### References:

- [1] Suresh, Indhu, P. S. N. S. R. Srikar, and R. K. Gangwar. "Integration of ML methods with CR model-based optical diagnostic for the estimation of electron temperature in Ga laser produced plasma." *Physics of Plasmas* 31.11 (2024).
- [2]. P S N S R Srikar et al, J. Phys. D: Appl. Phys. 58 415204 (2025)
- [3] Gedeon, V., Gedeon, S., Lazur, V., Nagy, E., Zatsarinny, O., & Bartschat, K. *Physical Review A*, 92(5), 052701, (2015).
- [4] <https://www.nist.gov/pml/atomic-spectra-database>

## Laser-collision induced fluorescence for measurement of collision rate coefficients in helium plasmas

**V. Meier<sup>1</sup>, A. Renggli<sup>1</sup>, F.A. Sobczuk<sup>1</sup>, C. Stollberg<sup>1</sup>, R. Jacquier<sup>1</sup>, P. Guittienne<sup>1</sup>, M. Baquero-Ruiz<sup>1</sup>, I. Furno<sup>1</sup>**

<sup>1</sup>*Ecole Polytechnique Fédérale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015, Lausanne, Switzerland*

Optical emission spectroscopy (OES) techniques are widely used to infer electron density ( $n_e$ ) and temperature ( $T_e$ ) in plasma devices. The interpretation of spectral line intensities relies on collisional-radiative models (CRMs) to retrieve plasma parameters. The accuracy of OES diagnostics is currently limited by uncertainties in the atomic data used within the CRMs. The cross-sections for electron impact (de)excitation (EIE/EID) in particular often carry large or poorly defined uncertainties that propagate all the way into results from OES diagnostics [1]. To address this issue, we propose to measure EIE/EID rates using laser-collision induced fluorescence (LCIF). The principle of LCIF is to pump an excited atomic state with a laser. A portion of this excited population transitions to a neighboring state through electron collisions, before radiatively decaying at a different wavelength. The intensity and temporal evolution of the LCIF signal depend on EIE/EID rates, themselves function of  $n_e$  and  $T_e$ . Using available EIE/EID rates from databases, LCIF has been applied to locally measure  $n_e$  and  $T_e$  [2-4] and is planned to be deployed on the ITER and T-15MD tokamak divertors [5,6]. In contrast, we plan to probe the EIE/EID rates by measuring  $n_e$  and  $T_e$  independently with Thomson scattering (TS).

This work presents proof of concept measurements of LCIF signals in helium plasmas. Experiments are performed on the Resonant Antenna Ion Device (RAID) [7], a linear helicon plasma source, where both LCIF and TS diagnostics are installed. RAID can sustain helium plasmas at densities up to  $n_e \leq 6 \cdot 10^{19} \text{ m}^{-3}$  in steady state operation, allowing for high sensitivity measurements. The plasma reaches temperatures up to  $T_e \leq 7 \text{ eV}$ , relevant to tokamak divertors [8]. To pump different atomic transitions, we use a pulsed laser, continuously tunable between 190 nm and 2000 nm. The 28 ps pulse duration allows for a clear separation of timescale between the excitation and decay processes. We use an ICCD with a 0.5 ns gate duration to measure the fluorescence.

We show experimental measurements of LCIF signals in different pumping schemes and discuss the extraction of EIE/EID rates based on previous works [9].

This research was supported by the SNSF grant 10003135.

- [1] E. Flom et al., Nuclear Materials and Energy 33, 101269 (2022)
- [2] K. Tsuchida et al., Plasma Phys. 25, 991 (1983)
- [3] E. A. Hartog et al., Phys. Rev. Lett. 62, No 13 (1989)
- [4] K. Dzierżega et al., J. Appl. Phys. 80 (6), 3196–3201 (1996)
- [5] A. V. Gorbunov et al., Plasma Phys. Rep. 38, No. 7, 574–578 (2012)
- [6] D. D. Krivoruchkoa et al., Plasma Phys. Rep. 49, No. 6, 687–699 (2023)
- [7] R. Jacquier et al., Fus. Eng. Des. 192, 113614 (2023)
- [8] B. Linehan et al., Nucl. Fus. 63, 3 (2023)
- [9] R. Denkelmann et al., J. Phys. B: At. Mol. Opt. Phys. 32, 4635 (1999)

## Electron and positron driven molecular processes for W and WH for application in plasma systems

**Pinal Mer<sup>1</sup>, and Chetan Limbachiya<sup>1</sup>**

<sup>1</sup>*The Applied Physics Department, The Maharaja Sayajirao University of Baroda, Vadodara, India*

Tungsten (W) and tungsten-based materials have been widely recognized for their excellent thermophysical properties, making them highly suitable for fusion applications, and they have been recommended as plasma-facing materials for the International Thermonuclear Experimental Reactor (ITER) and the Demonstration Power Plant (DEMO) [1]. The present work is focused on electron ( $e^-$ ) and positron ( $e^+$ ) scattering from tungsten (W) atoms and tungsten hydride (WH) molecules over a wide energy range, from the threshold up to 5 keV. These species are important plasma-facing components in the International Thermonuclear Experimental Reactor (ITER) [2]. We report elastic ( $Q_{el}$ ), inelastic ( $Q_{inel}$ ), total ionization ( $Q_{ion}$ ), summed electronic summed excitation ( $\sum Q_{exc}$ ), positronium formation ( $Q_{PS}$ ), direct ionization ( $Q_{D-ion}$ ), and total ( $Q_T$ ) cross-sections, calculated using the Spherical Complex Optical Potential (SCOP) method [3] and the Complex Scattering Potential–Ionization Contribution (CSP-ic) method [4,5]. These cross-sections serve as essential input data for plasma modelling and transport processes [6]. To the best of our knowledge, no prior experimental data are available for electron and positron interactions with the W atom and WH molecule. To provide an alternative approach and further validation of the present results, we also employed the Binary Encounter Bethe (BEB) method [7,8] to calculate  $Q_{D-ion}$  for positrons and  $Q_{ion}$  for electrons.

### References

- [1] S. E. Huber *et al.*, *J. Chem. Phys.*, **150**, 024306 (2019).
- [2] F Blanco *et al.* *Plasma Sources Sci. Technol.*, **26** 085004,(2017).
- [3] N. Thakkar, M. Swadia, M. Vinodkumar, N. Mason, and C. Limbachiya, *Plasma Sources Sci. Technol.*, **30**, 085008,(2021).
- [4] P. Mer and C. Limbachiya, *Phys. Chem. Chem. Phys.*, **26**, pp. 21504–21512, (2024).
- [5] P. Mer and C. Limbachiya, *Phys. Plasmas*, **32**, 113305 (2025).
- [6] P. Mer, N. Barad and C. Limbachiya, *J. Appl. Phys.*, **138**, 133303 (2025)
- [7] Y. K. Kim and M. E. Rudd, *Phys. Rev. A.*, **50**, 3954–3967, (1994).
- [8] K. Fedus and G. P. Karwasz, *Phys. Rev. A.*, **100** 1–14, (2019).



## ELECTRON-INDUCED VIBRATIONAL TRANSITIONS AND DISSOCIATIVE RECOMBINATION OF $H_2^+$ AND $BH^+$

**R.Mezlini<sup>1</sup>, R. Hassaine<sup>1,2</sup>, N. Pop<sup>1,3</sup>, K. Chakrabarti<sup>4</sup>, V. Laporta<sup>5</sup>, Å. Larson<sup>6</sup>, A.E. Orel<sup>7</sup>, M. Telmini<sup>8</sup>, J.Zs. Mezei<sup>1,9</sup> and I.F. Schneider<sup>1</sup>**

<sup>1</sup>*LOMC – Laboratory of Waves and Complex Media, Le Havre Normandie University, France*

<sup>2</sup>*Department of Physics & Astronomy, College of Science, Purdue University, West Lafayette, IN 47907, USA*

<sup>3</sup>*Fundamental of Physics for Engineers Department, Politehnica University Timisoara, 300223 Timisoara, Romania*

<sup>4</sup>*Department of Mathematics, Scottish Church College, Kolkata, India*

<sup>5</sup>*Istituto per la Scienza e Tecnologia dei Plasmi, CNR, Bari, Italy*

<sup>6</sup>*Department of Physics, Stockholm University, AlbaNova University Center, 106 91 Stockholm, Sweden*

<sup>7</sup>*Department of Chemical Engineering and Materials Science, University of California, Davis, California 95616, USA*

<sup>8</sup>*LSAMA, Department of Physics, Faculty of Science of Tunis, University of Tunis El Manar, 2092 Tunis, Tunisia*

<sup>9</sup>*Institute of Nuclear Research of the Hungarian Academy of Sciences, PO Box 51, Debrecen 4001, Hungary*

Hydrogen molecular ions and their isotopologues play a central role in the edge plasmas of magnetic fusion devices, as they originate from the fuel species hydrogen, deuterium, and tritium interacting with plasma-facing components. In particular,  $H_2^+$  and its isotopologues ( $D_2^+$ ,  $HD^+$ ) are of primary importance for dissociative recombination and ro-vibrational excitation processes governing plasma kinetics.

In this poster, we present recent and ongoing calculations of electron-induced dissociative recombination and vibrational transitions for  $BH^+$  and  $H_2^+$ , focusing on effective cross sections and rate coefficients. Results are presented for  $H_2^+$  at high electron energies ( $E > 3$  eV), following the MQDT methodology previously developed for  $BeH^+$  [1], as well as for  $D_2^+$  at low electron energies ( $E < 1$  eV), where rotationally resolved calculations are applicable, extending recent studies on  $H_2^+$  and  $HD^+$  [2].

The molecular structure data for  $H_2^+$  and  $BH^+$  are computed by the quantum chemistry methods as we did for  $BeH^+$  [3] and continuously improved using the Halium code [3]. These data, including potential energy curves and electronic couplings, are then used in dynamical calculations to obtain dissociative recombination and vibrational transition cross sections and rate coefficients.

More recently, boronized tungsten walls have been considered as plasma-facing materials in fusion devices, leading to the presence of boron hydride ions in the edge plasma. Consequently,  $BH^+$  has emerged as a species of growing interest. In this context, we investigate electron- $BH^+$  collisions using the Multichannel Quantum Defect Theory (MQDT).

The presented results provide valuable theoretical support for spectroscopic diagnostics and plasma modeling of edge fusion plasmas, while ongoing developments in molecular structure calculations ensure increasingly accurate descriptions of electron–molecule interactions.

- [1] E. Djuissi et al., *Phys. Rev. A* 108, 062820 (2023).
- [2] R. Hassaine et al., *Astron. Astrophys.* 703, A220 (2025).
- [3] S. Niyonzima et al., *Phys. Rev. A* 87, 022713 (2013).
- [4] M. Telmini and C. Jungen, *Phys. Rev. A* 68, 062704 (2003).

## Construction of atomic collisional radiative models for use in solar atmospheric modelling

**Nicolas A. Mori<sup>1</sup> and Giulio Del Zanna<sup>1</sup>**

*<sup>1</sup>DAMTP, Centre of Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK*

For the solar atmosphere the most important atomic transitions originate from hydrogen and helium atoms which compose over 98% of the Sun by mass. Recently, several new collisional radiative models have been produced for helium and its ion [1,2]. These models aimed to address shortcomings in the He recombination spectra and collisional excitation rates present in previous models through the use of more accurate atomic data. Between the present and previous models differences were found to be far above the target accuracy of 1% for predicted emissivity. Following from these models, similar models for atomic hydrogen are to be constructed, but for inclusion into Radiative Transfer/MHD codes reduced models are being built, also considering the evolution of the populations with time dependence. Thereafter, these models will also be modified for inclusion into the Solar Atomic Modelling Suite (SAMS) project [3]. The models will have wide-ranging applications in astrophysics and laboratory plasmas.

[1] G. Del Zanna, P. J. Storey, N. R. Badnell, and V. Andretta. *ApJ*, 239, 869 (2020).

[2] G. Del Zanna and P. J. Storey, *Mon. Not. R. Astron. Soc.*

[3] <https://sams-project-uk.github.io/>

## Electron collision studies on boron hydride species in fusion edge plasmas

**N. Mukherjee<sup>1</sup>, A. Bhattacharyya<sup>2</sup>, and K. Chakrabarti<sup>1</sup>**

<sup>1</sup>*Department of Mathematics, Scottish Church College, Kolkata – 700006, India*

<sup>2</sup>*Department of Physics, University of Calcutta, Kolkata – 700009, India*

Wall conditioning is an essential requirement for reliable tokamak operation, as plasma performance is highly sensitive to impurities release from the plasma-facing components (PFCs) [1-3]. Boronization is a highly effective wall conditioning technique that deposits boron layer on the PFCs, where it acts as an efficient getter for oxygen and other residual impurities, reducing radiative losses and improve plasma performance in fusion systems [1,2,4]. As ITER transitions to a full tungsten machine with boronized walls [5], plasma-material interactions involving boronized surfaces lead to release of boron atoms and ions, thereby introducing boron-based impurities into the plasma and influencing plasma edge chemistry and impurity transport. Considering the plasma's predominant hydrogenic composition, boron hydride species and their isotopologues are likely to be the leading source of impurities within the plasma. Accurate molecular and collision data for these species are therefore essential for modeling impurity transport, plasma flow and deposition processes. Using the R-matrix method [1,6], we have performed low energy electron collision calculations on both neutral BH and its molecular cation BH<sup>+</sup> at their equilibrium geometries [1,6]. For BH, we have constructed a good quality target wavefunction at the CI level to compute the state resolved elastic, electronic excitation, differential, and momentum transfer cross sections, together with an estimate of electron-impact dissociation cross section [6]. Additionally, a <sup>2</sup>Π resonant state of BH<sup>-</sup> was also reported, providing insights into pathways for electronic processes which occur via transient anionic states [6]. Extending this work to BH<sup>+</sup>, we also conducted a detailed electron-ion collision study on this molecular ion calculating electronic excitation cross sections from the BH<sup>+</sup> ground state to the low-lying excited states [1]. Alongside this, we have also investigated isotope effects at low energies, and calculated the cross sections for other processes providing pathways for rotational excitation and ionization [1]. We have also performed structure calculations on these molecules and obtained their Rydberg bound and resonant states. The data supporting our findings are openly available in the CollisionDB repository of IAEA [7].

[1] N. Mukherjee, A. Bhattacharyya, K. Chakrabarti, *Phys. Plasmas* **32**, 102502 (2025).

[2] J. Winter, *Plasma Phys. Control. Fusion* **38**, 1503 (1996).

[3] L. Peng, J. Sun, Z. Sun, F. Gao, X. Bonnin, J.Y. Liu, *Nucl. Mater. Energy* **26**, 100937 (2021).

[4] A. Bortolon, R. Maingi, A. Nagy, J. Ren, J. Duran, A. Maan, D. Donovan, J. Boedo, D. Rudakov, A. Hyatt, T. Wilks, M. Shafer, C. Samuell, M. Fenstermacher, E. Gilson, R. Lunsford, D. Mansfield, T. Abrams, R. Nazikian, *Nucl. Fusion* **60**, 126010 (2020).

[5] ITER, see <http://www.iter.org/>.

[6] K. Chakrabarti, N. Mukherjee, A. Bhattacharyya, J. Tennyson, *Plasma Phys. Control. Fusion* **67**, 025009 (2025).

[7] CollisionDB, see <https://amd.is.iaea.org/db/collisiondb/> for database of plasma collisional processes.

## Steady-State-Informed Latent Dynamics Surrogate Model for TD-NLTE Atomic Kinetics

**Jeongwoo Nam<sup>1,2</sup>, Minsang Cho<sup>3</sup>, William Anderson<sup>3</sup>, Youngsoo Choi<sup>3</sup>, Haewon Jeong<sup>4</sup> and Byoung-Ick Cho<sup>1,2</sup>**

<sup>1</sup>*Department of Physics and Photon Science, Gwangju Institute of Science and Technology (GIST), Gwangju 61005, Republic of Korea*

<sup>2</sup>*Center for Relativistic Laser Science, Institute for Basic Science (IBS), GIST, Gwangju 61005, Republic of Korea*

<sup>3</sup>*Lawrence Livermore National Laboratory, 7000 East Avenue, Livermore, California 94550, United States of America*

<sup>4</sup>*Department of Electrical and Computer Engineering, University of California, Santa Barbara (UCSB), California 93106, United States of America*

Time-dependent non-local thermodynamic equilibrium (TD-NLTE) atomic calculation plays a key role in plasma simulations across a wide range of applications, including inertial confinement fusion. Inline coupling of TD-NLTE calculation with radiation–hydrodynamic simulations enables more accurate modeling by reducing the gap between simulations and experiments, but its practical use remains limited by the high computational cost. We develop a physics-informed surrogate TD-NLTE model based on the Latent Space Dynamics Identification (LaSDI) framework [1], with the key novelty being the explicit incorporation of NLTE steady-state constraints. High-dimensional atomic population data generated by SCFLY [2] are compressed into a low-dimensional latent space using a nonlinear autoencoder with a mixed reconstruction loss that captures both microscopic population accuracy and macroscopic observables. The latent dynamics are identified using Sparse Identification of Nonlinear Dynamics with control (SINDyC) [3], driven by time-dependent temperature and density histories. To ensure physically meaningful long-term behavior, the analytically derived latent steady state is explicitly constrained to match reference NLTE steady-state solutions, in addition to enforcing Hurwitz stability. These preliminary results demonstrate the feasibility of a steady-state–constrained latent dynamics approach for efficient and physically consistent TD-NLTE modeling, paving the way toward practical inline coupling with radiation–hydrodynamic simulations.

[1] Fries, William D., Xiaolong He, and Youngsoo Choi, *Comput. Methods Appl. Mech. Eng.* **399**, 115436 (2022).

[2] H.-K. Chung et al., *AIP Conf. Proc.* 1811, 020002 (2017).

[3] S. L. Brunton, J. L. Proctor, and J. N. Kutz, *IFAC-PapersOnLine* 49, 710 (2016).

**Effective cross sections and rates coefficients derived from collision-induced rotational excitation of  $\text{HCl}^+(X^2\Pi)$  with  $\text{He}(^1S)$ : Isotopic effects**

**Joseph Ngueleo Baldagui<sup>†</sup>, Théophile Tchakoua<sup>\*</sup>, Jean Jules Fifen<sup>‡</sup>, Mama Nsangou<sup>†,‡</sup>**

<sup>†</sup>*Department of Physics, Faculty of Science, University of Maroua, P.O. Box. 814, Maroua, Cameroon.*

<sup>\*</sup>*Leiden Institute of Chemistry, Gorlaeus Laboratories, Leiden University, P.O. Box 9502, 2300 RA Leiden, The Netherlands*

<sup>‡</sup>*Department of Physics, Faculty of Science, University of Ngaoundere, P.O. Box. 454, Ngaoundere, Cameroon.*

<sup>†,‡</sup>*Department of Physics, Faculty of Science, University of Maroua, P.O. Box. 814, Maroua, Cameroon.*

In this work, we studied the helium-induced collisional excitation of the radical ion  $\text{HCl}^+$ . Our work focuses on calculating three-dimensional potential energy surfaces (PES) to study the interaction due to the collision between  $\text{HCl}^+$  and He, and on analyzing the influence of the isotopic effect on cross sections and collision rates. For *Ab initio* calculations of PES  $^2A'$  and  $^2A''$  of  $\text{HCl}^+(X^2\Pi)$ -He complex, we used the RCCSD(T)-F12 method with cc-pVQZ-F12 basis sets. These surfaces have been fitted using the Reproducing Kernel Hilbert Space (RKHS) method and were submitted to the close-coupling approach in order to work out the inelastic integral cross sections. Collision cross sections taking into account the fine structures of  $\text{HCl}^+$  have been performed for kinetic energies up to  $3500\text{ cm}^{-1}$  and the thermal excitation rates for kinetic temperatures from 0K to 400 K. It appears that the difference in the cross section and collisional rate coefficients for the  $\text{H}^{35}\text{Cl}^+$  and  $\text{H}^{37}\text{Cl}^+$  colliding with He was found to be negligible. In contrast, a significant difference in effective cross-sections and collision rates between  $\text{HCl}^+$ -He and  $\text{DCl}^+$ -He was observed to the extent that it is impossible to make estimation of collision rates of deuterated species from those of the hydrogenated species.

## Absolute calibration of various crystal types and geometries for National Ignition Facility x-ray spectrometers

**P. Parker<sup>1</sup>, M. Dozieres<sup>1</sup>, C. M. Krauland<sup>1</sup>, S. Stoupin<sup>2</sup>, N. Thompson<sup>2</sup>, J. Buscho<sup>2</sup>, J. Corbin<sup>2</sup>, J. Huckins<sup>2</sup>, T. Goyal<sup>2</sup>, M. Beach<sup>2</sup>, M. B. Schneider<sup>2</sup>, J. Seely<sup>3</sup>**

<sup>1</sup>*General Atomics, San Diego, 92186, California, USA*

<sup>2</sup>*Lawrence Livermore National Laboratory, Livermore, 94550, California, USA*

<sup>3</sup>*National Institute of Standards and Technology, Gaithersburg, MD 20899, USA*

Absolute calibration of crystals within x-ray spectrometers fielded at the National Ignition Facility (NIF) is critical to understand the plasma source properties that they aim to diagnose. The spectrometer calibration station (SCS) located at the Lawrence Livermore National Laboratory was established in recent years to perform integrated spectrometer throughput measurements in order to fully characterize a given crystal's response. Full NIF spectrometer or duplicate surrogate hardware is placed in a vacuum chamber adjacent to the x-ray source chamber, such that the crystal-to-source standoff can be adjusted to its actual position from a target in the NIF. We use a steerable electron beam to bombard an anode foil as our x-ray source. The anode foil is housed in a rotatable turret, allowing for an easy selection of various elements without breaking vacuum. This combined with a voltage of up to 30 kV, for the electron beam power supply, allows for x-ray line emission of up to 30 keV photon energies. The crystal response is determined by taking a ratio of the crystal diffracted signal to the source emission spectrum for a given set of x-ray lines within the spectrometer's bandwidth. We use an absolutely calibrated Amptek silicon drift detector to infer the source emission signal. The SCS further characterizes crystal attributes by illuminating any spectral warping or defects that can only be seen with x-rays. Spectral dispersion is also mapped so that theoretical design can be vetted. Here we present the calibration data from various NIF x-ray spectrometers of different geometries and crystal types. We show that often each crystal is unique and requires individual calibration, regardless of identical design and fabrication specifications.

Corresponding Author Email: [parker76@llnl.gov](mailto:parker76@llnl.gov)

This work performed under the auspices of the U.S. Department of Energy by General Atomics under Contract 89233119CNA000063 and by Lawrence Livermore National Laboratory under Contract DE-AC52-07NA27344.  
LLNL-ABS-866201

## Near-Surface Hydrogen Inventory Response to Picosecond Laser Pulses in Tungsten

**M. Popova<sup>1,2</sup>, D. Matveev<sup>1</sup>, S. Brezinsek<sup>1,2</sup>, C. Kawan<sup>1,2</sup>, E. Wüst<sup>1,2</sup>, G. Sergienko<sup>1</sup>, M. Zlobinski<sup>1</sup>**

<sup>1</sup> *Forschungszentrum Jülich GmbH, Institute of Fusion Energy and Nuclear Waste Management – Plasma Physics, 52425 Jülich, Germany*

<sup>2</sup> *Mathematisch-Naturwissenschaftliche Fakultät, Heinrich-Heine-Universität Düsseldorf, Düsseldorf, Germany*

Tungsten is the leading candidate for plasma-facing components in fusion devices. In future fusion reactors, neutron irradiation creates material defects that trap hydrogen isotopes, including the fusion fuel tritium and deuterium. Monitoring the hydrogen inventory is mandatory for nuclear safety and important for the efficient use of fuel. Laser-based diagnostic methods such as Laser-Induced Breakdown Spectroscopy (LIBS) and Laser-Induced Ablation with detection of volatile species by Quadrupole Mass Spectrometry (LIA-QMS) provide quantitative, depth-resolved assessments of the local fuel inventory and release. They are offering a controlled measurement of near-surface retention on short time scales for in-situ and ex-situ applications

To describe the effect of transient material heating on fuel release beyond the ablated layer, a one-dimensional modelling framework is developed in FEniCS/FESTIM for picosecond laser pulse trains. The simulations address depths below 1 mm. Transient heat conduction with temperature-dependent properties is coupled to a minimal near-surface trapping-detrapping scheme. Initial near-surface fuel fractions are in the percent to sub-percent range. Coarse ablation is represented by removing a thin surface layer tens of nanometers thick at the start of each pulse, which updates both the thermal and hydrogen-isotope boundary conditions. Per-pulse release is obtained by integrating the simulated desorption flux. Evolution over fuel release is analyzed for tens to hundreds of pulses.

Here in this modelling approach, the initial fuel distribution is taken from NRA depth profiles measured on proton-irradiated and self-damaged, deuterium-decorated tungsten samples which act as reference.

Results show that heating-induced release dominates over the contribution attributable to the imposed ablation layer release. Early pulses are characterized by rapid mobilization of the near-surface fuel reservoir, leading to pronounced, prompt desorption transients. Consequently, when ablation is applied at the start of subsequent pulses, it largely removes a layer already partially depleted by prior heating. With a simple defect-annealing term, model predictions are compared to LIA-QMS-derived depth reconstructions, yielding reasonable agreement.

The modelling framework supports the interpretation of LIA-QMS data for self- or proton damaged tungsten as proxy for neutron-damaged tungsten and clarifies the respective roles of fast heating and material ablation in pulse-train experiments with ps-laser systems.

## Optical Emission Spectroscopy in Helium plasmas for validation of collisional-radiative model on the Resonant Antenna Ion Device

A. Renggli, V. Meier, F.A. Sobczuk, C. Stollberg, M. Baquero Ruiz, R. Jaquier, P. Guittienne, I. Furno

*Ecole Polytechnique Fédérale de Lausanne (EPFL), Swiss Plasma Center (SPC), CH-1015 Lausanne, Switzerland*

Helium (He) is an important element in plasma physics due to its abundance in space and laboratory plasmas, as well as its relative simplicity in atomic structure. Optical Emission Spectroscopy (OES), combined with collisional radiative models (CRMs) can be used to determine the electron density  $n_e$  and temperature  $T_e$ . However the relation between plasma parameters and spectral line intensity is complex and present CRMs have been found to break down under certain conditions, particularly at low temperatures in tokamak divertors, which emphasizes the further need for development and validation of He CRMs [1].

The intensity of a given spectral line is proportional to both the density  $n_i$  of the emitting state and the Einstein coefficient  $A_{ij}$ , i.e.  $\varepsilon_{ij} \propto n_i A_{ij}$ . The distribution of the excited states is described by the CRM as a function of the plasma parameters, accounting for both radiative and collisional processes. Development of an accurate CRM is challenging and requires an exact description of the atomic structure, inclusion of all relevant processes, availability of precise reaction rates, as well as a correct treatment of transport processes, such as opacity.

We are currently investigating He plasmas on the Resonant Antenna Ion Device (RAID) [2] for validation of the CRM CoRa-He [3] developed at SPC, EPFL. RAID produces steady-state plasmas with  $n_e \approx 0.3 - 5 \cdot 10^{19} \text{ m}^{-3}$ ,  $T_e \lesssim 5 \text{ eV}$ , relevant for tokamak divertor physics. Plasma emission is collected through an optical fibers bundle and guided to an Isoplane SCT 320 spectrometer ( $f = 320 \text{ mm}$ ,  $\lambda$ : 200-1300 nm, 150 l/mm grating) to obtain radial emission profiles, which are then interpreted using CoRa-He, accounting for the plasma's environment and opacity effects. Complementary diagnostics, particularly RAID's high resolution Thomson scattering (TS) system, provides precise and independent characterization of the plasma parameters, used as input to the CRM.

We present OES measurements of He plasmas in RAID alongside CRM simulations and discuss the influence of opacity and transports on these measurements. Future work on the CRM will include several updates, such as the addition of higher level ( $n > 4$  for He I), level-mixing between singlet and triplet states [4] as well as the development of a more complex opacity model. Furthermore, the laser induced fluorescence (LCIF) diagnostic implemented on RAID offers another powerful tool to validate the CRM when the plasma is perturbed out of its equilibrium state [5].

This work was funded by the Swiss National Science Foundation, Grant No 10003135

- [1] B. L. Linehan *et al.*, Nucl. Fusion **63** 036021, (2023)
- [2] I. Furno *et al.*, EPJ Web of Conferences **157**, 03014, (2017)
- [3] L. R. Kadi, PhD thesis, *Collisional-radiative modeling and ultrafast laser-induced fluorescence in a helicon plasma device*, Ecole Polytechnique Fédérale de Lausanne, (2025)
- [4] M. Goto, J. Quant. Spectrosc. Radiat. Transfer **76**, 331-344, (2003)
- [5] R. Denkelmann *et al.*, J. Phys. B: At. Mol. Opt. Phys. **32** 4635, (1999)

## EMC3-EIRENE predictions of radiative detachment scenarios in W7-X equipped with a tungsten based divertor

Daniil Ryndyk<sup>1</sup>, Derek Harting<sup>1</sup>, Shuai Xu<sup>1</sup>, Heinke Frerichs<sup>2</sup>, Yuhe Feng<sup>3</sup>, Florian Effenberg<sup>4</sup>, Sebastijan Brezinsek<sup>1,5</sup> and the W7-X team

<sup>1</sup>(Presenting author underlined) Forschungszentrum Jülich GmbH, Institute of Fusion Energy and Nuclear Waste Management – Plasma Physics, 52425 Jülich, Germany

<sup>2</sup>University of Wisconsin-Madison, Madison, WI, USA

<sup>3</sup>MPG IPP Greifswald, 17491 Greifswald, Germany

<sup>4</sup>PPPL, Princeton, NJ 08543, USA

<sup>5</sup>Faculty of Mathematics and Natural Sciences, Heinrich Heine University Düsseldorf, 40225 Düsseldorf, Germany

Wendelstein 7-X (W7-X) operates with carbon-fiber composite (CFC) plasma-facing components (PFC) forming the island divertor used for particle and power exhaust. Erosion of PFCs leads to an influx of carbon (C) into the edge plasma where C radiation significantly cools down the divertor in the applied hydrogen plasmas. Raising the radiation in the divertor by hydrogen fueling and/or impurity seeding of low-Z species like N<sub>2</sub> or Ne allows access to the “radiative detachment” regime with reduced target heat and particle loads [1,2] at high electron densities and moderate electron temperatures in the divertor of about 10 eV. The seeded impurities replace partially the intrinsic impurity C as the proton impact energy falls below the physical sputtering threshold. However, carbon-based PFCs are not tolerable for a future fusion reactor due to unacceptable fuel retention as shown e.g. in JET.

A promising alternative PFC material is tungsten (W), which shows in general low sputtering, low fuel retention, and good thermomechanical properties. As a first systematic approach to study the impact of potential W PFCs in W7-X, we use EMC3-EIRENE as plasma boundary simulation code for 3D magnetic configurations. When C as intrinsic radiator is absent, impurity seeding is unavoidable for plasma operation at elevated power and particle fluxes [3]. We investigate detachment scenarios with the standard magnetic configuration of W7-X equipped with a W divertor in the current divertor design and analyze the dependence of detachment relevant parameters (e.g. neutral particle density and target heat loads) on Ne impurity concentration and operation conditions (e.g. heating power and plasma density). First results indicate a deeper penetration of recycled H neutrals into the confined region as a consequence of the increased reflection probability of H ions at the strike lines. It is anticipated that low impact energies and cold divertor operation with electron temperatures below 5 eV are required to restrict W erosion caused by the seed impurities to avoid significant W accumulation in the core. At these temperatures, volume recombination processes become increasingly relevant and hence are also investigated since up to now, the research on volume recombination in the island divertor has been limited to the C divertor [4]. Both the hydrogen recycling and seed impurity concentration need to be adapted to get access to this reactor-relevant regime and eventually, a change of the island divertor geometry with higher baffeling might be unavoidable.

[1] Y. Feng *et al*, Nucl. Fusion **61**, 086012 (2021).

[2] F. Effenberg *et al*, Nucl. Fusion **59**, 106020 (2019).

[3] S. Brezinsek *et al*, Nucl. Fusion **59**, 096035 (2019).

[4] Y. Feng *et al*, Nucl. Fusion **65**, 066008 (2025).

## A general approach to calculating ion-atom cross sections, applied to proton collisions with oxygen

K. Schrick<sup>1</sup>, C. T. Plowman<sup>1</sup>, and N. A. Mori<sup>2</sup>, I. Bray<sup>1</sup> and D. V. Fursa<sup>1</sup>

<sup>1</sup>*Department of Physics, Curtin University, Perth, WA 6845, Australia*

<sup>2</sup>*Department of Applied Mathematics and Theoretical Physics, Centre for Mathematical Sciences, University of Cambridge, Wilberforce Road, Cambridge CB3 0WA, UK*

Collisions between ions and atoms are found in a wide range of environments, including planetary atmospheres [1], fusion plasmas [2], and hadron therapy for cancer treatment [3]. To adequately model and analyse these environments, accurate cross sections are required for a range of atomic processes, including ionisation and excitations. Calculating these cross sections is particularly challenging for collisions with multi-electron atoms, as it is not feasible to account for the electrons individually.

The convergent close-coupling (CCC) method for collisions between positrons and light atoms combines wave functions generated using the multi-configurational Hartree Fock method [4] with Laguerre-based pseudostates [5] to model atomic processes. This method has been shown to produce accurate cross sections for a variety of targets [6, 7, 8]. However, for an ionic projectile, a fully quantum-mechanical method is not practical or necessary for collision energies greater than 1 keV, where the cross sections are most significant. Instead, the semi-classical method can be used to simplify the scattering problem. We have adapted the general and scalable semi-classical scattering equation solver from Ref. [9] to the general atomic structure code [10] used for light projectiles in Ref. [6, 7, 8].

As the first step, we apply our new suite of codes to the proton-oxygen system, for impact energies in the keV to MeV range. These particular cross sections have applications in Auroral modelling due to the prevalence of atomic oxygen in the upper atmosphere [1], and as a molecular constituent through independent-atom approaches [11]. Previous calculations approximated the field generated by the many electrons in the oxygen atom using model potentials [12]. However, the present approach provides a substantially more accurate description of the electronic structure and therefore a more detailed depiction of the underlying physics.

- [1] E. J. Schaefer, *Journal of Geophysical Research* (1896-1977) **68**, 1175 (1963).
- [2] O. Marchuk, *Phys. Scr.* **89**, 114010 (2014).
- [3] E. Scifoni, W. Tinganelli, W. K. Weyrather, M. Durante, A. Maier, and M. Krämer, *Physics in medicine and biology* **85**, 10.1088/0031-9155/58/11/3871 (2013).
- [4] C. Froese Fischer, *Computer Physics Communications* **64**, 369 (1991).
- [5] I. Bray and A. T. Stelbovics, *Phys. Rev. A* **46**, 6995 (1992).
- [6] N. A. Mori, L. H. Scarlett, I. Bray, and D. V. Fursa, *Phys. Rev. A* **107**, 032817 (2023).
- [7] N. Mori, L. Scarlett, I. Bray, and D. Fursa, *The European Physical Journal D* **77**, 182 (2023).
- [8] N. Mori, L. Scarlett, I. Bray, and D. Fursa, *The European Physical Journal D* **78**, 19 (2024).
- [9] C. T. Plowman, L. H. Scarlett, M. C. Zammit, I. Bray, and D. V. Fursa, *Phys. Rev. A* **112**, 042811 (2025).
- [10] O. Zatsarinny, *Computer Physics Communications* **174**, 273 (2006).
- [11] N. A. Mori, I. Bray, and D. V. Fursa, *The European Physical Journal D* **78**, 28 (2024).
- [12] T. Kirchner, H. J. Lüdde, M. Horbatsch, and R. M. Dreizler, *Phys. Rev. A* **61**, 052710 (2000).

## Interparticle Coulombic electron capture beyond the virtual photon approximation

Jan Šenk<sup>1,2</sup>, Vincent Graves<sup>3</sup>, Jimena D. Gorfinkiel<sup>3</sup>, Přemysl Koloreňč<sup>2</sup>, and Nicolas Sisourat<sup>1</sup>

<sup>1</sup>*Sorbonne Université, CNRS, Laboratoire de Chimie Physique – Matière et Rayonnement, UMR 7614, F-75005 Paris, France*

<sup>2</sup>*Institute of Theoretical Physics, Faculty of Mathematics and Physics, Charles University, V Holešovičkách 2, 180 00 Prague, Czech Republic*

<sup>3</sup>*School of Physical Sciences, The Open University, Walton Hall, Milton Keynes MK7 6AA, United Kingdom*

Interparticle Coulombic electron capture (ICEC) is an environment-enabled electron capture process where the excess energy is released by ionization or excitation of a neighboring particle. Both the capturing particle and the neighbor can be atoms, molecules, or ions. It is one of the plethora of electron capture (or electron attachment) processes that are relevant to astrophysics [1] and plasma physics [2]. ICEC was theoretically predicted in 2009 [3, 4] and is expected to be significantly more efficient than the competing photorecombination. To date, it has been studied using analytical approximations [5], theoretical models [6], and *ab initio* calculations [7]. It has yet to be experimentally observed.

The asymptotic approximation derived in [4] has been shown to severely underestimate the ICEC cross section when the distance between the two particles is comparable to their size. It treats the process as two independent events: virtual photorecombination at the capturing particle and virtual photoionization of the neighbor. They are linked by the transfer of the excess energy. It therefore neglects the correlation between the electrons of the two particles and, most importantly, the possibility of electron transfer from the neighbor to the capturing particle. We have improved on the asymptotic approximation by proposing an analytical model of the electron transfer mechanism of ICEC. We have benchmarked it with *ab initio* R-matrix scattering calculations for several systems [8].

In comparison to the original asymptotic approximation, our model improved the agreement with the *ab initio* results at relevant inter-particle distances while keeping its simplicity. The model ICEC cross sections can be evaluated with a simple analytical formula using tabulated properties of the participating particles.

- [1] F. Carelli, T. Grassi, and F. A. Gianturco, *A&A* **549**, A103 (2013).
- [2] H. P. Summers, W. J. Dickinson, in *Recombination of Atomic Ions*, edited by W. G. Graham, W. Fritsch, Y. Hahn, and J. A. Tanis, NATO ASI Series, Vol. 296 (1992).
- [3] K. Gokhberg, L. S. Cederbaum, *J. Phys. B: At. Mol. Opt. Phys.* **42**, 231001 (2009).
- [4] K. Gokhberg, L. S. Cederbaum, *Phys. Rev. A* **82**, 052707 (2010).
- [5] A. Molle, O. Zatsarinny, T. Jagau, A. Dubois, N. Sisourat, *J. Chem. Phys.* **158**, 134306 (2023).
- [6] F. M. Pont, A. Bande, L. S. Cederbaum, *J. Phys.: Condens. Matter* **28**, 075301 (2016).
- [7] N. Sisourat, T. Miteva, J. D. Gorfinkiel, K. Gokhberg, L. S. Cederbaum, *Phys. Rev. A* **98**, 020701 (2018).
- [8] J. Šenk, V. Graves, J. D. Gorfinkiel, P. Koloreňč, N. Sisourat, *J. Chem. Phys.* **161**, 174113 (2024).

## Full-dimensional quantum dynamic study for the H atom scattering from graphene surface

Lei SHI<sup>1</sup>

<sup>1</sup> *Laboratoire Ondes et Milieux Complexes-UMR6294, CNRS, Université Le Havre Normandie*

The adsorption of hydrogen atoms on graphene has received significant attention due to its importance in astrophysics, hydrogen storage, and semiconductor technologies. A full-dimensional potential energy surface (PES) for a periodic graphene surface containing 24 carbon atoms and one hydrogen atom was recently developed [1,2]. Based on this PES, classical molecular dynamics (MD) simulations were used to study hydrogen atom scattering from graphene. However, for certain collision conditions, discrepancies were observed between experimental results and MD predictions.

To clarify these differences, full-dimensional quantum dynamics (QD) simulations were performed. The original neural-network PES was refitted into a sum-of-products representation using the Monte-Carlo Canonical Polyadic Decomposition (MCCPD) method [3]. The nuclear wave function was then propagated using the multilayer multiconfiguration time-dependent Hartree (ML-MCTDH) method [4,5,6], with a multilayer tree optimized through hierarchical clustering [7]. Sticking probabilities and scattering distributions were obtained using quantum flux-based approaches.

The comparison between QD and MD simulations showed very good agreement [8,9], demonstrating that classical dynamics captures much of the essential physics. Importantly, the analysis revealed that remaining discrepancies with experimental observations arise primarily from PES limitations and genuine quantum effects, such as zero-point energy and resonance phenomena. Furthermore, phonon-resolved analysis highlighted the key role of longitudinal graphene phonons in the collision and adsorption mechanisms.

[1] Jiang, H., Kammler, M., Ding, F., Dorenkamp, Y., Manby, F. R., Wodtke, A. M., Miller, T. F., Kandratsenka, A., Bünemann, O. *Science* 364(6438), 379-382 2019.

[2] Wille, S., Jiang, H., Bünemann, O., Wodtke, A. M., Behler, J., Kandratsenka, A. *Physical Chemistry Chemical Physics* 22(45), 26113–26120 2020.

[3] Schröder, M. *Journal of Chemical Physics* 152(2), 2020.

[4] Beck, M. H., Jackle, A., Worth, G. A., Meyer, H. D. *Physics Report* 324(1), 1-105 2000.

[5] Manthe, U. *Journal of Chemical Physics* 128(16), 2008.

[7] Mendive-Tapia, D., Meyer, H. D., Vendrell, O. *Journal of Chemical Theory and Computation* 1, 2023.

[6] Wang, H., Thoss, M. *Journal of Chemical Physics* 119, 2003.

[8] Shi, L., Schröder, M., Meyer, H. D., Pelaez, D., Wodtke, A. M., Golibrzuch, K., Schönemann, A. M., Kandratsenka, A., Gatti, F., *Journal of Chemical Physics* 159, 2023.

[9] Shi, L., Schröder, M., Meyer, H. D., Pelaez, D., Wodtke, A. M., Golibrzuch, K., Schönemann, A. M., Kandratsenka, A., Gatti, F., *The Journal of Physical Chemistry A* 129 (7), 1896-1907, 2025.

# Relativistic Calculations Of Transition Parameters Of Doubly Ionized Argon

**Akansha Singhal , Lalita Sharma**

*Department of Physics, IIT Roorkee, Roorkee 247667 , India*

Atomic structure parameters for inert gas ions are of significant importance due to their wide uses in fusion plasma modeling and diagnostics. In fusion devices, inert gases can exist in various charge states therefore, reliable atomic data are essential to support ongoing projects focused on the design and development of fusion reactors.

Limited data are available for the transition parameters of doubly ionized argon ( $\text{Ar}^{2+}$ ). To address this gap, we have carried out detailed calculations of energy levels and transition parameters, including wavelengths, weighted oscillator strengths, transition probabilities, and line strengths, for electric dipole (E1) and magnetic dipole (M1) transitions. The computations were performed using the fully relativistic multiconfiguration Dirac–Hartree–Fock method implemented in GRASP2018 [2], which incorporates both Breit interaction and quantum electrodynamics effects. These calculations were conducted for the lowest 230 fine-structure levels of sulphur-like  $\text{Ar}^{2+}$ , arising from the configurations  $3s^23p^4, 3p^6, 3s^23p^34\ell$  ( $\ell = s, p, d, f$ ),  $3s3p^5, 3s^23p^33d$ , and  $3p^53d$ .

Our calculated energy levels show excellent agreement with the 125 experimentally measured [1,3] levels reported in the NIST database. Furthermore, good consistency is observed between the published line strength data and our calculated results. To further validate the accuracy of our calculations, we compared the line strengths obtained in both the length and velocity gauges, which also exhibit strong agreement.

[1] Moore, C. E. (1949). Atomic energy levels as derived from the analyses of optical spectra (Vol. 1). US Department of Commerce, National Bureau of Standards.

[2] Fischer, C. F., Gaigalas, G., Jönsson, P., & Bieroń, J. (2019). GRASP2018—A Fortran 95 version of the general relativistic atomic structure package. *Computer Physics Communications*, 237, 184-187

[3] Luna, F. R. T., Bredice, F., Cavalcanti, G. H., & Trigueiros, A. G. (2000). Atomic transitions for the doubly ionized argon spectrum, Ar III. *Brazilian Journal of Physics*, 30, 386-391.

## Photoionization, Rayleigh, and Raman scattering cross sections for the hydrogen molecule

Adam J. C. Singor<sup>1</sup>, Mark C. Zammit<sup>1</sup>, Liam H. Scarlett<sup>2</sup>, Igor Bray<sup>2</sup>, and Dmitry V. Fursa<sup>2</sup>

<sup>1</sup>*Theoretical Division, Los Alamos National Laboratory, Los Alamos, New Mexico 87545, USA*

<sup>2</sup>*Department of Physics and Astronomy, Curtin University, Perth, Western Australia 6102, Australia*

Photon scattering cross sections have proved essential in many applications, such as calculating opacities, modeling plasma and radiative transport, analyzing planetary atmospheres, and Raman spectroscopy. In particular, Raman spectroscopy of H<sub>2</sub> plays an important role in analyzing hydrogen storage techniques, monitoring ortho-to-para conversion, and monitoring nuclear waste. Atomic and molecular hydrogen are abundant in the interstellar medium. Hence, photon scattering cross sections for H<sub>2</sub> and its ion are of particular interest in astrophysics.

Photon–molecule scattering processes have been well understood to second order in perturbation theory since the development of the Kramers–Heisenberg–Waller (KHW) matrix element [1, 2] in the mid-1920s. The convergent close-coupling method, which has previously been used to study electron scattering from atoms and molecules as well as single and double photoionization [3], has been applied to photoionization and radiative association of the hydrogen molecule. We have calculated rovibrationally-resolved photoionization cross sections for the ground electronic state of H<sub>2</sub>. Rayleigh and Raman scattering cross sections have been calculated for transitions between all rovibrational levels of the  $X^1\Sigma_g^+$  ground electronic state of H<sub>2</sub>, resulting in a total of 9582 Rayleigh and Raman cross sections [4]. This work represents the most comprehensive study of photon scattering on molecular hydrogen. Thermally-averaged photoionization, Rayleigh, and Raman scattering cross sections have been produced for a system in local thermodynamic equilibrium.

[1] H. A. Kramers and W. Heisenberg, *Z. Phys.* **31**, 681 (1925).

[2] I. Waller, *Z. Phys.* **51**, 213 (1928).

[3] I. Bray, D. V. Fursa, A. S. Kheifets, and A. T. Stelbovics, *J. Phys. B: At. Mol. Opt. Phys.* **35**, R117 (2002).

[4] A. J. C. Singor, L. H. Scarlett, M. C. Zammit, I. Bray, and D. V. Fursa, *J. Chem. Phys.* **161**, 244304 (2024).

## GEOSH: Chemical Equilibrium Calculations of Ideal Gas Mixtures

Sarah M. Stangl<sup>1</sup>, Cristiano Nisoli<sup>1</sup>, Mark C. Zammit<sup>1</sup>, Eddy M. C. Timmermans<sup>1</sup>, Amanda J. Neukirch<sup>1</sup>, Christopher M. Mauney<sup>1</sup>, Jeffery A. Leiding<sup>1</sup>

<sup>1</sup>*(Presenting author underlined) Los Alamos National Laboratory*

Understanding the relative abundances of atoms, molecules, and ions in a gaseous mixture at equilibrium is necessary when modeling gaseous mixtures in a wide range of applications such as planetary and stellar atmospheres. To simplify the problem, we apply the law of mass action and Saha-Langmuir equations to express the chemical species in terms of the atomic species present in the system. By normalizing chemical species and eliminating electron capture and ionization degrees of freedom, we reduce the problem to a set of non-linear equations describing the conservation of each atomic nuclei and charge species. We introduce GEOSH, a code designed to solve for the relative concentrations of chemical species by solving a system of non-linear equations using a Newton-Raphson solver while determining the thermodynamic quantities of the system in equilibrium.

## 1D Imaging X-ray Spectroscopy for Inferring Time-Integrated, Spatially-Resolved Volumetric Temperature Distributions in Relativistic Laser–Solid Interactions

**Radka Štefaníková<sup>1,2,+</sup>, Thomas E. Cowan<sup>1,2</sup>, Kateřina Falk<sup>1</sup>, Thomas Kluge<sup>1</sup>, Michaela Kozlová<sup>1</sup>, Irene Prencipe<sup>1</sup>, Hans-Peter Schlenvoigt<sup>1</sup>, Ulrich Schramm<sup>1,2</sup>, Karl Zeil<sup>1</sup>, Josefine Metzkes-Ng<sup>1</sup> and Michal Šmíd<sup>1</sup>**

<sup>1</sup>*Helmholtz-Zentrum Dresden-Rossendorf, Dresden, Germany*

<sup>2</sup>*Technische Universität Dresden, 01062 Dresden, Germany*

<sup>+</sup>*Currently at Marvel Fusion GmbH, 80339 Munich, Germany*

An experimental approach for reconstructing volumetric temperature distributions in solid-density plasmas generated by ultra-relativistic laser–solid interactions is presented. In contrast to previous diagnostics that quantified lateral or longitudinal temperature profiles separately, our method enables the retrieval of time-integrated temperature distributions across both spatial dimensions simultaneously. The technique combines spatially resolved x-ray emission spectroscopy with a minimal radiation-transport model to estimate the spatial extent of temperature-dependent emission regions. Applied to laser-driven plasmas in micron-scale metallic foil targets, the analysis reveals a layered and anisotropic heating structure: the hottest regions remain localized near the laser-irradiated front surface, while emission from cooler plasma extends deeper into the target bulk. As the method relies solely on standard x-ray imaging spectroscopy, it is compatible with high-repetition-rate laser systems and offers a practical, scalable diagnostic for probing volumetric heating in high-energy-density and fusion-relevant plasmas.

## Vibrationally resolved core excitation of the $\text{CN}^+$ molecular ion

Maria Tatsch<sup>1</sup>, Anastasia Andreeva<sup>2</sup>, B. Michel Döhning<sup>1</sup>, Pierre-Michel Hillenbrand<sup>3</sup>, Michael Martins<sup>2</sup>, Alfred Müller<sup>1</sup>, Simon Reinwardt<sup>2</sup>, Jörn Seltmann<sup>4</sup>, Julius Schwarz<sup>2</sup>, Florian Trinter<sup>5</sup>, Stefan Schippers<sup>1</sup>

<sup>1</sup>*I. Physikalisches Institut, Justus-Liebig-Universität Gießen*

<sup>2</sup>*Institut für Experimentalphysik, Universität Hamburg*

<sup>3</sup>*GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt*

<sup>4</sup>*Deutsches Elektronen-Synchrotron DESY, Hamburg*

<sup>5</sup>*Fritz-Haber-Institut der Max-Planck-Gesellschaft, Berlin*

Near-edge x-ray absorption fine structure spectroscopy (NEXAFS) is a powerful tool to probe electronic core excitations. We measured the electronic structure of the  $\text{CN}^+$  ion vibrationally resolved to gain insights towards accurate descriptions of core-excited diatomic molecules. So far, there is no other experimental data on the  $\text{CN}^+$  ion, consequently theoretical data is also scarce. First theoretical studies of the core excitation of the C atom of  $\text{CN}^+$  were published only recently [1].

The experiment took place at the Photon Ion end station at PETRA III (PIPE) [2]. The PIPE setup is a permanently installed user facility at the XUV beamline (P04) of the PETRA III synchrotron radiation facility operated by DESY in Hamburg. The brilliant synchrotron light of PETRA III enables vibrationally resolved core-excitation spectroscopy and provides access to high resolution measurements. The production of  $\text{CN}^+$  ions is quite challenging due to its toxicity and also due to its chemical characteristics. The  $\text{CN}^+$  ions were produced in an electron cyclotron resonance ion source (ECR) from a mixture of CO and  $\text{N}_2$  gases [3]. The ion beam was mass-over-charge analyzed with a dipole magnet and the  $\text{CN}^+$  ions were selected and merged with the counter-propagating photon beam over a length of ca. 1.7 m. The collimated  $\text{CN}^+$  ion current in the interaction region was 2 nA. The photon energy was scanned across the carbon and nitrogen *K* edges.  $\text{CN}^{2+}$  product ions resulting from the photon-ion interaction were separated from the ion beam with a demerging magnet and detected with a single particle detector. The measured ion yields as a function of photon energy exhibit vibrationally resolved resonances associated with the excitation of a 1s electron to unoccupied molecular levels. On our poster, we will present the current status of our data analysis.

[1] J. Li et al., Phys. Rev. A 111, 052803 (2025).

[2] S. Schippers et al., X-Ray Spectrometry 49, 11 (2020).

[3] D. S. Belic et al., Phys. Rev. A 95, 052702 (2017).

## Initial predictive modeling of plasma-wall interactions using ERO2.0 for W7-X with tungsten wall divertor

Georgii Timkovskii<sup>1</sup>, Juri Romazanov<sup>1</sup>, Sebastijan Brezinsek<sup>1,2</sup>, Daniil Ryndyk<sup>1</sup>, Henri Kumpulainen<sup>1</sup> and W7-X Team<sup>3</sup>

<sup>1</sup> *Forschungszentrum Jülich GmbH, IFN-1 - Plasma Physics, Jülich, Germany*

<sup>2</sup> *Mathematisch-Naturwissenschaftliche Fakultät, HHU Düsseldorf, Düsseldorf, Germany*

<sup>3</sup> *Max-Planck-Institut für Plasmaphysik, 17491 Greifswald, Germany*

Plasma-wall interaction (PWI) is a critical aspect of fusion device operation, influencing material lifetime, fuel retention, and overall plasma performance. In addition to interpretive modelling used to analyze and understand physics experiments, predictive modeling with validated codes like ERO2.0 becomes essential for anticipating system behavior and guiding design or operational decisions as required for an exchange of the wall material in Wendelstein 7-X (W7-X).

ERO2.0 is a fully kinetic Monte Carlo code dealing with PWI processes at the surface and incorporating phenomena such as drifts, ionization, and impurity-ion collisions in the plasma. While there have been multiple ERO2.0 studies of stellarators - including cases with some tungsten components - no systematic investigation exist for a fully metallic stellarator with a tungsten divertor.

In this work, PWI and impurity transport for the W7-X stellarator are studied with focus on the planned wall material exchange from a carbon wall to a tungsten wall. The influence of the resolution of existing 3D grids on the resulting PWI and impurity transport is analyzed.


## Machine Learning-Assisted Diagnostics of Laser-Induced Aluminium Plasma

Akash Uniyal, Lalita Sharma

*Department of Physics, IIT Roorkee, Roorkee 247667, India*

Laser-induced plasmas (LIPs) are gaining special attention nowadays due to their unique physical and chemical characteristics. These plasmas exhibit a transient nature with their properties varying with space and time. On one hand, where their versatile nature makes them suitable for a wide range of applications, at the same time this brings challenges in their accurate diagnostics [1]. The optimization of these plasmas for certain applications involves the correct determination of the plasma parameters. Techniques like laser induced breakdown spectroscopy (LIBS) when coupled with theoretical collisional-radiative (CR) modelling approach serves as a reliable way to diagnose these parameters accurately [2]. However, the real-time diagnostics is often infeasible with conventional CR modelling approach as these models are computationally intensive.

In light of this, people are approaching towards data driven methodologies namely machine learning (ML) for fast diagnostics of these plasmas [3, 4, 5, 6]. Due to the availability of open source softwares and resources and enhancement in data acquisition techniques, ML based approaches serves as efficient tools in the field of plasma spectroscopy. ML models once trained with CR model generated synthetic spectral intensity data can then be employed for rapid estimation of plasma parameters. In this context, we have developed two supervised learning based ML models namely – Random Forest (RF) and Deep Neural Network (DNN) for aluminium (Al) plasma. These models are trained and tested with CR model simulated spectral intensity data and then employed for predicting the electron temperature from experimental observables. Hyperparameter optimization is done using random search strategy to enhance predictive performance. The accuracy of these models is evaluated by measuring performance functions including MSE and  $R^2$  values. The electron temperature values as predicted by CR model and ML models are compared showing reasonably good agreement also shown in the attached figure along with this abstract. Details of the models will be discussed in the poster presentation. Overall, this study highlights the potential of ML based methods as an effective way for real-time diagnostics of non-equilibrium plasmas.



- [1] S. S. Harilal, M. C. Phillips, D. H. Froula, K. K. Anoop, R. C. Issac, and F. N. Beg, *Rev. Mod. Phys.* 94, 035002 (2022).
- [2] N. Ghosh, B. R. Geethika, J. Thomas, and L. Sharma, *Plasma Sources Sci. Technol.* 34, 055015 (2025).
- [3] A. N. Bakhtiyari, Y. Wu, D. Qi, and H. Zheng, *Optik* 272, 170297 (2023).
- [4] P. S. N. S. R. Srikar, I. Suresh, and R. K. Gangwar, *Spectrochim. Acta B* 215, 106909 (2024).
- [5] I. Suresh, P. S. N. S. R. Srikar, and R. K. Gangwar, *Phys. Plasmas* 31, 113501 (2024).
- [6] P. S. N. S. R. Srikar, I. Suresh, and R. K. Gangwar, *J. Phys. D: Appl. Phys.* 58, 415204 (2025).

## Abstract for Joint ICTP-IAEA School on Atomic and Molecular Process in Plasmas

**T. Van Hoomissen<sup>1</sup>, H. R. Hasson<sup>2</sup>, D. Schneider<sup>1</sup>, K. Chandler<sup>2</sup>, F. Kraus<sup>3</sup>, P. W. Moloney<sup>4</sup>, P. Hartigan<sup>5</sup>, C. C. Kuranz<sup>6</sup>, D. B. Schaeffer<sup>1</sup>**

<sup>1</sup>(Presenting author underlined) University of California, Los Angeles

<sup>2</sup>Sandia National Laboratories

<sup>3</sup>Princeton Plasma Physics Laboratory

<sup>4</sup>Imperial College London

<sup>5</sup>Rice University

<sup>6</sup>University of Michigan

This document is the template for contributed abstracts (posters and/or talks) and gives the guidelines for preparing and submitting abstracts for the *Joint ICTP-IAEA School on Atomic and Molecular Processes in Plasmas* to be held online from 4 May 2026 to 8 May 2026.

Time-resolved spectroscopy is a critical diagnostic for probing the evolution of plasma conditions in high-energy-density experiments [1]. The streaked visible spectroscopy (SVS) diagnostic at Sandia National Laboratories' Z Machine provides temporally and spatially resolved measurements of plasma emission, enabling inference of fundamental plasma parameters such as electron temperature and density. In this work, we analyze SVS data from the MagShockZ platform, which studies high-Mach-number quasi-perpendicular magnetized collisionless shocks. Using PrismSPECT, a collisional-radiative spectral analysis code, we model emission spectra across a range of electron temperatures and densities and compare the simulated spectra to the experimental measurements [2]. We present the first results of the electron temperature and density evolution of the background aluminum wire array plasma, and we compare the results to the MARZ campaign, which featured similar aluminum wire array plasmas.

[1] J. L. Porter, M. A. Gilmore, and J. E. Bailey, *Rev. Sci. Instrum.* 88, 123501 (2017).

[2] J. Colgan, M. W. C. Dharma-wardana, and F. G. Serpa, *High Energy Density Phys.* 26, 12 (2018).

Title: Streaked Visible Spectroscopy Analysis for Laser-Driven Collisionless Shocks on Z

## Characterisation of Neutral–Plasma Interactions for Gas Puff Imaging Using the DEGAS-2 Code

**Ruchi Varshney<sup>1,2</sup>, Ritu Dey<sup>3</sup>, and D. Raju<sup>1,2</sup>**

<sup>1</sup>*Institute for plasma Research, Bhal Village, Gandhinagar, Gujrat, 382428, India*

<sup>2</sup>*Homi Bhabha National Institute, Anushaktinagar, Mumbai, Maharashtra, 400094, India*

<sup>3</sup>*Department of physics, Indian Institute of Technology Tirupati, Yerpedu, 517619, India*

The edge plasma of a tokamak is well suited for studying turbulence because its lower temperatures and densities allow the use of diagnostics based on atomic and molecular processes, such as Gas Puff Imaging (GPI) [1]. Understanding turbulence in this region is important since it influences the boundary conditions of the core plasma.

Using DEGAS-2 [2,3], we examine how injected neutrals interact with the plasma and how the visible emission varies with puff rate. The puffed neutrals remain non-perturbative but produce much stronger emission than background neutrals, highlighting the importance of the edge and SOL region. Our results show that excited atoms created through molecular dissociation are a major source of the observed photons, supporting improved interpretation of edge turbulence measurements. We also present a comparative study of helium and hydrogen puffing to examine differences in their emissivity.

[1] S.J. Zweben et al., Phys. Plasmas 9 (2002) 1981

[2] D. P. Stotler, B. LaBombard, J. L. Terry, and S. J. Zweben, J. Nucl. Mater. 313–316, 1066 (2003).

[3] D.P. Stotler, C.F.F. Karney, Contribution Plasma Phys. 34

## Self-Energy Line Broadening in Dense Plasmas

**Jackson R. White<sup>1,2</sup>, Thomas A. Gomez<sup>3,4</sup>, and Charles E. Starrett<sup>2</sup>**

<sup>1</sup>*Department of Astronomy, University of Texas at Austin, Austin, TX 78712, USA*

<sup>2</sup>*X Computational Physics Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

<sup>3</sup>*Department of Nuclear and Radiation Sciences, University of Michigan, Ann Arbor, MI 48109, USA*

<sup>4</sup>*Department of Astrophysical and Planetary Sciences, University of Colorado, Boulder, CO 80305, USA*

Spectral line shapes describe the distribution of opacity from bound-bound electronic transitions, and are sensitive to different broadening mechanisms. In many dense laboratory and astrophysical plasmas the dominant broadening mechanism is pressure broadening, or Stark broadening, caused by collisions between radiating atoms and neighboring plasma particles [1]. Stark-broadened line shape models traditionally begin from an isolated-atom picture, and account for plasma effects perturbatively. This isolated-atom approach breaks down at high (near-solid) plasma densities, where many-body effects become more important. In this poster we present a new approach to modeling electron Stark broadening in dense plasmas, using self-energy calculations from configuration-resolved average atoms with integer-occupied core electron orbitals. Using the Excited States Method [2], we move beyond the isolated-atom picture by incorporating plasma effects self-consistently using density functional theory (DFT) techniques. We also demonstrate how this self-energy approach allows for dynamic screening, in contrast to traditional line shape theory that relies on static Debye screening. Finally, we compare our new line shape calculations to semi-analytic and simulation-based line shape codes for a range of different plasma conditions, and discuss applications of this work to recent high-energy-density laboratory opacity experiments.

[1] T. Gomez, T. Nagayama, P. Cho, D. Kilcrease, C. Fontes, M. Zammit, *J. Phys. B* **55**, 034002 (2022).

[2] C. Starrett, T. Thelen, C. Fontes, D. Rehn, *Phys. Rev. E* **109**, 035201 (2024).

# Dielectronic Recombination Study on $\text{Sr}^{4+}$ , $\text{Se}^{4+}$ , $\text{Ga}^{4+}$ and $\text{In}^{4+}$ Ions

Yanlan Xu<sup>1</sup>, Xiaobin Ding<sup>1,2,3</sup>, and Chenzhong Dong<sup>1,2,3</sup>

<sup>1</sup> Key Laboratory of Atomic and Molecular Physics and Functional Materials of Gansu Province, College of Physics and Electronic Engineering, Northwest Normal University, Lanzhou, 730070, China

<sup>2</sup> Gansu International Scientific and Technological Cooperation Base of Laser plasma Spectroscopy, Lanzhou, 730070, China

<sup>3</sup> Gansu Provincial Research Center for Basic Disciplines of Quantum Physics, Lanzhou, 730070, China

Dielectronic recombination (DR) is a crucial process in astrophysical plasmas and laboratory fusion devices, directly regulating plasma ionization balance, energy transport, and radiative spectral properties[1]. Löbbling et al.[2] analyzed ultraviolet spectra from the BD−22°3467 of a DAO-type white dwarf, and reported the first detection of spectral lines from trans-iron elements. The intense radiative field drives the extreme enrichment of these elements in the stellar atmosphere, and the DR processes of their ionized states play a crucial role in regulating the stellar atmosphere. Given the higher abundances of Sr, Se, Ga, and In, we calculated the DR cross sections of 4+ ions of these element.




Figure 1: DR cross section for  $\text{Sr}^{4+}$ ,  $\text{Se}^{4+}$ ,  $\text{Ga}^{4+}$  and  $\text{In}^{4+}$

Firstly, using the Flexible Atomic Code based on the Dirac-Fock-Slater method[3], we computed the energy levels, radiative transition rates, and autoionization rates for these four ions. Next, under the approximations of isolated resonance and independent processes, we calculated the DR cross sections with the Voigt profile, where the full width at half maximum (FWHM) of the Gaussian broadening was set to 10 meV.

Figure 1 shows the DR cross sections of these four ions. The DR cross section of  $\text{Sr}^{4+}$  is the largest (on the order of  $10^{-17}\text{cm}^2$ ), while those of the other three ions are of the order of  $10^{-18}\text{cm}^2$ . The DR cross section of each ion exhibits distinct density in different energy intervals; e.g.,  $\text{Sr}^{4+}$  shows density in 20–35 eV and  $\text{In}^{4+}$  in 40–60 eV. Among them, the strong resonance peaks with large cross sections significantly enhance the intensity of recombination reactions in this energy segment, thereby increasing the occurrence probability of electron recombination processes within this energy interval. In subsequent work, we will calculate their DR rate coefficients and analyze their impacts on the plasma.

[1] A. Burgess, ApJ, **139**, 776 (1964).

[2] L. Löbbling, M. A. Maney, et al., MNRAS, **492**, 528-548 (2020).

[3] M.F. Gu, Can. J. Phys. **86**, 675-689 (2008).