

A diagrammatic approach to composite, rotating impurities.

G. Bighin and M. Lemeshko

Institute of Science and Technology Austria


Trieste, July 4th, 2017

Impurity problems

Definition: one (or a few particles) interacting with a many-body environment.

How are the properties of the particle modified by the interaction?

Still $\mathcal{O}(10^{23})$ degrees of freedom...




Impurity problems

Definition: one (or a few particles) interacting with a many-body environment.

How are the properties of the particle modified by the interaction?

Still $\mathcal{O}(10^{23})$ degrees of freedom...


Quasiparticle description?



From impurities to quasiparticles

Structureless impurity: translational degrees of freedom/linear momentum exchange with the bath.

Most common cases: electron in a solid, atomic impurities in a BEC.



From impurities to quasiparticles

Structureless impurity: translational degrees of freedom/linear momentum exchange with the bath.

Most common cases: **electron in a solid**, atomic impurities in a BEC.




Image from: F. Chevy, Physics **9**, 86.

From impurities to quasiparticles

Structureless impurity: translational degrees of freedom/linear momentum exchange with the bath.

Most common cases: electron in a solid, **atomic impurities in a BEC**.




Image from: F. Chevy, Physics **9**, 86.

From impurities to quasiparticles

Structureless impurity: translational
degrees of freedom
momentum

This scenario can be formalized in terms of
quasiparticles using the **polaron**.

Most common cases: electron in a
solid, **atomic impurities in a BEC**.




Image from: F. Chevy, Physics **9**, 86.


From impurities to quasiparticles

Structureless impurity: translational degrees of freedom, no internal degrees of momentum

This scenario can be formalized in terms of **quasiparticles** using the **polaron**.

Most common cases: electron in a solid, **atomic impurities in a BEC**.

Image from: F. Chevy, Physics **9**, 86.



Composite impurity: translational *and* internal (i.e. rotational) degrees of freedom/linear and angular momentum exchange.

From impurities to quasiparticles

Structureless impurity: translational degrees of freedom, momentum

This scenario can be formalized in terms of **quasiparticles** using the **polaron**.

Most common cases: electron in a solid, **atomic impurities in a BEC**.

Image from: F. Chevy, Physics **9**, 86.

What about a **rotating particle**? Can there be a **rotating analogue of the polaron quasiparticle**? The main difficulty: the **non-Abelian $SO(3)$ algebra** describing rotations.


and
f
entum

The angulon

A composite impurity in a bosonic environment can be described by the angulon Hamiltonian^{1,2,3,4} (angular momentum basis: $\mathbf{k} \rightarrow \{k, \lambda, \mu\}$):

$$\hat{H} = \underbrace{\underbrace{\hat{\mathbf{B}}\hat{\mathbf{J}}^2}_{\text{molecule}}}_{\text{phonons}} + \underbrace{\sum_{k\lambda\mu} \omega_k \hat{b}_{k\lambda\mu}^\dagger \hat{b}_{k\lambda\mu}}_{\text{phonons}} + \underbrace{\sum_{k\lambda\mu} U_\lambda(k) \left[Y_{\lambda\mu}^*(\hat{\theta}, \hat{\phi}) \hat{b}_{k\lambda\mu}^\dagger + Y_{\lambda\mu}(\hat{\theta}, \hat{\phi}) \hat{b}_{k\lambda\mu} \right]}_{\text{molecule-phonon interaction}}$$

- Linear molecule.
- Derived rigorously for a molecule in a weakly-interacting BEC¹.
- Phenomenological model for a molecule in any kind of bosonic bath³.



¹R. Schmidt and M. Leshchko, Phys. Rev. Lett. **114**, 203001 (2015).

²R. Schmidt and M. Leshchko, Phys. Rev. X **6**, 011012 (2016).

³M. Leshchko, Phys. Rev. Lett. **118**, 095301 (2017).

⁴Y. Shchadilova, "Viewpoint: A New Angle on Quantum Impurities", Physics **10**, 20 (2017).

The angulon


A composite impurity in a bosonic environment can be described by the angulon Hamiltonian^{1,2,3,4} (angular momentum basis: $\mathbf{k} \rightarrow \{k, \lambda, \mu\}$):

$$\hat{H} = \underbrace{\hat{B}\hat{J}^2}_{\text{molecule}} + \underbrace{\sum_{k\lambda\mu} \omega_k \hat{b}_{k\lambda\mu}^\dagger \hat{b}_{k\lambda\mu}}_{\text{phonons}} + \underbrace{\sum_{k\lambda\mu} U_\lambda(k) \left[Y_{\lambda\mu}^*(\hat{\theta}, \hat{\phi}) \hat{b}_{k\lambda\mu}^\dagger + Y_{\lambda\mu}(\hat{\theta}, \hat{\phi}) \hat{b}_{k\lambda\mu} \right]}_{\text{molecule-phonon interaction}}$$

This talk: toy potential. Can be connected to real PESs³.

Model for a molecule in a weakly-interacting BEC¹.

- Phenomenological model for a molecule in any kind of bosonic bath³.



¹R. Schmidt and M. Lemeshko, Phys. Rev. Lett. **114**, 203001 (2015).

²R. Schmidt and M. Lemeshko, Phys. Rev. X **6**, 011012 (2016).

³M. Lemeshko, Phys. Rev. Lett. **118**, 095301 (2017).

⁴Y. Shchadilova, "Viewpoint: A New Angle on Quantum Impurities", Physics **10**, 20 (2017).

Composite impurities and where to find them

Strong motivation for the theoretical study of composite impurities comes from many different fields. Composite impurities are realized as:

- **Molecules** embedded into **helium nanodroplets** (rotational spectra, rotational constant renormalization).




Image from: J. P. Toennies and A. F. Vilesov, *Angew. Chem. Int. Ed.* **43**, 2622 (2004).

Composite impurities and where to find them

Strong motivation for the theoretical study of composite impurities comes from many different fields. Composite impurities are realized as:

- **Molecules** embedded into **helium nanodroplets** (rotational spectra, rotational constant renormalization).




Image from: J. P. Toennies and A. F. Vilesov, *Angew. Chem. Int. Ed.* **43**, 2622 (2004).


Composite impurities and where to find them

Strong motivation for the theoretical study of composite impurities comes from many different fields. Composite impurities are realized as:

- Molecules embedded into helium nanodroplets (rotational constant renormalization).

Rotational spectrum

Renormalized lines (smaller effective B)



Gas phase


in ^4He

Image from: J. P. Toennies and A. F. Vilesov, *Angew. Chem. Int. Ed.* **43**, 2622 (2004).

Composite impurities and where to find them

Strong motivation for the theoretical study of composite impurities comes from many different fields. Composite impurities are realized as:

- **Molecules** embedded into **helium nanodroplets** (rotational spectra, rotational constant renormalization).
- **Ultracold molecules** and ions.




B. Midya, M. Tomza, R. Schmidt, and M. Lemesko,
Phys. Rev. A **94**, 041601(R) (2016).

Composite impurities and where to find them

Strong motivation for the theoretical study of composite impurities comes from many different fields. Composite impurities are realized as:

- **Molecules** embedded into **helium nanodroplets** (rotational spectra, rotational constant renormalization).
- **Ultracold molecules** and ions.
- Electronic excitations in **Rydberg atoms**.




Pfau group, Nature **502**, 664 (2013).

Composite impurities and where to find them

Strong motivation for the theoretical study of composite impurities comes from many different fields. Composite impurities are realized as:

- **Molecules** embedded into **helium nanodroplets** (rotational spectra, rotational constant renormalization).
- **Ultracold molecules** and ions.
- Electronic excitations in **Rydberg atoms**.
- Angular momentum transfer from the **electrons** to a **crystal lattice**.

Path integral description for the angulon



Main reference: GB and M. Lemesko, arXiv:1704.02616

Path integral description for the angulon

The path integral in QM describes the transition amplitude between two states with a weighted average over all trajectories, S is the classical action.

$$G(x_i, x_f; t_f - t_i) = \langle x_f, t_f | x_i, t_i \rangle = \int \mathcal{D}x \, e^{iS[x(t)]}$$



Path integral description for the angulon

The **angulon's Green function** is calculated in the same way. We need

- Molecular coordinates: two **angles** (θ, ϕ) describing the orientation of the molecule.
- An infinite number of **harmonic oscillators** $b_{k\lambda\mu}$ to describe the bosonic bath.

$$G(\theta_i, \phi_i \rightarrow \theta_f, \phi_f; T) = \int \mathcal{D}\theta \mathcal{D}\phi \prod_{k\lambda\mu} \mathcal{D}b_{k\lambda\mu} e^{i(S_{\text{mol}} + S_{\text{bos}} + S_{\text{mol-bos}})}$$

Path integral description for the angulon

The **angulon's Green function** is calculated in the same way. We need

- Molecular coordinates: two **angles** (θ, ϕ) describing the orientation of the molecule.
- An infinite number of **harmonic oscillators** $b_{k\lambda\mu}$ to bosonic bath.

Derived from the Hamiltonian

$$G(\theta_i, \phi_i \rightarrow \theta_f, \phi_f; T) = \int \mathcal{D}\theta \mathcal{D}\phi \prod_{k\lambda\mu} \mathcal{D}b_{k\lambda\mu} e^{i(S_{\text{mol}} + S_{\text{bos}} + S_{\text{mol-bos}})}$$

Path integral description for the angulon

The **angulon's Green function** is calculated in the same way. We need

- Molecular coordinates: two **angles** (θ, ϕ) describing the orientation of the molecule.
- An infinite number of **harmonic oscillators** $b_{k\lambda\mu}$ to describe the bosonic bath.

$$G(\theta_i, \phi_i \rightarrow \theta_f, \phi_f; T) = \int \mathcal{D}\theta \mathcal{D}\phi \prod_{k\lambda\mu} \mathcal{D}b_{k\lambda\mu} e^{i(S_{\text{mol}} + S_{\text{bos}} + S_{\text{mol-bos}})}$$

Critically the environment ($b_{k\lambda\mu}$) can be **integrated out exactly**

$$G(\theta_i, \phi_i \rightarrow \theta_f, \phi_f; T) = \int \mathcal{D}\theta \mathcal{D}\phi e^{iS_{\text{eff}}[\theta(t), \phi(t)]}$$



and included in an effective action S_{eff} .

Path integral description for the angulon

A closer look at the effective action:

$$S_{\text{eff}} = \underbrace{\int_0^T dt B \mathbf{J}^2}_{S_0} + \underbrace{\frac{i}{2} \int_0^T dt \int_0^T ds \sum_{\lambda} P_{\lambda}(\cos \gamma(t, s)) \mathcal{M}_{\lambda}(|t - s|)}_{S_{\text{int}}}$$

- A term describing a **free molecule** $\sim B \mathbf{J}^2$.
- A **memory term** accounting for the many-body environment, a function of the angle $\gamma(t, s)$ between the angulon position at different times.



Path integral description for the angulon



A closer look at the effective action:

Legendre polynomials


Memory kernel

$$S_{\text{eff}} = \underbrace{\int_0^T dt \mathbf{B} \mathbf{J}^2}_{S_0} + \underbrace{\frac{i}{2} \int_0^T dt \int_0^T ds \sum_{\lambda} P_{\lambda}(\cos \gamma(t, s)) \mathcal{M}_{\lambda}(|t - s|)}_{S_{\text{int}}}$$

- A term describing a **free molecule** $\sim \mathbf{B} \mathbf{J}^2$.
- A **memory term** accounting for the many-body environment, a function of the angle $\gamma(t, s)$ between the angulon position at different times.



Path integral description for the angulon



- The many-body problem is reformulated in terms of a **self-interacting free molecule**.
- Time-non-local interaction (cf. Caldeira-Leggett, polaron, more generally: open quantum systems)
- The **interaction term** is very difficult to treat: it encodes exactly the many-body nature of the problem.


Diagrammatic theory of angular momentum in a many-body bath

We treat the interaction as a **perturbation**

$$G = \int \mathcal{D}\theta \mathcal{D}\phi e^{iS_0 + iS_{\text{int}}} = \int \mathcal{D}\theta \mathcal{D}\phi e^{iS_0} (1 + iS_{\text{int}} - \frac{1}{2}S_{\text{int}}^2 + \dots) = G^{(0)} + G^{(1)} + G^{(2)} + \dots$$

The result can be interpreted as a **diagrammatic expansion** (solid lines represent a free rotor, dashed lines are the interaction)

- $G^{(0)}(\theta_i, \phi_i \rightarrow \theta_f, \phi_f; T)$ is the Green's function for a free rotor 

- $G^{(1)}(\theta_i, \phi_i \rightarrow \theta_f, \phi_f; T)$ is the one-loop correction 

- $G^{(2)}(\theta_i, \phi_i \rightarrow \theta_f, \phi_f; T)$ is the two-loop correction



- and so on...

Angulon spectral function

Let us use the theory! The plan is simple:

1. Self-energy (Σ)
2. Dyson equation to obtain the angulon Green's function (G)
3. Spectral function (\mathcal{A})

Angulon spectral function

Let us use the theory! The plan is simple:

1. Self-energy (Σ)
2. Dyson equation to obtain the angulon Green's function (G)
3. Spectral function (\mathcal{A})

First order:

The diagram shows an equation: $\bigcirc \Sigma =$ followed by a horizontal line with three segments. The first segment is labeled $\lambda_1 \mu_1$ and has a dot at its right end. The second segment is a dashed arc labeled $\lambda_2 \mu_2$ above it, connecting the first dot to a second dot. The third segment is labeled $\lambda \mu$ and starts from the second dot.

Equivalent to a simple, 1-phonon variational Ansatz (cf. Chevy Ansatz for the polaron)

$$|\psi\rangle = Z_{LM}^{1/2} |0\rangle |LM\rangle + \sum_{\substack{k\lambda\mu \\ jm}} \beta_{k\lambda j} C_{jm,\lambda\mu}^{LM} b_{k\lambda\mu}^\dagger |0\rangle |jm\rangle$$

Angulon spectral function

Let us use the theory! The plan is simple:

1. Self-energy (Σ)
2. Dyson equation to obtain the angulon Green's function (G)
3. Spectral function (\mathcal{A})

Second order:


$$\textcircled{\Sigma} = \begin{array}{c} \lambda_1 \mu_1 \\ \curvearrowright \\ \lambda_2 \mu_2 \quad \lambda_3 \mu_3 \quad \lambda_4 \mu_4 \quad \lambda_5 \mu_5 \end{array} + \begin{array}{c} \lambda_{s1} \mu_{s1} \quad \lambda_{s3} \mu_{s3} \\ \curvearrowright \quad \curvearrowright \\ \lambda_2 \mu_2 \quad \lambda_4 \mu_4 \quad \lambda_5 \mu_5 \end{array}$$

Angulon spectral function

Let us use the theory! The plan is simple:

1. Self-energy (Σ)
2. Dyson equation to obtain the angulon Green's function (G)
3. Spectral function (\mathcal{A})

Dyson equation

$$\text{angulon} = \text{quantum rotor} + \text{many-body field} \circ \Sigma$$


Angulon spectral function

Let us use the theory! The plan is simple:


1. Self-energy (Σ)
2. Dyson equation to obtain the angulon Green's function (G)
3. Spectral function (\mathcal{A})

Finally the spectral function allows for a study the whole excitation spectrum of the system:

$$\mathcal{A}_\lambda(E) = -\frac{1}{\pi} \text{Im } G_\lambda(E + i0^+)$$


Angulon spectral function

Angulon **spectral function** as a function of the density:



Angulon spectral function


Angulon **spectral function** as a function of the density:



Key features:

1. Low density
2. Intermediate instability
3. High density

Angulon spectral function: low density




Density range: from ultra-cold atoms to superfluid helium.

Low density: **free rotor spectrum**, $E \sim L(L + 1)$.

Many-body-induced fine structure: upper phonon wing (one phonon with $\lambda = 0$, isotropic interaction).

Angulon spectral function: instability




Intermediate region: **angulon instability**.

Corresponding to the emission of a phonon with $\lambda = 1$ (due to anisotropic interaction).

Experimental observation: I. N. Cherepanov, M. Lemeshko, “Fingerprints of angulon instabilities in the spectra of matrix-isolated molecules”, arXiv:1705.09220.

Angulon spectral function: high density



High density: the **two-loop corrections** start to be relevant.

Conclusions

- The problem of angular momentum redistribution in a many-body environment has been treated through the **path integral formalism** and reformulated in terms of **diagrams**.
- It allows for a simple, compact derivation of angulon properties, including higher order terms.
- Future perspectives:
 - Dynamics.
 - Diagrammatic Monte Carlo.

Thank you for your attention.



Der Wissenschaftsfonds.

This work was supported by the
Austrian Science Fund (FWF), project
Nr. P29902-N27.

