



Voronoi Volume Function

A new probe of cosmology & galaxy evolution

Aseem Paranjape

Inter-University Centre for Astronomy & Astrophysics (IUCAA), Pune

High Energy Physics Seminar, ICTP

June 3, 2020

(in collaboration with Shadab Alam)

arXiv:2001.08760 (MNRAS in press)



Voronoi tessellation


Given N_{trc} tracers at positions $\{\mathbf{x}_t\}$ with $1 \leq t \leq N_{\text{trc}}$, the Voronoi tessellation is a partition of space into N_{trc} cells $\{\mathcal{C}_t\}$ such that, for a given tracer t , \mathcal{C}_t is the set of points closer to t than to any other t' .

— Used in various fields such as meteorology, epidemiology, geophysics, computational fluid dynamics (e.g. AREPO) etc.

— Several applications in cosmology too. E.g., cosmic web classification, void identification, etc.

This talk:

**Volume function of Voronoi cells
of 3d clustered tracers**



SciPy implementation of Voronoi tessellation of uniformly distributed 2d tracers.



Voronoi tessellation

simulation:

WMAP7 Λ CDM

tracers:


mass-thresholded
haloes at $z = 0$

tessellation:

Monte Carlo
algorithm

colour:

$$1 + \delta_{\text{trc}} = (n_{\text{trc}} V)^{-1}$$





Outline

- ◆ Definition of Voronoi volume function (VVF)
- ◆ Known results and analytical expectations
 - Connection to void probability function
- ◆ Results from simulations
 - Effects of cosmology, substructure and RSD
- ◆ Preliminary comparisons with GAMA results



Voronoi volume function

Definitions

If $V(t)$ is volume of cell \mathcal{C}_t containing tracer t then

$$\langle V \rangle = \frac{1}{N_{\text{trc}}} \sum_t V(t) = \frac{V_{\text{tot}}}{N_{\text{trc}}} = n_{\text{trc}}^{-1}$$

Define

$$y \equiv V / \langle V \rangle = n_{\text{trc}} V$$

We will denote the probability distribution $p(y)$ as the Voronoi volume function (VVF).

Clearly $\langle y \rangle = \int dy p(y) y = 1$.



Voronoi volume function

Uniformly distributed (Poisson) tracers

For **uniformly distributed (Poisson)** tracers, $\langle y^2 \rangle$ is known analytically [*Gilbert 1962*]

$$\langle y^2 \rangle_{\text{Poisson}} = \frac{8\pi^2}{3} \int_0^\infty dz z^2 \int_{-1}^1 d\mu \frac{1}{v(z, \mu)^2} \simeq 1.179$$

where $v(z, \mu) = \frac{\pi}{3} \left[2z^3 + 3\mu z(z^2 + 1) - (3\mu^2 z^2 + 1)z + 3(1 - \mu z)T + 2T^{3/2} \right]$
with $T = |z^2 + 1 - 2\mu z|$.

Although $p(y)$ is not known analytically, accurate fitting functions exist, e.g.:

$$p_{\text{Poisson}}(y) = \frac{c b^{a/c}}{\Gamma(b/c)} y^{a-1} \exp(-by^c)$$

with $a = 4.8065$, $b = 4.06342$, $c = 1.16391$ [*Tanemura 2003*]



Voronoi volume function

Clustered tracers

For **clustered** tracers, generalising [Gilbert 1962] we can write

$$\langle y^2 \rangle = \frac{8\pi^2}{3} \int_0^\infty dz z^2 \int_{-1}^1 d\mu \frac{1}{v(z, \mu)^2} n_{\text{trc}}^2 \int_0^\infty dV_U V_U \exp(W_0(n_{\text{trc}}, V_U))$$

where $\exp(W_0(n_{\text{trc}}, V))$ is the *void probability function* for the volume V [White 1979]:

$$W_0(n_{\text{trc}}, V) = \sum_{k=1}^{\infty} \frac{(-n_{\text{trc}} V)^k}{k!} \bar{\xi}_k(V) \equiv (-n_{\text{trc}} V) \chi(n_{\text{trc}}, V)$$

where $\bar{\xi}_k(V)$ is the connected k -point correlation function averaged over V .


Thus $\langle y^2 \rangle$ depends on the infinite hierarchy of tracer correlation functions.

[For Poisson distributed tracers, $\chi(n_{\text{trc}}, V) = 1$ and we recover $\langle y^2 \rangle_{\text{Poisson}} \simeq 1.179$.]



Simulation suite

Dark matter only GADGET-2 N-body runs



Cold Dark Matter

P13 ($\Omega_m = 0.315, h = 0.673, \sigma_8 = 0.829$)


L150_N512

WMAP7 ($\Omega_m = 0.276, h = 0.7, \sigma_8 = 0.811$)

L150_N1024, L300_N1024, L600_N1024

P18 ($\Omega_m = 0.306, h = 0.678, \sigma_8 = 0.815$)

L200_N1024



Alternate Dark Matter (paired to CDM)

WMAP7 warm DM ($m_{\text{DM}} = 0.4 \text{ keV}$)

L150_N1024




P18 ballistic DM ($z_* = 10^5, f_{\text{bdm}} = 0.5$)

L200_N1024






Simulation suite

WMAP7 cdm



WMAP7 wdm

P18 cdm




P18 bdm




VVF shape

Dependence on halo mass





tracers:
mass-thresholded
haloes in real space






VVF percentiles + std dev

Dependence on halo mass and redshift




tracers:
mass-thresholded haloes in real space








VVF percentiles + std dev

Dependence on halo clustering



[AP, Hahn & Sheth 2018]





tracers: (mass + bias)-thresholded haloes in real space at $z = 0$



VVF percentiles + std dev

Effects of substructure and redshift-space distortions




tracers: mass-thresholded (sub)haloes in real and redshift space at $z = 0$



Comparison with data

Subhalo abundance matching: CDM simulations




tracers: ($v_{\text{peak}} + \text{bias}$)-thresholded (sub)haloes in redshift space at $z = 0$ and $z = 0.1$



Comparison with data

Subhalo abundance matching: cosmology dependence





Conclusions

- ★ Voronoi volume function is a novel probe of nonlinear structure, sensitive to variety of tracer properties
 - Halo mass
 - Substructure content
 - Kinematics (RSD)
 - Large-scale clustering
 - Redshift evolution

- ★ Cosmology dependence (*cdm* / *wdm*):
 - VVF of mass-selected samples essentially universal
 - Weak cosmology dependence in realistic (SHAM) samples

- ★ Cosmology dependence (*bdm*):
 - Strong sensitivity to oscillatory features in $P(k)$