

Introduction to the lonosphere

Sandro M. Radicella

Regional Workshop on GNSS and Space Weather Rabat (Morocco), 9-13 May 2022

This lecture




- ✓ The atmospheric system
- ✓ Formation of the ionosphere
- ✓ Ionospheric structure
- ✓ Ionospheric variations



The atmospheric system

The Earth System





A System of Interacting Systems

The atmospheric system



We use a number of variables to describe the atmosphere:


- Temperature
- Composition and Mixing ratio
- Ionization



•••


Temperature





THERMOSPHERE: Temperature increases steadly with altitude because is heated mainly by absorption of EUV and XUV radiation through dissociation of molecular oxygen. Temperature is highly variable with time of day and solar activity.

MESOSPHERE: Temperature decreases with altitude because ozone density decreases faster than the increase of incoming radiation.




STRATOSPHERE: Temperature increases with altitude due to heating from the ozone which absorbs the solar ultra-violet radiation that penetrates down to these altitudes.

TROPOSHERE: Temperature decreases with altitude. Heated mainly by the ground, absorbs solar radiation and re-emits it in the infra-red.


Atmospheric composition:



Ground level



Above 100 Km



Mixing Ratio




Turbulent mixing: *lower and middle atmosphere*

- ✓ Does not depend on molecular weight
- ✓ Tends to be independent of height

Diffusion: *upper atmosphere*

- ✓ Mean molecular weight of mixture gradually decreases with height.
- ✓ Only lightest gases are present at higher levels.
- ✓ Each gas behaves as if it were
- ✓ alone.




Near 100km: diffusion = turbulent mixing.

Density drops-of exponentially with height

A closer look to the atmospheric pressure





Atmospheric Hydrostatic Equilibrium



Pressure Gradient:

$$\frac{dp}{dz} = -g(z)\rho$$

height derivative of pressure equals acceleration of gravity times density

Perfect Gas Law:

$$p = nkT = \frac{\rho}{M}kT$$

Approximation: If g and T are not functions of z, then:

$$\frac{dp}{dz} = -p\frac{Mg}{kT} = -\frac{p}{H}$$

$$H = \frac{kT}{Mg}$$

H = scale height (e-folding distance)


$$\frac{dp}{p} = -\frac{dz}{H}$$

$$p(z) = p(z_0) \exp\left[-\frac{z - z_0}{H}\right]$$

Scale Height



In various scientific contexts, a "Scale Height" is a distance over which a quantity decreases by a factor of e.



The scale height is the height at which the atmosphere would extend if it were all compressed into one of constant density (the rectangular area in the above diagram)



Formation of the ionosphere


Photochemical processes in the atmosphere



The atmosphere of the Earth is made up of a large number of chemical constituents.

Major constituents are N_2 , O_2 and Ar, but many more constituents are produced in the atmosphere by *photochemical processes of solar origin* or at the surface by different natural processes and human activity.

Photochemical processes play a fundamental role in the middle and upper atmosphere including the ionosphere.




Main photochemical absorption processes of solar radiation



Photodissociation
$$XY + hv \rightarrow X + Y$$

(wavelength > 130 nm)

Photoionization $X + hv \rightarrow X^+ + e$ (wavelength < 100 nm)



Ion species recombination processes



Radiative Recombination


 $X^+ + e \rightarrow X + hv (\alpha_R)$

Dissociative Recombination

 $XY^+ + e \rightarrow X + Y (\alpha_D)$

Ion-Ion Recombination

XY⁺ + Z⁻→neutrals (α₁)




Formation of the Ionosphere



Solar UV and X radiation impinges at angle χ_{ν} and a flux I_{∞} on the top of the atmosphere.

Solar radiation is absorbed in the upper atmosphere and ionizes the neutral atmosphere




Chapman layer theory



Named for Sydney Chapman, who first derived mathematically the shape of such a distribution.

The theory assumes:

- A monocromatic ionizing radiation from the sun,
- A single neutral constituent to be ionized distributed exponentially (i.e., with a constant scale height),
- Photochemical equilibrium



Basic equations of solar radiation absorption in the atmosphere (1)



- $H = k_B T_n / m_n g$, H is the neutral scale height, with g being the gravitational acceleration at height z = 0, where the density is n_0 .
- According to radiative transfer theory, the incident solar radiation / diminishes with altitude along the ray path in the atmosphere.
- σ_{ν} is the radiation absorption cross section for radiation (photon) of frequency ν .

$$n_n(z) = n_0 \exp(-z/H)$$

$$dI = \sigma_{\nu} n_n \frac{dz}{\cos \chi_{\nu}} I$$

Basic equations of solar radiation absorption in the atmosphere (2)



Solving for the intensity yields:

$$I(z) = I_{\infty} \exp \left[-\frac{\sigma_{\nu} n_0 H}{\cos \chi_{\nu}} \exp(-z/H) \right]$$


$$q_{\nu}(z) = \kappa_{\nu} \sigma_{\nu} n_0 I_{\infty} \exp \left[-\frac{z}{H} - \frac{\sigma_{\nu} n_0 H}{\cos \chi_{\nu}} \exp(-z/H) \right]$$

The photoionization rate per unit volume $q_{\nu}(z)$, is proportional to the ionization efficiency, κ_{ν} , and absorbed radiation:

$$q_{\nu}(z) = \kappa_{\nu} \sigma_{\nu} n_{n} I(z)$$

This equation describes the formation of the Chapman layer and represents the basis of the theory of the photochemical processes in the atmosphere.

- Radiation intensity decreases and neutral density increases with decreasing altitude.
- ✓ As a consequence ion production reaches a maximum and after that decreases, forming a layer with a clearly defined peak.



Basic equations of solar radiation absorption in the atmosphere (3)



Assuming dissociative recombination and quasi-neutrality ($n_e = n_i$), the continuity equation for n_e reads:

$$\frac{dn_e}{dt} = q_{v,e} - \alpha_T n_e^2$$



Ionospheric structure






Layered structure varying with time and solar activity due to different ionization production and loss processes


Transport processes become important in the F2 region and upper, including ambipolar diffusion and wind-induced drifts along B and electromagnetic drifts across B.


E and F1 regions behave as a Chapman layer dominated by photochemical processes.

At the E region heights sporadic thin layers can be formed with electron densities above the background values.

D region is characterized by the presence of negative ions due to the attachment of electrons to neutrals







Continuity equation and lon transport in the F region



Formed ions and electrons (P), tend to recombine (L) but are also affected by transport with a plasma drift V.

$$\frac{\partial n_e}{\partial t} = P - L - div(n_e V)$$

F region chemistry (1)



Above ca. 150 km ion-electron production is by EUV (10-90 nm)

$$O + h\nu \rightarrow O^+ + e$$

$$N_2 + h\nu \rightarrow N_2^+ + e$$

F region chemistry (2)



Recombination is a two-stages process:

$$O^+ + N_2 \rightarrow NO^+ + N$$
 (attachment like), rate $\propto \beta[O^+]$

This reaction controls the loss rate at high heights

$$NO^+ + e \rightarrow N + O$$
 rate $\propto \alpha [NO^+][n_e]$

This reaction controls the rate at low heights

$$\frac{dn_e}{dt} = q(z,\chi) - \alpha_D n_e^2 - \beta n_e$$

This is the continuity equation (without transport) for electron density

F region chemistry (3)



Continuity equations for the ionized species

$$\frac{dn_{e}}{dt} = q - \alpha_{D} n_{e}[NO^{+}]$$

$$\frac{d[O^{+}]}{dt} = q - \beta[N_{2}][O^{+}]$$

$$\frac{d[NO^{+}]}{dt} = \beta[N_{2}][O^{+}] - \alpha_{D} n_{e}[NO^{+}]$$

$$\beta[N_{1}] >> \alpha_{D} : [NO^{+}] >> [O^{+}] \Rightarrow a = \alpha n^{2}$$

$$\beta[N_2] >> \alpha n_e \; ; [NO^+] >> [O^+] \; \Rightarrow \; q = \alpha n_e^2$$

$$\beta[N_2] << \alpha n_e \ ; [NO^+] << [O^+] \ \Rightarrow \ q = \beta[N_2] n_e$$

F region chemistry (4)



Assuming photochemical equilibrium in the F region

$$q \propto I_{\infty}[O]$$

$$n_e = (q/\alpha)^{1/2} \propto (I_{\infty}[O])^{1/2}$$

At F2 heights

$$n_e \propto [O]/[N_2]$$

More exactly:

At F2 heights:
$$n_e \propto I_{\infty}[O]/[N_2]$$

This result is important to explain same aspects of the variability of the F2 electron density

NmF2 and [O]/[N2]



The annual and semi-annual amplitudes (normalized) and phases for NmF2 and [O/N2] at example stations

Yu, T. et al. (2004), Global scale annual and semiannual variuations of daytime NmF2 in the high solar activity years, J. of Atmos. and Solar-Terr. Physics 66, 1691-1701




Fig. 8. The variation for the daytime NmF2 at Canberra Station, Port Stanley Station and Singapore Station are respectively, shown in the upper panel of 8(a)-(c), and the [O/N2] at those stations are shown in the bottom panel.

High Latitude

NmF2 shows the same pattern as [O/N2].
NmF2 appears controlled by [O/ N2].

Middle Latitude

Port Stanley, NmF2 shows semi- annual variation and [O/N2] a dominant annual variation. NmF2 only partly controlled by the [O/N2].

Low Latitude

NmF2 and [O/N2] show semi-annual variation but not annual components.
[O/ N2] plus other mechanisms contribute to NmF2 variation.

Transport in the F region (I)



Ions and electrons are also affected by transport with a plasma drift V

$$\frac{dn_e}{dt} = q(z, \chi) - \beta [N_2] n_e - \frac{d(n_e V)}{dh}$$

W being the upward drift velocity

$$\frac{dn_e}{dt} = q(z,\chi) - \beta[N_2]n_e - \frac{d(n_eW)}{dh}$$

Electron density in the F region



Under day-time equilibrium conditions

- (1) $N_m F_2 \approx q_m / (\beta [N_2])_m$ (2) Below the peak: $N \approx q / \beta [N_2]$ (3) Well above the peak $N \propto e^{-z/2}$

F region structure in summary




- ✓ The lowest region (F1), where photochemistry dominates.
- ✓ A transition region from chemical to diffusion (lower F2).
- ✓ The upper region, or topside, where diffusion dominates
- ✓ In the F2 (including the topside) the presence of transport processes, influenced by the geomagnetic field, became important.



Ionospheric variations

Ionospheric variations





Two types of variations



Climate

Variations occurring in cycles.

Can be predicted with

reasonable accuracy

Weather

Variations mostly due to Solar induced Space Weather but also caused by coupling with lower atmosphere

Introduction to the topic




We will concentrate on the variations of the F2 layer through two parameters that are related to the peak electron density and the total electron content in the ionosphere.

The starting point will be a mention to experimental techniques used to derive these parameters.

The ionosonde and the ionogram



A radar that transmits pulses of a sweep of frequencies usually from 1 to up to 20 MHZ



As the frequency increases, the wave penetrates further in the ionosphere

Critical frequencies and virtual heights





The frequency at which the wave penetrates the layer foF2, foF1, foE

$$N = 0.124 \times 10^{11} (f)^{2}$$
$$N[m^{-3}], f[MHz]$$

lonosonde

vertical circuit





VIRTUAL HEIGHT

The height reached by radio wave if the group velocity is equal to the velocity of light in vacuum.


The presence of the ionosphere slows down the group velocity h'F2, h'F, h'E

Total electron content



The total electron content (TEC) is the total number of electrons along a path between a transmitter and a receiver

Can be obtained by different means, mainly from GNSS and satellite born altimeters



$$N_T = \int_s N(s)$$

1 TEC unit =
$$10^{16} m^{-2}$$

Diurnal, Seasonal and Solar Activity variations of foF2








Figure 1.5: Monthly median values of NmF2 as recorded at Rome in April, June, October and December for the years of maximum activity 1990-1991 (max of solar cycle 22) and 2001-2002 (max of solar cycle 23).



From: L. Perna, Ionospheric plasma response to the anomalous minimum of the solar cycle 23/24: modeling and comparison with IRI-2012

PhD Thesis, Università di Bologna, 2017






Day-to-day variability of NmF2


Variation of NmF2 at Slough for every day during four 2-month periods in 1973-1974.

H. Rishbeth, M. Mendillo / Journal of Atmospheric and Solar-Terrestrial Physics 63 (2001) 1661–1680




Global variations of foF2 April (R12=100)





Geographical variations: the equatorial anomaly





Nava B., Radicella S.M., Pulinets S. and Depuev V. "Modelling bottom and topside electron density and TEC with profile data from topside ionograms", Advances in Space Research, V. 27, pp. 31-34, 2001.

From the Chapman theory electron density should maximise over the geographic equator at equinox.

Actually it maximises 15-20 degrees of geomagnetic latitude N and S, with small minimum at the equator

Due to the presence of the geomagnetic field:
the 'fountain effect"

Equatorial Anomaly and the "fountain effect"




The Equatorial Electrojet drives the F-region behavior

E-field is zonal (along latitude lines)

Magnetic field is meridional (along longitudes)

Plasma drift is vertically upwards

Plasma
descends
down the
magnetic lines
N and S of the
geomagnetic
equator



The Equatorial Anomaly of the ionosphere and the "fountain effect"


Diurnal and Seasonal development of the Equatorial Anomaly



The monthly averaged equatorial ionospheric anomaly contour chart of vertical TEC in geographic latitude (Taiwan sector: 6.0 UT = 14.0 LT)

September 1996 - August 1997.

C.-C. Wu et al. / Journal of Atmospheric and Solar-Terrestrial Physics 66 (2004) 199–207




Vertical TEC diurnal and day-to-day variations (1)



TEC(10**16) ebre Lat=40.8N Lon=0.5E

Middle Latutude




GPS derived vertical TEC at 5 min interval for Roquetes (Lat. 40.8°, Lon. 0.5° E, Mag. Dip 57°), October 2000



Vertical TEC diurnal and day-to-day variations (2)


TEC(10**16) nklg Lat=00.4N Lon=9.7E

Low Latutude



GPS derived vertical TEC at 5 min interval for Libreville (Lat. 0.4° N, Long. 9.7° E, Mag. Dip –25°), October 2000.

Vertical TEC Meridional cross section and day-to-day Variations



Low latitude Ionospheric irregularities




Radar observations of equatorial night-time F region often reveal rising plumes or bubbles of decreased electron density that penetrate into the topside reaching very high altitudes.

The generation of rising bubbles is initiated by a seed perturbation at the bottomside of a rising F layer.

Ionospheric irregularities seen by different techniques




JULIA radar plumes



Hyosub Kil et al., J. Geophys. Res. 119, Issue 7, 2014

Ionogram spread-F



D 100 200 400 600 800 1000 1500 3000 [km] MUF 9.3 9.4 10.0 10.9 12.3 14.2 19.8 34.6 [MHz] 280fk128h 50 kHz 5.0 km / DGS-256 ASON 908 / 7.9 5 345.6 E

ShowIonogram v 1.0

TEC depletion and amplitude scintillation





Fig. 2. Example of the measurements at Cape Verde (18 October 2010). The top plot shows the TEC measurements and also the elevation angle of the satellite. Bottom plot shows S4 index measurements at same location. In this example the depletion depth is about 7 TECU.

V. V. Paznukhov et al., Ann. Geophys., 30, 675–682, 2012

Ionospheric Storms



A very complex phenomenon that needs at least a full lecture by itself



Nava, B., J. Rodríguez-Zuluaga, K. Alazo- Cuartas, A. Kashcheyev, Y. Migoya-Orué, S. M. Radicella, C. Amory-Mazaudier, and R. Fleury (2016), Middle- and low-latitude ionosphere response to 2015 St. Patrick's Day geomagnetic storm, J. Geophys. Res. Space Physics, 121, 3421–3438, doi:10.1002/2015JA022299.

Tropospheric induced ionospheric variations



From the paper:

"Day-to-day ionospheric variability due to lower atmosphere perturbations" by H.-L. Liu, V. A. Yudin, and R. G. Roble; GEOPHYSICAL RESEARCH LETTERS, VOL. 40, 665–670.

This study demonstrates that the thermosphere-ionosphere-mesosphere electrodynamics general circulation model (TIEGCM) constrained by the atmosphere community climate model (WACCM) simulations is capable of reproducing observed features of day-to-day variations in the F2 region at low latitudes.

LIU ET AL: DAY-TO-DAY IONOSPHERIC VARIABILITY





Figure 4. Daily values of NmF2 (gray), their mean values (black solid), and the standard deviation (dashed) for (a) 51.25° N/0 longitude, (b) equator/75°W, and (c) 51.25° S/57.5°W (all geographic). (d) Mean values (shades) and standard deviation (lines) of NmF2 for LT1300. Contour intervals: 2.5×10^{10} m⁻³.

Under constant solar minimum and geomagnetically quiet conditions the meteorological driving may contribute comparably with geomagnetic forcing to the ionospheric day-to-day variability.

Earthquake and Tsunami ionospheric variations





Ionospheric variations induced by the Tohoku-Oki earthquake and tsunami of March 11, 2011.

The map shows changes in the Total Electron Content and sea surface heights.

Image by NASA/JPL-Caltech.



This lecture gives only a pale idea of the complexity of the ionosphere but I hope it awakes more interest in this fascinating part of our environment.

