

Dissipation-driven Strange Metal Behavior in High- T_c Superconducting Cuprates

S. Caprara¹, C. Di Castro¹, M. Grilli¹, G. Seibold²

¹*Dipartimento di Fisica, Università di Roma 'Sapienza', 00185, Rome, Italy*

²*Institut für Physik, BTU Cottbus-Senftenberg, D-03013 Cottbus, Germany*

Besides the mechanism responsible for high critical temperature superconductivity, the grand unresolved issue of the cuprates is the occurrence of a strange metallic state above the so-called pseudogap temperature T^* . Even though such state has been successfully described within a phenomenological scheme, the so-called Marginal Fermi-Liquid theory, a microscopic explanation is still missing. However, recent resonant X-ray scattering experiments [1] identified a new class of charge density fluctuations characterized by low characteristic energies and short correlation lengths, which are related to the well-known charge density waves. These fluctuations are present over a wide region of the temperature-vs-doping phase diagram and extend well above T^* . A recent investigation showed that charge density fluctuations can explain the strange metal phenomenology [2]. Therefore, charge density fluctuations are likely the long-sought microscopic mechanism underlying the peculiarities of the metallic state of cuprates. A more recent analysis [3] also showed that the damping of these fluctuations can cast in a single and new framework the linear-in- T resistivity of cuprates, sometimes extending down to very low T , and the surprising low-temperature divergence of the specific heat occurring at a critical doping p^* slightly larger than the optimal doping. Finally, an even more recent analysis [4] shows that in 2D the interplay between overdamped charge density fluctuations and diffusive electron modes due to quenched disorder can interplay to provide a microscopic generic mechanism for the above phenomenology. This scheme puts in a completely different light the strange metal issue and provides a general perspective to account for the strange metal behavior in many other system near quantum critical points, like heavy fermions, ruthenates, iron-based superconductors, and twisted bilayer graphene.

References

- [1] R. Arpaia, S. Caprara, R. Fumagalli, G. De Vecchi, Y.Y. Peng, E. Andersson, D. Betto, G.M. De Luca, N.B. Brookes, F. Lombardi, M. Salluzzo, L. Braicovich, C. Di Castro, M. Grilli, and G. Ghiringhelli, *Science* **365**, 906 (2019)
- [2] G. Seibold, R. Arpaia, Y. Y. Peng, R. Fumagalli, L. Braicovich, C. Di Castro, M. Grilli, G. Ghiringhelli, and S. Caprara, *Commun. Phys.* **4**, 7 (2021).
- [3] S. Caprara, C. Di Castro, G. Mirarchi, G. Seibold, and M. Grilli, *Commun. Phys.* **5**, 1 (2022).
- [4] S. Caprara, C. Di Castro, M. Grilli, and G. Seibold, arXiv:2205.10876