

"COLLEGE ON SOIL PHYSICS"

12 March - 6 April 2001

SOIL STRUCTURE

M. Pagliai

Istituto Sperimentale per lo Studio
e la Difesa del Suolo
Firenze, Italy

These notes are for internal distribution only

SOIL STRUCTURE

M. Pagliai

Istituto Sperimentale per lo Studio e la Difesa del Suolo. Piazza D'Azeglio 30, 50121 Firenze,
Italy

Definition

Soil structure may be defined either as "the shape, size and spatial arrangement of individual soil particles and clusters of particles (aggregates)" or as "the combination of different types of pores with solid particles (aggregates)". Soil structure has generally been defined in the former way and measured in terms of aggregate characteristics. These can be related to plant growth only empirically. In fact, it is the pore shape, the pore size distribution and the pore arrangement which affect many of the most important processes in soil that influence plant developments such as storage and movement of water and gases, solute movements and ease of root growth. For this reason measurements of pore space are increasingly being used to characterize soil structure. In fact, between the particles arranged singly or in aggregates, there is an intricate system of pore spaces on which plant roots, micro-organisms and soil fauna depend for the storage and movement of water and air.

Genesis of soil structure

Soil structure is a dynamic property and it is subjected to genesis and degradation processes. The main factors that affect the genesis of soil structure are represented by the effect of cations, by the interaction between clay particles under the influence of soil water content (wetting and drying cycles) and temperature, by the effect of organic matter, which is the main agent of aggregate stabilization, by root growth and by the action of soil macro and micro-organisms. Possible steps leading to the genesis of soil structure can be summarized as follows: flocculation of soil particles, microaggregation, aggregation and stabilization of aggregates. The main factors that affect the degradation of soil structure are the long-term intensive cultivations, which deplete the soil organic matter content, soil erosion, soil compaction, the formation of surface crusts and the formation of a compacted layer along the soil profile (e.g. ploughpan). Possible steps leading to the degradation of soil structure can be summarized as follows: destabilization or mechanical destruction of soil aggregates, disaggregation, microdisaggregation and dispersion of soil particles.

The genesis of soil structure depends on the presence of cementing substances, even though a structure can sometimes be recognized also in the absence of cementing substances

(Sequi, 1978). In many clay soils, for example, water can be naturally drained in the cracks formed during the drying processes following the shrinkage of the inorganic component of the soil and such cracks can never completely close during the year. However, the laboratory analysis of these types of soils reveals a low value of permeability. The alternation of wetting and drying cycles can be, therefore, an important mechanism of the genesis of soil structure also independently of the presence of cementing substances. Fig. 1 shows the quantification of the influence of wetting and drying cycles on soil porosity in different types of soils. In the sandy loam soil the total porosity showed a slight increase during the wetting and drying cycles. On the contrary, in the clay loam soil and in the silty clay soil the total porosity strongly increased until the 11th wetting and drying cycle, and at the end of the 16th it showed a slight decrease (Sartori et al., 1985; Pagliai et al., 1987). The micro and macroscopic observations revealed that the large degree of porosity was represented by intra-aggregate pores, which surrounded or separated the newly formed soil aggregates, thus confirming that wetting and drying cycles produce aggregation and this capacity is strongly related to the content of clay in soil. The slight decrease of porosity after the 11th cycle may be ascribed to a decrease of the microbial activity in soil during the continuous repetition of wetting and drying cycles. Utomo and Dexter (1982) demonstrated the importance of microbial activity in the formation of soil aggregates during wetting and drying cycles, and the formation of soil aggregates is related to soil porosity. However, the genesis of soil structure mainly depends on the presence of cementing substances. Organic matter is the main cementing substance and the main processes of soil aggregation are of a biological origin. In soils with an appreciable content of clay, the primary particles tend, under favourable conditions, to group themselves into structural units known as secondary particles or aggregates. Such aggregates are not characterized by any fixed size or stability. The visible aggregates, which are generally of the order of several millimetres to several centimetres in diameter, are often called peds or macroaggregates. The literature concerning the complex interrelationship of physical, biological and chemical reactions involved in the formation of soil aggregates is very wide; for example, the review of Harris et al. (1966) listed nearly four hundred references. Sequi (1978) summarized the main mechanisms involved in soil aggregation as the linkage between organic matter and mineral

constituents, the adhesion of living organisms to mineral constituents, and physical actions of living organisms on inorganic constituents.

Linkage between organic matter and mineral constituents. - Organic matter is responsible for the formation of real bonds between organic and inorganic soil components and this is perhaps the most widely studied topic in soil aggregation (Greenland, 1965a,b; Oades, 1984). Authors often deal with organic matter without considering the presence of living organisms but this approach appears incomplete, except for peaty soils, where organic matter represents a very unusual pedological entity. Peat can be considered as a complex of mummified materials, like a transmission stage towards the formation of lignite and fossilization. However, living organisms represent a consistent percentage of soil organic carbon. Jenkinson and Powlson (1976) in their study on soil biomass determination showed that in several arable and uncultivated soils, containing from 0.84 to 3.49% of organic matter, the living organisms constituted from about a thirtieth to one fiftieth of the organic matter. Plant roots, in a broad sense, must be added to the soil biomass. Roots exert pressure which compresses aggregates. Water uptake by roots causes differential dehydration, shrinkage, and the opening of numerous small cracks. Moreover, root exudations and the continual death of roots and particularly of root hairs promote microbial activity which results in the production of humic cements. Since these binding substances are transitory, as they are susceptible to further microbial decomposition, organic matter must be replenished and supplied continually if aggregate stability is to be maintained in the long run. However, in some cases roots and rhizosphere organisms produce a range of acids that can stimulate the production of dispersible clay (Oades, 1984). Such an effect has been described by Reid and Goss (1982) and Reid et al. (1982), particularly for maize plants during the early weeks of growth. The acids produced could break organic matter-Fe, Al, Ca-clay bonds by lowering the pH and/or complexing the metal ions. Drying of the soil by plant growth or by air drying leads to an increase in the formation and stability of soil aggregates, presumably due to the increased sorption and effectiveness of organic binding agents, probably due to polysaccharides which are exuded from roots in addition to the soluble exudates (Oades, 1978).

Polysaccharides, among the many microbial products capable of binding soil aggregates, are prominent. Such materials are attached to clay surfaces by means of cation bridges, hydrogen bondings, van der Waals forces, and anion adsorption mechanisms. Polysaccharides, in particular, consist of large, linear, and flexible molecules capable of forming multiple bonds with several particles at once. The action of polysaccharides has been the subject of much research. Carbohydrates represent one quarter of soil organic matter and are essentially derived from plant polysaccharides in roots and plant debris. Other polysaccharides in soils are mucilages associated with roots (Oades, 1978), or microbial mucilages (Hepper, 1975). Cheshire et al. (1983) suggested that polysaccharides are the most important cements present in soils. It is widely demonstrated that the effectiveness of microbial polysaccharides in promoting soil aggregation is strictly correlated with their molecular weight. Pagliai et al. (1980) showed that microbial dextrans with a higher molecular weight appeared most efficient in improving pore size distribution and the stability of soil aggregates. Thus the molecular weight or length of the molecules is considered to be an important factor in improving the pore size distribution of a soil or the water stability of aggregates. The longer the molecules are the more able they are to link particles together in the walls of pores to stabilize the pores against the destructive forces of wetting.

Micromorphological techniques can give useful contributions in the studies dealing with the interaction of organic matter-soil structure by means of the microscopic examination of soil thin sections. Fig. 2 shows accumulation of organic matter distributed as a coat along the walls of elongated pores. These coats on pore walls can effectively seal pores from the adjacent soil matrix, thus stabilizing the pore walls against the destructive forces of water and assuring the functionality of the pores. These favourable conditions, with respect to soil structure, are not permanent. In fact, when the organic matter is totally decomposed and mineralized it loses its capability as a cementing substance, therefore the pore walls collapse and close the pore. This is the first step of soil structural degradation; to avoid this it is necessary to assure an adequate turnover of organic matter in soil. Pagliai and Vittori Antisari (1993) confirmed that the increase of micro and macro porosity and the improvement of soil structure in cultivated soils strictly depend on the addition of organic materials, including waste organic matter.

Because clays and organic materials are polyanions they can be bridged by polyvalent cations, such as Ca, Mg, Al, Fe. Soil organisms accumulate inorganic constituents as an essential part of their environment. Soil alkali-soluble organic matter itself contains large amounts, up to about 40%, of ash composed mainly of iron, aluminium and silica. Most of the iron is firmly bound to organic matter, as it is not removable by ion-exchange resins (Sequi et al., 1975). The main evidence for the role of cation bridges has been based on the increased disaggregation of soil after treatment with complexing agents or after treatments with acids (Hamblin and Greenland, 1977). The complexing agents most commonly used have been pyrophosphate and acetylacetone, which aimed particularly at Al and Fe. It has been shown consistently that pyrophosphate aids the disruption of aggregates beyond that due to saturation of the system with Na. Giovannini and Sequi (1976a, b) showed that treatments with acetylacetone in benzene, a reagent which essentially extracts metals bound to the organic matter, led to a substantial decrease in the water stability of soil aggregates. They suggested that metals can behave as bridges of nets composed of polymeric chains of soil organic matter.

Humification processes generally lead to a concentration of aromatic substances in soil organic matter. Such substances exert two main influences in soil. Firstly they complex, protect, and stabilize organic compounds in general (Haider et al., 1975). Griffiths and Burns (1972) showed that tannic acid stabilizes soil aggregates treated with polysaccharides. This effect can be due to the protection from microbial degradation by phenolic substances; the protective effect is general for most enzymes in soil (Sequi, 1978). The second influence depends on the hydrophobic nature of aromatic substances. The soil may be regarded as a system of hydrophilic components, with organic matter as unique hydrophobic constituent, due to its aromatic substances. Water repellency affects water infiltration, evaporation and water flow (Debano, 1975), and consequently influences soil structure. The commonly reported occurrence of water repellency in soil after fire (Giovannini and Lucchesi, 1987) is an interesting aspect of the same problem.

Adhesion of living organisms to mineral constituents. - The soil microflora can be regarded as aquatic and it is reasonable to regard bacterial cells as part of the colloidal system in soil (Oades, 1984). As is well known, microorganisms are not easily leached or washed out

from soils, because they are attached to either clay or humic substances. The size of cells are on average similar to that of clay particles and these cells are ionogenic and in many cases amphoteric due largely to carboxyl and amino groups. Most microbial cells have a net negative charge at the pH of the soil. These negatively charged bioparticles can attract clay surfaces sufficiently close as to be adsorbed. Marshall (1971) showed the adhesion of microorganisms to clay particles including the fundamental forces of interaction between organisms and surface and the implications for the organisms involved and the surfaces acting as adsorbed.

Physical actions of living organisms on inorganic constituents. - Micro and macro living organisms can also cause mechanical actions on inorganic constituents in several ways. Unicellular organisms, during proliferation surround and link soil particles. Other filamentous growth forms, such as fungi hyphae, explore pores and entangle mineral constituents. Plant roots grow in the most favourable directions, forming a three dimensional net whose thickness is conditioned by physical and chemical fertility. Developing roots also compress the adjacent soil, so reducing the pore size. As the roots and hyphae decompose the fragments become the centre of water stable aggregates, because of mucilages produced during decomposition of the organic fragments interact with clay (Oades, 1984). Therefore, plant roots, such as fungal hyphae and bacterial colonies may well share physical actions with cementing properties of slime layers and exudates.

Organisms which have their own movement contribute to the migration of soil particles. In their movements these organisms abandon their catabolic residues in different places and this products can act as cementing substances.

Water adsorption by plant roots and other organisms causes a strong modification of the physical status of the adjacent soil and, for example, shrinkage phenomena can occur. Another important effect common to all living organisms, is their influence on the soil air composition. Such changes cause variations in oxidation potentials and redistribution by diffusion (Sequi, 1978).

Aggregate stability

As already said soil structure is a dynamic property and the degree of aggregation is a time-variable property, as aggregates form, degrade and reform periodically. A visible example of this cycle can be seen in cultivated soils where the granular or crumbly structure originated by the tillage operation visibly and rapidly deteriorates under destructive forces, such as, for example, rainfall impact (causing slaking, swelling, shrinkage, crusting and erosion) (Fig. 3). Therefore, soils vary in the degree to which they are vulnerable to externally applied natural or man-made destructive forces. Aggregate stability is a measure of such vulnerability and can be defined as the capacity of cohesive forces between soil particles to resist the externally applied destructive forces. The evaluation of aggregate stability and of the size of stable aggregates is important because it is the size of aggregates which determines their susceptibility to water erosion. Moreover, the size of aggregates determines the volume of pore space in soils.

Since the stability of aggregates and pores decreases on wetting dry soil, treatments in water are commonly used to determine the aggregate stability. Such treatments are mostly variations of the wet sieving method introduced by Tiulin (1933) and modified by several others (Kemper and Rosenou, 1986). In general, these methods consist in putting soil aggregates of a selected size onto a 0.2 mm mesh sieve which is moved mechanically in water. The dry weight of soil remaining on the sieve, expressed as a percentage of the original dry weight, is reported as aggregate stability. A deduction is made from both weights for sand grains too large to go through the sieve. Wet sieving is widely used for comparing effects of soil treatments and management practices on aggregate stability. To obtain reproducible results the method has to be rigorously standardized particularly as regards sample preparation, wetting procedure, and the handling and treating of wet aggregates. Slow wetting by capillarity from water under suction is commonly used to allow air to escape freely and so avoid explosive damage to weak aggregates. The aggregate stability is usually much lower when the samples are immersed rapidly.

Aggregate stability strongly depends on the soil organic matter. Its role ranges, as already said, from the production of cementing substances to the action of living organisms, roots, etc. Stability increases with a greater content of organic matter in the soils, especially among those with less than 2 per cent (Kemper and Koch, 1966). Iron and aluminium oxides

can act alone or in combination with organic matter to stabilize aggregates. The clay content also plays a strong influence, since stability arises internally from bonds between clay plates, packets of clay plates (domains), and other particles. However, the presence of clay particles does not in itself ensure stability. Sodium in the exchange complex of the clay can make it quite unstable.

Soil Porosity

Soil porosity represents the liquid and gaseous soil phases and its relation with the solid phase is explained in the chapter on soil physical composition. To characterize the pore system it is necessary, first of all, to determine the size distribution and shape of pores because the agronomic functions of pores depend on their size and shape. According to one of the most widely used classifications, that of Greenland (1977) reported in Table 1, the very fine pores less than $0.005\ \mu\text{m}$, called "bonding pores", are critically important in terms of the forces holding domains and aggregates of primary particles together; pores of less than $0.5\ \mu\text{m}$ are the "residual pores" for the chemical interactions at the molecular level; pores which have an equivalent pore diameter ranging from 0.5 to $50\ \mu\text{m}$ are the "storage pores", i.e. the pores that store water for plants and for micro-organisms; and the pores ranging from 50 to $500\ \mu\text{m}$ are those called "transmission pores" in which the movements of water are important for plants, and, moreover, they are the pores needed by feeding roots to grow into. The water content when pores larger than $50\ \mu\text{m}$ have drained, corresponds to the field capacity of the soil. The wilting point commences when most pores larger than approximately $0.5\ \mu\text{m}$ have emptied.

Pores larger than $500\ \mu\text{m}$ can have some useful effects on root penetration and water movement (drainage), especially in fine-textured soils. However, a high percentage of this latter type of pore (above 70-80% of the total porosity) in soils is usually an index of poor soil structure, especially in relation to plant growth. This is because surface cracks, which develop after rainfall, when the stability of soil aggregates is poor, belong to this size classe (Pagliai et al., 1981, 1983). Until now the necessary proportion of large pores for air and water transmission and easy root growth has generally been inadequately defined. In fact, adequate

storage pores (0.5-50 μm) as well as adequate transmission pores (50-500 μm) are necessary for plant growth (Greenland, 1981).

A traditional method to determine pore size distribution is based on moisture retention data, using the concept of "equivalent" sizes derived from a capillary model. This model relates the height of capillary rise (h) to pore radius:

$$\rho gh = 2\gamma \cos\theta / r$$

where ρgh is the suction (ρ is the density of water = 0.9982 Mg m^{-3} at 20°C , g is the acceleration due to gravity = 9.80 m s^{-2} , and h is the height of capillary rise expressed in metres), γ the surface tension of water (72.75 mNm^{-2} at 20°C), θ the solid-liquid contact angle which is here assumed to be zero, and r the radius of the capillary tube (m). This relationship can be pictured, for example, as a continuous graph, relating capillary diameter ($2r$) to corresponding negative pressure (suction) (Fig. 4). This figure illustrates that fine pores can exercise a larger pull than large pores. For example, a cylindrical pore diameter of $30 \mu\text{m}$ corresponds to a capillary rise of 1 m, (thus, suction of 1 m will withdraw water from pores with an effective diameter greater than $30 \mu\text{m}$) and a diameter of $100 \mu\text{m}$ with a relatively low capillary rise of 0.28 m. On this basis, a moisture characteristic curve can be used to show the amount of pore space (as given by the water content on a volume basis) that has pores smaller than a given effective size.

However, for fine-textured soils the calculation of pore size distribution in this way is likely to be invalid, for two reasons. First, it is unlikely that the calculated pore size distribution represents the pore size distribution of the soil at all moisture contents. Secondly, during water desorption most fine-textured soils shrink and particle rearrangement takes place (Aylmore, 1961).

Pore size distribution can most rapidly be determined over a wide range ($0.007 \mu\text{m}$ to $100 \mu\text{m}$) by mercury intrusion porosimetry, i.e. measuring the pressure required to force mercury into the pores. Such a pressure depends on the contact angle, the pore shape and the surface tension of the liquid. For cylindrically shaped pores the relationship between pressure and the minimum pore diameter which may be intruded was given by Washburn (1921) as:

$$P = -4\gamma \cos\theta / d$$

where P is the required pressure, γ the surface tension of the liquid, θ the contact angle and d the diameter of the pores.

Experimentally, the method consists of the following steps: one applies an increment of pressure to a suitable apparatus (mercury intrusion porosimeter) in which mercury is permitted to intrude into the pores of a sample; the volume intruded is measured when the liquid ceases to flow; and then the process is repeated under successively higher pressures to the limit of the apparatus. The volume of mercury intruded in a given step is taken as the total volume of pores with equivalent diameters smaller than that corresponding to the pressure causing the flow. By adding the results of all of the steps a cumulative volume distribution with respect to pore diameters is built up.

One disadvantage of this method is that the pores must be empty of water at the start of the measurement. Diamond (1970) and Lawrence (1977) summarised the existing method of drying the soil for this kind of analysis.

Using the technique of image analysis it is now possible to characterize soil structure by the quantification of soil porosity in all its aspects (pore shape, pore size distribution, irregularity, orientation, continuity, etc.) on thin sections, prepared from undisturbed soil samples (Bouma et al., 1977, 1982; Murphy et al., 1977a, b; Pagliai et al., 1983, 1984; Pagliai, 1988). This morphometric technique has the advantage that the measurement and the characterization of pore space can be combined with a visual appreciation of the type and distribution of pores in soil in a particular moment of its dynamic evolution. For this analysis it is necessary to prepare thin sections of soil following a procedure which consists in taking undisturbed soil samples using appropriate implements, containers and techniques taking care that the interior structure of the soil samples remains undisturbed. Then the soil samples, carefully packed, are transported to the laboratory, dried to avoid pronounced shrinkage phenomena, using appropriate methods, e.g. acetone replacement of the water (Murphy et al., 1986), and impregnated, under vacuum, with a polyester resin, which has the characteristic of polymerising slowly at room temperature without altering in any way the structure of the soil. Practically, this resin fills the pores of the soil. When the soil samples are hardened (generally after 4-6 weeks) they are made into vertically or horizontally thin sections by using

appropriate machines (Murphy, 1986). Their thickness is about 30 μm so that they can be analysed by the microscope in transmitted light. The size depends on the kind of machines available; for porosity measurement a size larger than 6X6 cm should be recommended. The image analysis can be used not only on soil thin sections but also on polished faces of large soil blocks impregnated directly in the field with (fairly cheap) materials such as paraffin wax (Dexter, 1988), or plaster of Paris (FitzPatrick et al., 1985), or resin (Moran et al., 1989).

The soil thin sections or photographs (Pagliai et al., 1984) are analysed with image analyzers, such as the Quantimet (Murphy, 1977a, b; Pagliai et al., 1983, 1984). Two-dimensional images obtained have to be transformed into data representing three-dimensional area percentages that are representative for three-dimensional volumes. Stereology techniques have been recently applied to achieve this objective (Ringrose-Voase and Bullock, 1984; Ringrose-Voase and Nortclif, 1977). Basic measurements of image analysis on pores include number, area, perimeter, diameters, projections, etc., and these are supplemented by derived measurements such as shape factors, size distribution, continuity, irregularity and orientation.

The shape factors allow division of pores into different shape groups such as, for example, more or less rounded (regular), irregular and elongated pores (Bouma et al., 1977; Pagliai et al., 1983). Pores of each shape group can be further subdivided into a select number of size classes according to either the equivalent pore diameter for rounded and irregular pores or the width for elongated pores. The equivalent pore diameters are calculated from the area of regular and irregular pores, while the width of elongated pores is calculated from their area and perimeter data using a quadratic equation because it is assumed that elongated pores are long narrow rectangles (Pagliai et al., 1984).

The regular pores are obviously those of a rounded shape and can be distinguished in two types according to their origin: the spherical pores formed by entrapped air during soil drying and the channels and chambers formed by biological activity (root growth and movement of soil fauna). Their distinction on soil thin sections is very evident because spherical pores (vesicles, according to Brewer, 1964) have very smooth walls, while channels, even though cut in a transversal way on thin section, present rough walls with deposits of

insect excrements or root exudates. The presence of many spherical pores of the first type (vesicles) creates a vesicular structure typical of soils with evident problems of degradation.

The irregular pores are the common soil voids with irregular walls (vughs, according to the micromorphological terminology of Brewer, 1964) and can be isolated (packing voids) or interconnected. The dominant presence of these pores produce the typical vughy structure (Bullock et al., 1985). In cultivated soils these pores can be originated by the effect of soil tillage implements.

The elongated pores can be distinguished in two types, i.e., cracks and thin fissures (planes). The former are typical of clay soils with a depleted soil organic matter content and they are visible at the surface when the soil is dry and has shrunk. The thin fissures are the most important, especially from an agronomic point of view, in fact, they are the typical transmission pores already mentioned. An adequate proportion of this type of pore (over 10% of the total porosity) generally creates an angular to subangular blocky structure of good quality. Obviously for this to be true it is necessary for these pores to be homogeneously distributed in the soil matrix. In fact, for these pores characterization by image analysis, besides the identification of their shape and width, must also determine their length. With the same procedure of width determination it is also possible to determine the length of these elongated pores, which may reflect their continuity, and it is well known that the flow of water through soil depends on the continuity of large pores. Therefore the analysis of pore patterns allows the characterization and prediction of flow processes in soils.

For root growth and water movement not only the size and continuity of elongated pores are important but also their irregularity and orientation. The ratio convex perimeter/perimeter or convex area/area of elongated pores gives information about their irregularity, tortuosity and re-entrancy. As regards water movement, for example, the very regular and the moderately regular elongated pores play a different role. The very regular elongated pores are flat and smooth pores with accommodating faces which tend to seal when the soil is wet and, thus prevent water movement. In contrast, the moderately regular elongated pores have walls which do not accommodate each other. Therefore, these pores permit water movement even when the soil is wet and fully swollen (Pagliai et al., 1984). The

ratio vertical/horizontal dimensions gives the orientation of elongated pores (Pagliai et al., 1984). It is easily understandable that many soil processes such as water movement, leaching, clay migration, etc., are strongly related to the orientation of pores in soil and these processes radically change depending on whether a vertical or horizontal pore orientation is dominant.

Types of soil structure

In general, three broad categories of soil structure can be recognized: single grained, aggregates and massive. The definition of the several types of soil structure is fundamental for a first qualitative evaluation of soil physical fertility. According to the definition of soil structure the different types of structure are originated by the combination and the spatial arrangement of different types of pores and aggregates.

The main types of soil structure can be summarized as follows:

Single grain structure. - Typical of sandy soils; the quartz (sand-sized) grains are completely loose and the fine materials in the intergranular spaces are very rare. The porosity is represented by packing voids delimited by quartz grains. When these grains are of a more or less uniform shape and size a rather compact grain structure can be originated.

Bridged/pellicular grain structure. - The sand-sized grains are bridged or coated by fine materials, usually clay, which can cement some grains to each other, so originating microaggregates.

Vughy structure. - There are no separated aggregates and the mass is broken up by scattered but not interconnected irregular pores, "vughs", and occasional channels and chambers (Bullock et al., 1985) (Fig. 5). Such a structure allows good soil aeration but the continuity of pores is limited. In the case of heavy rains ponding water can be present in the soil profile with consequent collapse of the soil structure. Such a type of structure is, therefore, unstable and transitory.

Vesicular structure. - There are no separated aggregates and the mass is broken up by rounded pores, "vesicles" (Bullock et al., 1985), originated by entrapped air during drying processes (Fig. 6). This kind of structure is an indicator of degraded soils.

Crumb structure. - The soil aggregates are more or less rounded, often rugose, well separated from each other and rather compact inside. The porosity is represented by pore space (packing voids) which separates the aggregates. This type of structure is often originated by anthropogenic activities (soil tillage).

Subangular blocky structure. - The soil aggregates are separated by elongated continuous pores (planes), are of different sizes and can be rather porous inside. Aggregate faces largely accommodate each other. From an agronomic point of view, this is the best type of soil structure because the continuity of elongated pores allows good water movement and facilitates root growth. Moreover, it is a rather stable soil structure.

Angular blocky structure. - The soil aggregates have angular edges and are separated by elongated pores of a regular shape. Aggregate faces normally accommodate each other. This is a typical structure of clay soils and it less stable than the subangular blocky structure because during the wetting process the soil swells and pores, which have accommodating walls, tend to seal up, thus, prevent water movement, resulting in soil degradation.

Platy structure. - The thin and flat soil aggregates are separated by elongated pores oriented parallel to the soil surface and, therefore, not continuous in a vertical sense. This leads, as a consequence, to a drastic reduction of water infiltration capacity. Soils with this type of structure are subject to water stagnation or runoff and erosion depending on their slope. The platy structure is typical of compact soils.

Prismatic structure. - The soil macroaggregates are divided into prisms separated by vertically oriented elongated pores with accommodating walls. Also this type of structure is typical of clay soils, especially in the B horizon, and is not very stable because the swelling of the soil when wet causes the closing of pores.

Massive structure. - The soil material is very compact, there are no visible separated aggregates. The porosity is very low and represented by small pores isolated in the soil matrix. This type of structure represents a bad "habitat" for plant development and is common in degraded soil with a low content of organic matter.

Complex structure. - The soil presents two or more types of soil structure.

Effects of agricultural management practices and treatments on soil structure

Many anthropogenic activities affect soil structure. To evaluate the impact of management practices on the soil environment it is necessary to quantify the modifications of soil structure. Parameters such as pore size distribution, pore shape, pore continuity and relative position of aggregates and pores are very important for evaluating induced modifications of soil structure, e.g., by different management practices such as soil tillage, application of manures, irrigation and soil compaction by wheel traffic.

Soil tillage. - Long-term intensive arable cultivation has negative effects on soil physical properties, particularly on soil structure, with resulting effects on soil erodibility and crop yields. The need to check the degradation of soil structure has caused farmers to consider no-till management as an alternative to conventional tillage. Abandoning traditional farming rotations and adopting intensive monocultures, without applications of farmyard manure or organic materials to the soil, has decreased the soil organic matter content with evident degradation of soil structure. The resulting soil porosity conditions are often unfavorable for crop growth (Pagliai et al., 1983; 1984; 1989; Shipitalo and Protz, 1987).

No-till management has been shown to increase the soil faunal populations (Barnes and Ellis, 1979) and this could ameliorate the detrimental effect of row crop production on soil structure. However, no-till management can not always be recommended because of reduced yields in some types of soil, probably due to the development of soil physical properties not conducive to plant growth (Pagliai and Pezzarossa, 1990). Therefore, the relationships between tillage practices, soil structure, and soil fauna are considered worthy of further investigation with the aim of preserving the soil environment.

Recent research has shown that management practices predominately affect soil porosity (Kooistra et al., 1990; Pagliai et al., 1989; Shipitalo and Protz, 1987). Management practices can also affect biological activity and can therefore affect both the formation and preservation of biopores, important for water movement and root development (Kooistra, 1991; Pagliai and De Nobile, 1993).

Tillage produces a loose and crumb structure with increased macroporosity which is dominated by large packing pores separating aggregates or clods. Such a crumb structure

produced by tillage is very unstable, especially in the surface layer, and after the first rainfall and wetting and drying cycles can be transformed into a more compact structure, often associated with the presence of a surface crust. This transformation is strictly related to the type of soil, the organic matter content and the aggregate stability. During the cultural cycle the soil structure changes. Therefore, in order to study the modifications induced by different tillage treatments it is necessary to plan an appropriate sampling strategy during the crop cycle. A sampling in the period of crop ripening is particular important because at that time good soil conditions are critical for crop development (Pagliai et al., 1989). In the case of studying the effect of different tillage treatments on crusting soils a sampling before seedling emergence is very important.

Applications of manures. - The application of organic materials to the soil has also been shown to enhance both soil porosity and pore size distribution. Fig. 7 summarizes results obtained in a long-term field experiment on a sandy loam soil in which annual application of sewage sludges and composts from urban refuse were compared to the applications of farmyard manure or chemical fertilizers (Pagliai et al., 1981; 1983a). The application rates were calculated on the organic carbon basis and were equivalent to 50 metric tons/ha of manure. The soil was planted to corn and the tillage was the same for all treatments, i.e., conventional ploughing to a depth of 25 cm. Data refer to the annual sampling in the period of corn ripening. Soil pore space significantly increased in all years, after treatment with all organic materials compared to treatment with chemical fertilization alone, irrespective of the sampling year.

Pore shape and size distribution were also affected by the application of organic materials. The proportion of storage (30-50 μm) and transmission pores (elongated pores, 50-500 μm) was greater in soil treated with any type of organic material than in soil treated with chemical fertilizers only (Fig. 8). Similar results were obtained with the application of pig slurry to a silty clay soil. In this experiment, a positive relationship between the rate of application and the increase of soil porosity was found. It also emerged that the time of application was important, with the best results being obtained for spring applications (Pagliai et al., 1983a; 1985). The increase of pore space and the modification of the pore system in soil

treated with these organic materials was associated with an increase of aggregate stability (Pagliai et al., 1981). Thus, from these cited data there is clear evidence that sludges, composts and livestock effluents can improve the physical properties of soil in a similar way to manure.

Irrigation. - Intensive continuous cultivation is not the only cause of soil structure degradation; other management practices like irrigation are also important, especially in the longer term. Mathieu (1982) showed that after 15 years of impounding irrigation in clay soils, the structural conditions appeared greatly changed: massive structure, modification of the pore shape and pore size distribution, increased migration of clay particles from the ploughed horizon Ap to the B horizon. Pezzarossa et al. (1991) found that in a peach orchard near Verona (Italy) soil porosity of the surface layer decreased during the irrigation season and that the decrease was significantly greater when irrigation was by impounding rather than under sprinkler-irrigation, due to a reduction in elongated pores (Fig. 9). The latter was associated with a lower water content in the surface soil and reduced root density. The larger amount of water applied in this system caused progressive soil compaction, resulting in decreased porosity and structural degradation.

Fig. 10 visually explains the results of Fig. 9: in fact, in the sprinkler-irrigated soil a subangular blocky structure can be observed, while in the soil irrigated by impounding the structure was more compact, rather massive, with drastic reduction of both the total amount and the size and continuity of elongated pores. This caused decreased water infiltration and, under these conditions, 30% of applied water was lost by run-off. It is clearly intuitive that run-off along the interrow transported nutrients, particularly nitrogen and potassium. For example, results reported by Pagliai (1992) showed that the losses of potassium amounted to 25% of the quantity applied as fertilizer (150 kg/ha/year of K_2O).

Sprinkler irrigation consumes only half the amount of water needed for impounding and causes no evident damage to soil structure, so the loss of water by run-off is negligible but, unfortunately, this method is more expensive and complicated for the farmer. However, for the real conservation of environmental resources and over all to reduce risks of soil structure degradation, erosion and pollution, it is necessary to consider and rationalize all the management practices.

Soil compaction by wheel traffic. - Soil compaction is caused by a combination of natural and man-made forces. The latter are mainly related to vehicle wheel traffic and passage of tillage implements and generally have a much greater compactive effect than natural forces such as raindrop impact, soil swelling and shrinking and root enlargement especially because trends in agricultural engineering over the last few decades have resulted in machines of a greater size and weight.

Fig. 11 shows the results of an experiment dealing with the modifications of soil porosity and structure induced by tractor wheels in a clay sandy loam soil and for how long such modifications are apparent (Pagliai et al., 1992). In the compacted topsoil, porosity greatly decreased (3-4 times), particularly that of elongated pores. At the same time, the size of these pores was reduced and other observations indicated a modification of their orientation. This effect was even more evident in the case of several passes over the same area. Microscopic observation revealed that these elongated pores were thin fissures parallel to the soil surface, forming a platy structure, and had no vertical continuity (Fig. 12). As a consequence water infiltration was reduced, causing an increase of the risk of soil erosion. After seven months the modifications induced by compaction were still evident with respect to the uncompacted topsoil. Bullock et al. (1985b) also found that soil structure could regenerate after compaction in clay soils, although over a period of some years. Bresson and Zambaux (1990) concluded that the noted decrease in the porosity of biopores and fissures due to compaction, associated with increased pH and decreased organic matter due to continuous cultivation, may alter the cohesion of the isotropic ultramicrofabric and make compaction irreversible. Coulon and Bruand (1989) confirmed that compaction modified the pore space geometry and fabric of the elementary particles.

The decrease of soil porosity causes, as already stated, strong structural modifications in the compacted areas, and it is strongly related to the cone penetrometer resistance. Fig. 13 shows a good correlation, in the surface compacted layer, between soil porosity and resistance to cone penetration.

The formation of compacted layers along the profile (ploughpans) - Continually ploughing the soil may cause the formation of a ploughpan (ploughsole) at the lower depth

limit of cultivation (Fig. 14). Fig. 15 shows that the macroporosity, measured by image analysis, was high just above the ploughpan; decreased to a very low value in the ploughpan, and slightly increased in uncultivated soil below. In this case the ploughpan is only a few centimetres deep but other studies have shown that the ploughpan can reach a thickness of up to 10 cm. Fig. 15 also shows that other types of tillage such as the chiesel ploughing or minimum tillage could prevent or reduce the formation of the ploughpan. Microscopic observations revealed that in the first few centimetres of the ploughpan there were a few, very thin, elongated poroids parallel to the surface of the ploughpan. Some irregular pores were present in the lower part of the ploughpan. The presence of such dense layers at the lower depth limit of cultivation may greatly reduce water drainage and hamper root development at depth. Kooistra et al. (1984) observed that the continuity of large pores, made visible by stained channels, in thin sections of an undisturbed ploughpan in a sandy loam soil was greater than that in a ploughpan disturbed by deep rototilling, which had discontinuous (unstained) packing pores between the aggregates. This means that once the ploughpan is formed it is very difficult to recover a structure which allows good drainage.

REFERENCES

- Aylmore, L.A.G. 1961. Hydraulic and swelling of clay mineral systems. Ph. D. Thesis, Univ. of Adelaide.
- Barnes, B.T. and Ellis, F.B. 1979. Effect of different methods of cultivation and direct drilling, and disposal of straw residues, on populations of earthworms. *Journal of Soil Science*, 30: 669-679.
- Bouma, J., Jongerius, A., Boersma, O.H., Jager, A. and Schoonderbeek, D. 1977. The function of different types of macropores during saturated flow through four swelling soil horizons. *Soil Science Society of America Journal*, 41: 945-950.

- Bouma, J., Belmans, C.F.M. and Dekker, L.W. 1982. Water infiltration and redistribution in a silt loam subsoil with vertical worm channels. *Soil Science Society of America Journal*, 46: 917-921.
- Bresson, L.M. and Zambaux, C. 1990. Micromorphological study of compaction induced by mechanical stress for a Dystrichreptic Fragiudalf. In: L.A. Douglas (Editor), *Developments in Soil Science 19. Soil Micromorphology: A Basic and Applied Science*. Elsevier, Amsterdam, 1990. pp. 33-40.
- Brewer, R. 1964. *Fabric and Mineral Analysis of Soils*. John Wiley, New York, 470 pp.
- Bullock, P., Fedoroff, N., Jongerius, A., Stoops, G. and Tursina, T. 1985a. Handbook for soil thin section description. *Waine Res. Pub., Wolverhampton*.
- Bullock, P., Newman, A.C.D. and Thomasson, A.J. 1985b. Porosity aspects of the regeneration of soil structure after compaction. *Soil and Tillage Research*, 5: 325-341.
- Cheshire, M.V., Sparking, G.P. and Mundie, C.M. 1983. Effect of periodate treatment of soil on carbohydrate constituents and soil aggregation. *Journal of Soil Science*, 34: 105-112.
- Coulon, E. and Bruand, A. 1989. Effects of compaction on the pore space geometry in sandy soils. *Soil and Tillage Research*, 15: 137-152.
- Debano, L.F. 1975. Infiltration, evaporation and water movement as related to water repellency. In: W.C. Moldenhauer (ed.), *Soil Conditioners*. Soil Sci. Soc. Am. Spec. Pub. n. 7, Madison, Wisconsin, pp. 155-164.
- Dexter, A.R. 1988. Advances in characterization of soil structure. *Soil and Tillage Research*, 11: 199-238.
- Diamond, S. 1970. Pore size distribution in clays. *Clays and Clay Minerals*, 18: 7-23.
- Fitz Patrick, E.A., Makie, L.A. and Mullins, C.E. 1985. The use of plaster of Paris in the study of soil structure. *Soil Use Management*, 1: 70-72.
- Giovannini, G., Lucchesi, S. and Giachetti, M. 1987. The natural evolution of burned soil: a three-year investigation. *Soil Science*, 143: 220-226.
- Giovannini, G. and Sequi, P. 1976a. Iron and aluminium as cementing substances of soil aggregates. I. Acetylacetone in benzene as an extractant of fractions of soil iron and aluminium. *Journal of Soil Science*, 27: 140-147.

- Giovannini, G. and Sequi, P. 1976b. Iron and aluminium as cementing substances of soil aggregates. II. Changes in stability of soil aggregates following extraction of iron and aluminium by acetylacetone in a non-polar solvent. *Journal of Soil Science*, 27: 148-153.
- Greenland, D.J. 1965a. Interactions between clays and organic compounds in soil Part I. Mechanisms of interaction between clays and defined organic compounds. *Soil Fertilizer*, 28: 415.
- Greenland, D.J. 1965b. Interactions between clays and organic compounds in soil Part II. Adsorption of soil organic compounds and its effect on soil properties. *Soil Fertilizer*, 28: 521.
- Greenland, D.J. 1977. Soil damage by intensive arable cultivation: temporary or permanent? *Philosophical Transactions of the Royal Society of London*, 281: 193-208.
- Greenland, D.J. 1981. Soil management and soil degradation. *Journal of Soil Science*, 32: 301-322.
- Griffiths, E. and Burnes, R.G. 1972. Interaction between phenolic substances and microbial polysaccharides in soil aggregation. *Plant & Soil*, 36: 599-612.
- Haider, K., Martin, J.P. and Filip, Z. 1975. Humus biochemistry. In: E. Paul and A.D. McLaren (eds), *Soil Biochemistry*, vol. 4, pp. 195-244. Marcel Dekker, New York.
- Hamblin, A.P. and Greenland, D.G. 1977. Effect of organic constituents and complex metal ions on aggregate stability of some East Anglian soils. *Journal of Soil Science*, 28: 410-416.
- Harris, R.F., Chester, G. and Allen, O.N. 1966. Dynamic of soil aggregation. *Advances in Agronomy*, 18: 107.
- Hepper, C.M. 19875. Extracellular polysaccharides of soil bacteria. In: N. Walker (ed.), *Soil Microbiology*, pp. 93-100. Butterworths, London.
- Jenkinson, D.S. and Powlson, D.S. 1976. The effect of biocidal treatments on metabolism in soil. V. A method for measuring soil biomass. *Soil Biology Biochemistry*, 8: 209-213.
- Kemper, W.D. and Rosenou, R.C. 1986. aggregate stability and size distribution. *Methods of Soil Analysis, Part I. Physical and Mineralogical Methods. Agronomy Monograph No. 9* (second edition). pp. 425-442.

- Kemper, W.D. and Koch, E.J. 1966. Aggregate stability of soils from western portions of the United States and Canada. U.S. Dep. Agric. Tech. Bull. 1355.
- Kooistra, M.J. 1991. Micromorphological approach to the interactions between soil structure and soil biota. *Agriculture, ecosystems and environment*, 34: 315-328.
- Kooistra, M.J., Bouma, J., Boersma, O.H. and Jager, A. 1984. Physical and micromorphological characterization of undisturbed and disturbed ploughpans in a sandy loam soil. *Soil and Tillage Research*, 4: 405-417.
- Kooistra, M.J., Juo, A.S.R. and Schoonderbeek, D. 1990. Soil degradation in cultivated alfisols under different management systems in south western Nigeria. In: L.A. Douglas (ed.), *Soil Micromorphology*. Elsevier, Amsterdam, pp. 123-148
- Lawrance, G.P. 1977. Measurement of pore size in fine textured soil: a review of existing techniques. *Journal of Soil Science*, 28: 527-540.
- Marshall, K.C. 1971. Sorptive interactions between soil particles and microorganisms. In: A.D. McLaren and J. Skujins (eds), *Soil Biochemistry*, vol. 2. Marcel Dekker, New York.
- Mathieu, C., 1982. Effects of irrigation on the structure of heavy clay soils in north east Morocco. *Soil and Tillage Research*, 2: 311-329.
- Moran, C.J., McBratney, A.B. and Koppi, A.J., 1989. A rapid method for analysis of soil macropore structure. I. Specimen preparation and digital binary image production. *Soil Sci. Soc. Am. J.*, 53: 921-928.
- Murphy, C.P. 1986. Thin section preparation of soils and sediments. A B Academic Publishers, Herts. pp. 149.
- Murphy, C.P., Bullock, P. and Turner, R.H. 1977a. The measurement and characterization of voids in soil thin sections by image analysis: Part I. Principles and techniques. *Journal of Soil Science*, 28: 498-508.
- Murphy, C.P., Bullock, P. and Biswell, K.J. 1977b. The measurement and characterization of voids in soil thin sections by image analysis: Part II. Applications. *Journal of Soil Science*, 28: 509-518.

- Murphy, C.P., McKeague, J.A., Bresson, L.M., Bullock, P., Kooistra, M.J., Miedema, R. and Stoops, G. 1985. Description of soil thin sections: an international comparison. *Geoderma*, 35: 15-37.
- Oades, J.M. 1978. Mucilages at the root surface. *Journal of Soil Science*, 29: 1-16.
- Oades, J.M. 1984. Soil organic matter and structural stability: mechanisms and implications for management. *Plant and Soil*, 76: 319-337.
- Pagliai, M. 1988. Soil porosity aspects. *International Agrophysics*, 4: 215-232.
- Pagliai, M. 1991. Influenza della fertilizzazione organica sulle proprietà fisiche del terreno. In: N. Senesi and T.M. Miano (Editors), *Riciclo di Biomasse di Rifiuto e di Scarto e Fertilizzazione Organica del Suolo*. Patron Editore, Bologna, pp.139-143.
- Pagliai, M. 1992. Losses of Potassium through run-off erosion and "slash and burn" agriculture. *Proceedings 23rd IPI Colloquium*. Praga, Czechoslovakia 12-16 October 1992. pp. 345-362.
- Pagliai, M., Bisdom, E.B.A. and Ledin, S. 1983a. Changes in surface structure (crusting) after application of sewage sludges and pig slurry to cultivated agricultural soils in northern Italy. *Geoderma*, 30: 35-53.
- Pagliai, M. and De Nobili, M. 1993. Relationships between soil porosity, root development and soil enzyme activity in cultivated soils. *Geoderma*, 56: 243-256.
- Pagliai, M., Febo, P., La Marca, M. and Lucamante G. 1992. Effetti del compattamento provocato da differenti tipi di pneumatici su porosità e struttura del terreno. *Rivista di Ingegneria Agraria*, 3: 168- 176.
- Pagliai, M., Guidi, G. and La Marca, M. 1980. Macro and micromorphometric investigation on soil dextran interactions. *Journal of Soil Science*, 31: 493-504.
- Pagliai, M., Guidi, G., La Marca, M., Giachetti, M. and Lucamante, G. 1981. Effect of sewage sludges and composts on soil porosity and aggregation. *Journal Environmental Quality*, 10: 556-561.
- Pagliai, M., La Marca, M. and Lucamante, G. 1983b. Micromorphometric and micromorphological investigations of a clay loam soil in viticulture under zero and conventional tillage. *Journal of Soil Science*, 34: 391-403.

- Pagliai, M., La Marca, M. and Lucamante, G. 1985. Relationship between soil structure and time of landspreading of pig slurry. In: J.H. Williams, G. Guidi and P. L'Hermite (Editors), Long-term Effects of Sewage Sludge and Farm Slurries Applications. Elsevier Applied Science Publishers, London, pp. 45-56.
- Pagliai, M., La Marca, M. and Lucamante, G. 1987. Changes in soil porosity in remoulded soils treated with poultry manure. *Soil Science*, 144: 128-140.
- Pagliai, M., La Marca, M., Lucamante, G. and Genovese, L. 1984. Effects of zero and conventional tillage on the length and irregularity of elongated pores in a clay loam soil under viticulture. *Soil and Tillage Research*, 4: 433-444.
- Pagliai, M. and Pezzarossa, B. 1990. Structure and porosity of silty clay and clay soils in relation to different management practices. *Agricoltura Mediterranea*, 120: 110-116.
- Pagliai, M., Pezzarossa, B., Mazzoncini, M. and Bonari, E. 1989. Effects of tillage on porosity and microstructure of a loam soil. *Soil Technology*, 2: 345-358.
- Pagliai, M. and Vittori Antisari, L., 1993. Influence of waste organic matter on soil micro and macrostructure. *Bioresource Technology*, 43: 205-213.
- Pezzarossa, B., Pagliai, M. and Xiloyannis C. 1991. Effetti di differenti sistemi di irrigazione su alcune caratteristiche fisiche di un terreno investito a pescheto. *Rivista di Frutticoltura*, 7-8: 71-75.
- Reid, J.B. and Goss, M.J. 1982. Interactions between soil drying due to plant water use and decreases in aggregate stability caused by maize roots. *Journal of Soil Science*, 33: 47-53.
- Reid, J.B., Goss, M.J. and Robertson, P.D. 1982. Relationships between the decreases in soil stability affected by the growth of maize roots and changes in organically bound iron and aluminium. *Journal of Soil Science*, 33: 397-410.
- Ringrose-Voase, A.J. and Bullock, P. 1984. The automatic recognition and measurement of soil pore types by image analysis and computer programs. *Journal of Soil Science*, 35: 673-684.
- Ringrose-Voase, A.J. and Nortcliff, S., 1987. The application of stereology to the estimation of soil structural properties: a preview. In: N. Fedoroff, L.M. Bresson and M.A. Courty

- (Editors), Soil Micromorphology, Proc. VIIth Int. Working Meeting on Soil Micromorphology. Association Francaise pour l'Etude du Sol, Plasir, France, pp. 81-88.
- Ringrose-Voase, A.J. and Nys, C. 1990. One-dimensional image analysis of soil structure. II. Interpretation of parameters with respect to four forest soil profiles. *Journal of Soil Science*, 41: 513-527.
- Sartori, G., Ferrari, G.A. and Pagliai, M. 1985. Changes in soil porosity and surface shrinkage in a remoulded saline clay soil treated with compost. *Soil Science*, 523-530.
- Sequi, P. 1978. Soil structure - An outlook. *Agrochimica*, 22: 403-425.
- Sequi, P., Guidi, G. and Petruzzelli, G. 1975. Influence of metals on solubility of soil organic matter. *Geoderma*, 13: 153.
- Shipitalo, M.J. and Protz, R. 1987. Comparison of morphology and porosity of a soil under conventional and zero tillage. *Canadian Journal of Soil Science*, 67: 445-456.
- Tiulin, A.F. 1933. Considerations on genesis of soil structure and on methods for its determination. *Trans. 1st Com. Int. Soc. Soil Sci., Moscow*, vol. A. pp. 111-132.
- Utomo, W.M. and Dexter, A.R. 1982. Changes in soil aggregate water stability induced by wetting and drying cycles in non-saturated soil. *Journal of Soil Science*, 33: 623-637.
- Washburn, E.W. 1921. Note on a method of determining the distribution of pore size in a porous material. *Proc. nat. Acad. Sci. U.S.A.*, 53: 173-185.

Table 1 - Classification of soil pores according to their size. Modified from Greenland (1977).

Equivalent diameter μm (10^{-6}m)	Water Potential (bar)	Name
<0.005	>-600	Bonding space
0.005 -0.5	-600 /-6	Residual pores
0.5 -50	-6 /-0.06	Storage pores
50 -500	-0.06 /-0.006	Transmission pores
>500	<-0.006	Fissures

CAPTIONS TO FIGURES

Fig. 1 - Influence of wetting and drying cycles on porosity formed by pores larger than 30 μm in three different soils. Modified by Pagliai et al. (1987).

Fig. 2 - Microphotograph of a vertically oriented thin section from an undisturbed sample of a sandy loam soil showing the presence of organic matter (dark) as coats on pore (light) walls. Frame length 3 \times 5 mm.

Fig. 3 - Macrophotographs of vertically oriented thin sections prepared from undisturbed samples from the surface layer (0-10 cm) of a clay loam soil after the tillage (a) and after the first impact of rainfall and the following drying cycles (b). The transformation of a crumb/subangular blocky structure to a compact platy structure, associated with the presence of a surface crust, is evident. The rounded pores in the top layer in photo b are vesicles formed by entrapped air during the crusting processes. Frame length 3 \times 5 cm.

Fig. 4 - Example of relationships between pore size (equivalent diameter) and water potential in soil, according to the capillarity model.

Fig. 5 - Microphotograph of a vertically oriented soil thin section showing an example of vughy structure. The white areas represent the pores. Frame length 3 \times 5 mm.

Fig. 6 - Microphotograph of a vertically oriented soil thin section showing an example of vesicular structure. The white areas represent the pores. Frame length 3 \times 5 mm.

Fig. 7 - Effect of soil application of waste materials, farmyard manure and chemical fertilizers on soil porosity, expressed as a percentage of area occupied by pores per thin sections. Modified from Pagliai (1991).

Fig. 8 - Pore size distribution according to the equivalent pore diameter for regular and irregular pores and to the width for elongated pores in soil treated with farmyard manure in comparison with soil treated with chemical fertilizers.

Fig. 9 - Pore size distribution according to the equivalent pore diameter for regular and irregular pores and to the width for elongated pores, in the inter-row of a peach orchard sandy loam soil under two types of irrigation systems.

Fig. 10 - Macrophotographs of vertically oriented thin sections prepared from undisturbed samples from the surface layer (0-10 cm) of a peach orchard sandy loam under sprinkler

irrigation (a) and irrigated by impounding (b). The transformation from a subangular blocky structure in sprinkler-irrigated soil to a rather massive structure in soil irrigated by impounding is very evident. Frame length 3×5 cm.

Fig. 11 - Porosity and pore shape distribution in a clay loam soil uncompacted (control) and compacted by one pass and five passes of wheel traffic at the time of compaction (A) and seven months later (B). Modified from Pagliai et al. (1992).

Fig. 12 - Macrophotographs of vertically oriented thin sections prepared from undisturbed samples from the surface layer (0-10 cm) of uncompacted (a) and compacted (b) clay loam soil. The transformation from a subangular blocky structure of the uncompacted areas to a massive platy structure of the compacted areas. The interruption of pore continuity in the interface soil-atmosphere is also evident. Frame length 3×5 cm.

Fig. 13 - Correlation between soil porosity and resistance to cone penetration in the surface layer (0-10 cm) compacted by wheel traffic.

Fig. 14 - Macrophotographs of vertically oriented thin sections prepared from undisturbed samples from the 23-30 cm layer of a clay loam soil. The lower limit of cultivation (ploughpan) is visible. Frame length 3×5 cm.

Fig. 15 - Example of variation of soil porosity along the cultural profile, measured by image analysis and expressed as a percentage of area occupied by pores larger than 30 µm per thin section.






Fig. 2




Fig. 36




Fig. 3a




Fig. 4




Fig. 5





Fig 6







Fig. 8





Fig. 9




Fig. 106




Fig. 10a




Fig. 11




Fig. 126




Fig. 120




Fig. 13




Fig. 14




Fig. 15

HANDBOOK OF SOIL CONDITIONERS

SUBSTANCES THAT ENHANCE
THE PHYSICAL PROPERTIES OF SOIL

EDITED BY

ARTHUR WALLACE

*University of California—Los Angeles, and
Wallace Laboratories, El Segundo, California*

RICHARD E. TERRY

*Brigham Young University
Provo, Utah*



MARCEL DEKKER, INC.

NEW YORK • BASEL • HONG KONG

ISBN 0-8247-0117-8

The publisher offers discounts on this book when ordered in bulk quantities. For more information, write to Special Sales/Professional Marketing at the address below.

This book is printed on acid-free paper.

Copyright © 1998 by MARCEL DEKKER, INC. All Rights Reserved.

Neither this book nor any part may be reproduced or transmitted in any form or by any means, electronic or mechanical, including photocopying, microfilming, and recording, or by any information storage and retrieval system, without permission in writing from the publisher.

MARCEL DEKKER, INC.
270 Madison Avenue, New York, New York 10016
<http://www.dekker.com>

Current printing (last digit):
10 9 8 7 6 5 4 3 2 1

PRINTED IN THE UNITED STATES OF AMERICA

4

Use of Manures for Soil Improvement

Marcello Pagliai and Nadia Vignozzi *Ministry of Agriculture, Florence, Italy*

I. INTRODUCTION

Long-term intensive arable cultivation has negative effects on soil physical properties, particularly on soil structure, with resulting effects on soil erodibility and crop yields. The need to check the degradation of soil structure has caused farmers to consider reduced tillage management as an alternative to conventional tillage and the utilization of waste organic materials. Abandoning traditional farming rotations and adopting intensive monocultures, without applications of farmyard manure or organic materials to the soil, has decreased the soil organic matter content with evident degradation of soil structure. The resulting soil porosity conditions are often unfavorable for crop growth (Pagliai et al., 1983b, 1984, 1989; Shipitalo and Protz, 1987).

The gradual decrease in organic matter content in intensively cultivated soils, which may lead to deterioration of the soil physical status and possible erosion, is particularly worrying; especially in areas where climatic conditions cause the rapid decomposition of organic matter.

The decrease of soil organic matter control can therefore be prevented by additions of organic materials, which can also contribute to improved soil productivity. An abundant source of organic materials is animal manures from intensive livestock production. Application of animal manures to cultivated soils has been increasing steadily in the last decades because of the need for lower disposal costs and for the recycling of nutrient elements in the soil crop system. Some details of the main characteristics of some of these materials are reported in Table 1.

Few studies are available on the effects of long-term application of animal manures on soil physical properties that may influence soil productivity and prevent soil degradation. The agronomic utilization of animal manures may in-

Table 1 Main Characteristics of Some Animal Manures

Properties	Pig slurry	Cattle slurry	Poultry manure	Farmyard manure
pH	6.9	7.0	7.1	6.8
Water (%)	96.0	92.0	14.0	79.1
Organic matter ^a	50.0	52.0	34.8	64.1
Total N ^a	7.7	3.4	5.0	2.0
Total P ^a	2.5	1.0	2.2	0.8
Total K ^a	4.9	3.0	3.3	2.4
Ash ^a	22.2	28.0	37.0	25.0

^aExpressed as % dry weight.

Source: Modified from Pagliai et al., 1987; Pagliai and Vittori Antisari, 1993.

duce changes in the physical conditions and chemical composition of the soil, and knowledge of these processes is necessary in order to obtain dependable application systems. To evaluate the impact of management practices on the soil environment it is necessary to quantify the modifications of soil structure. According to Greenland (1981), porosity is the best indicator of soil structure conditions. In fact, adequate "storage pores" (0.5–50 μm equivalent pore diameter (e. p. d.), which allows the storage of water for plants and microorganisms) and adequate "transmission pores" (continuous pores ranging from 50 to 500 μm e.p.d., which allow water movement and easy root growth) are necessary for crops. Until now, not only has the necessary proportion of these pores generally been inadequately defined but also there has been a lack of knowledge of how porosity relates to crop yield due to the difficulty of finding an adequate method that allows the complete characterization of soil porosity. Micromorphometric analysis overcomes some of the problems associated with other methods of porosity analysis, because porosity observed by means of an image analyzer in two-dimensional thin soil sections can be related to three-dimensional pore size by utilizing the stereology principle (Ringrose-Voase, 1991).

II. SOIL PORE SYSTEM

Pore space measurements are increasingly being used to quantify soil structure, because it is the size, shape, and continuity of pores that affect many of the important processes in soils (Lawrence, 1977). Possibilities now exist for improving shape models and for creating more accurate models of pore space in soils by using micromorphology coupled with image analysis of photographs of thin soil sections or of impregnated soil blocks (Murphy et al., 1977a,b; Pagliai et al., 1983a,b; Moran et al., 1989; Moran and McBratney, 1992). This technique has the advantage that the measurement and characterization of the pore space can be combined with a visual appreciation of the type and distribution of the pores.

Although it only analyzes 2-dimensional pictures, this method provides useful information on the complexity of pore patterns in soils, not obtainable using other common methods such as mercury intrusion, water retention, and nitrogen sorption. Nowadays, however, with the improvement of software programs, it is possible to apply a mathematical program of stereology to the image analysis in order to characterize soil porosity in three dimensions (Ringrose-Voase and Bullock, 1984; Ringrose-Voase, 1991). This micromorphometric method based on image analysis can be used not only on thin soil sections but also on polished faces of large soil blocks impregnated directly in the field with fairly cheap materials such as paraffin wax (Dexter, 1988), or plaster of Paris (FitzPatrick et al., 1985).

Parameters such as pore size distribution, pore shape, and relative position of aggregates and pores are very important for evaluating induced modifications of soil structure, e.g., by addition of organic materials (Pagliai et al., 1981).

The improvement of soil physical properties and particularly soil porosity may reflect on soil biological and biochemical activities such as, for example, enzyme activity, which is a fundamental property determining the biological fertility of soils. However, very few have considered their reciprocal correlation (Sequi et al., 1985), though it is of utmost importance, since the poral spectrum of soil plays a determining role in the possible habitats of microbes, as it determines hydric conditions, aeration conditions, trophic conditions, and relation between organisms (predation and competition) (Couteaux et al., 1988). Pagliai and De Nobili (1993) have demonstrated that important biological activity such as that of soil enzymes is positively correlated with the amount of pores ranging from 30 to 200 μm . The quantification not only of the changes of soil porosity but also the relation of such changes with chemical and biochemical properties may give useful data for the evaluation of the efficiency of management practices like soil tillage (Pagliai, 1987b), addition of waste organic materials, etc., in order to maintain soil fertility to prevent the soil resource from degradation phenomena.

For a thorough characterization of soil macropores, the main aspects to be considered are not only the pore shape but also the pore size distribution, especially of elongated continuous pores, because many of these pores directly affect plant growth by easing root penetration and storage and transmission of water and gases. For example, according to Russell (1978) and Tippkötter (1983), feeding roots need pores ranging from 100 to 200 μm to grow into. According to Greenland (1977), pores of equivalent pore diameter ranging from 0.5 to 50 μm are the storage pores, which provide the water reservoir for plants and microorganisms, while transmission pores ranging from 50 to 500 μm (elongated and continuous pores) are important both in soil–water–plant relationships and in maintaining good soil structure conditions. Damage to soil structure can be recognized by decreases in the proportion of transmission pores. Pores larger than 500 μm , called fissures, are useful for root penetration and for drainage especially in fine-textured soils. Adequate proportion of all these types of pores are needed for good soil quality and plant growth.

A. Microporosity

In a long-term field experiment established on a silty clay soil (Vertic Cambisol, according to the FAO world map legend) planted with corn, the application of pig slurry and farmyard manure increased the microporosity (Fig. 1). Pig slurries were surface applied by a liquid manure spreader at the end of May, and the addition rates were 100, 200, and 300 metric tons (MG) ha⁻¹; the farmyard manure (50 MG ha⁻¹) was incorporated into the soil at the ploughing time (winter) and the control soil was given chemical fertilizers. Sampling for porosity measurement was carried out at the end of September after ten years of manure application. The microporosity was measured by the combination of scanning electron microscopy and image analysis (Pagliai et al., 1983b; Pagliai and Vittori Antisari, 1993). All type of pores increased, but while the increase of regular and irregular micropores was approximately the same in all treated soils, the increase of elongated micropores was more relevant. In soil treated with pig slurry such an increase seemed to be related to the amount of slurry added to the soil. The increase of elongated micropores was associated with the new formation of microaggregates, and moreover the increase of pores in this size range was important, since these pores are related to the amount of water available to plants. There are few data in the literature about the effect of manure on microporosity, but similar conclusions were also draw by Schjønning et al. (1994) on a sandy loam soil.




Figure 1 Effect of surface application of pig slurry (PS, 100, 200, and 300 MG ha⁻¹) and incorporation of farmyard manure (FYM) on soil microporosity (0.2–50 μm) on a silty clay soil expressed as a percentage of total area of pores per thin section. Total microporosity values followed by the same letter are not significantly different at the 0.05 level. (Part of these data are modified from Pagliai and Vittori Antisari, 1993.)

B. Macroporosity

In the same long-term field experiment on silty clay soil, mentioned above, total macroporosity (pores larger than 50 μm) increased in the soil treated with pig slurry and farmyard manure with respect to the control soil treated only with chemical fertilizers (Fig. 2). The increase of macroporosity was proportional to the amount of pig slurry added to the soil, while the effect of farmyard manure incorporation was equivalent to the application of 200 MG ha^{-1} of pig slurry. Differences between the control soils and soils treated with the highest amount of slurry (300 MG ha^{-1}) were highly significant (Pagliai and Vittori Antisari, 1993). This could be an important result, because this experimental field was located in an area of high density of pig livestock, and the soil was representative of this area. Therefore the agronomic utilization of such livestock effluents, besides the beneficial effect on the soil, resolves the great problem of the disposal of these wastes.

The effect of liquid manure (livestock effluents) on soil physical properties strongly depends on the time of landspreading. Figure 3 summarizes results obtained in the same field experiment where pig slurries were surface applied in February (instead of May) on the ploughed soil, approximately one month before the corn seeding. In comparison with Fig. 2 it is clear that, at the sampling time in September, i.e., in the period of corn ripening when good soil conditions




Figure 2 Effect of surface application of pig slurry (PS, 100, 200, and 300 MG ha^{-1}) and incorporation of farmyard manure (FYM) on soil macroporosity ($>50 \mu\text{m}$) on a silty clay soil expressed as a percentage of total area of pores per thin section. Total macroporosity values followed by the same letter are not significantly different at the 0.05 level. (Part of these data are modified from Pagliai and Vittori Antisari, 1993.)




Figure 3 Effect of surface application of pig slurry (PS, 100, 200, and 300 MG ha⁻¹), applied in winter, and incorporation of farmyard manure (FYM) on soil macroporosity (>50 μ m) on a silty clay soil expressed as a percentage of total area of pores per thin section. Total macroporosity values followed by the same letter are not significantly different at the 0.05 level. (Part of these data are modified from Pagliai and Vitori Antisari, 1993.)

are critical for crop development, the porosity in soil treated with pig slurry was lower than in soil where slurries were applied at the end of May. The lower addition rates did not show significant difference compared to the control, and the effect of the highest rate was lower than that of the farmyard manure. In May the soil temperature and humidity were optimal for biological activity, which was higher than in February, and the biological activity strongly contributed to improve soil structure through, for example, the formation of biopores. It was demonstrated that the decomposition of organic materials (like sewage sludges, livestock effluents) due to microbial activity was maximum in the first few weeks following their incorporation in soil (Tester et al., 1977; Terry et al., 1979). Pagliai et al. (1985) have also shown that the decomposition of livestock effluents in soil was quite rapid and the residual effect for the following crop was much lower than that of the farmyard manure. For this, the fertilization with animal slurries needs continuous applications of adequate rates of these materials. The amount of the addition rate depends on the type of soil and on its hydrological properties. The limiting factor to the utilization of a large amount of slurry is represented by the nitrogen losses by surface runoff or through the leaching along the soil profile (Brogan, 1981). This may cause a negative environmental impact due to the possible pollution of surface or ground waters.

Other limitations to the agronomic use of waste materials could be represented by the salinity of both soil and products used. Sartori et al. (1985) showed that the application of a compost, supplied by a municipal treatment plant, on a saline clay soil caused a strong decrease of soil porosity and thus leading, as a consequence, to a deterioration of soil physical properties. It is well known that compost and sewage sludges are characterized by relevant saline concentration (this aspect is developed in Chapter 7, 8 and 9 of this book). However, some animal manures may also create problems when applied to saline soil. On the contrary, when these manures are applied to non saline soil, they generally improve the soil physical properties.

Figure 4 summarizes results of laboratory trials to determine the effects of poultry manure applications to a nonsaline clay loam soil on soil porosity following wetting and drying cycles (Pagliai et al., 1987). In control samples the total porosity increased until the 11th wetting and drying cycles, and at the end of the 16th it showed a slight decrease. This means that the studied soil had a good suitability to soil structure regeneration. In poultry-manure treated samples the total porosity was higher than in the control samples. During the wetting and drying cycles it increased until the 11th cycle and then slightly decreased. This slight decrease of porosity after the 11th cycle in both treated




Figure 4 Effects of the addition of poultry manure (PM) on soil porosity, expressed as percentage of total area occupied by pores larger than 50 μm per thin section, following wetting and drying cycles. (Modified from Pagliai et al., 1987.) The poultry manure was applied at two rate levels: low rate 0.75% by weight of soil (both weight of poultry manure and soil were calculated on dry-matter basis) and high rate 7.5%.

and control samples may be ascribed to a decrease of the microbial activity in soil during the continuous repetition of wetting and drying cycles. Utomo and Dexter (1982) demonstrated the importance of microbial activity in the formation of soil aggregates during wetting and drying cycles, and the formation of soil aggregates is related to soil porosity. A comparison of data from the samples treated with the low rate and those treated with the high rate of poultry manure shows that the porosity in these samples was not proportional to the increase of the poultry manure added to soil. Therefore these data indicate that adding poultry manure to soils can improve the physical properties of soil, such as porosity. This parameter is very important for evaluating the effect of the application of waste materials on soil structure, and therefore it is also very important for evaluating the quality of these materials to be applied to soils.

C. Pore Shape and Size Distribution

Figure 5 represents an example of the variations of pore size and shape distribution in soil treated with organic materials compared to untreated soil. Generally, in soils supplied with animal manure the increase of elongated pores mainly involved those in the range of transmission pores (50–500 μm). Such an increase together with those of storage pores (<50 μm), was a clear sign of improved soil physical conditions and therefore of more favorable soil conditions for plant development.

Similar results were also obtained by Kladivko and Nelson (1979a), who reported an increase of porosity in the range of transmission pores following the application of sewage sludges on silty and clay loam soils, thus confirming an improvement of soil pore system. Such results also indicated the similar action of the organic matter contained in sewage sludges and in animal manure. However, several authors reported an increase of soil porosity and a decrease of bulk density after addition of animal manure (Khaleel et al., 1981; Anderson et al., 1990; Rose, 1991).

D. Type of Soil Structure

The variation of soil porosity following manure application directly reflects the soil structure. Indeed, the microscopic observations of thin sections, prepared from undisturbed soil samples collected in field experiments where the applications of manures were tested, generally reveal differences between samples of untreated soil and those treated with several types of manures. The increase of porosity and, overall, the increase of elongated pores in the range of transmission pores (50–500 μm) in soils given manures generally produced a sub-angular blocky structure homogeneously distributed along the Ap horizon (Fig. 6), which allowed, as already mentioned, more favorable conditions for plant development. The microscopic examinations of thin sections of treated soils also revealed the presence of organic matter such as wall coating of elongated





Figure 5 Pore size distribution of pores larger than 50 μm, determined by image analysis on thin section according to the equivalent pore diameter for regular and irregular pores and width for elongated pores, in control plots of the same silty clay soil of Fig. 2 and in plots treated with 300 MG ha⁻¹ of pig slurry applied at the end of May. (Modified from Pagliai and Vittori Antisari, 1993.)

pores (Fig. 7). Such organic matter coatings could effectively seal pores from the adjacent soil matrix, stabilizing soil structure against the destructive action of water, and assuring the functionality of the pores. In untreated soils the lower porosity, especially in the size range of transmission pores, produced a more compact soil structure with larger aggregates rather compact inside. In many




Figure 6 Photomicrograph of a vertically oriented thin section prepared from undisturbed soil samples collected in the surface layer (0–10 cm) of a silty clay soil treated with 300 MG ha⁻¹ of pig slurry. Picture taken under plain polarized light (pores appear white). A subangular blocky structure is evident.

cases, such as clay and silty loam soils, the large aggregates were separated by elongated pores parallel to the soil surface originating a platy structure (Fig. 8).

III. AGGREGATE STABILITY

The binding together of individual soil particles forms aggregates, and the strongly bounded aggregates give rise to what is known as well-structured soil that has greater resistance to the destructive forces of water (raindrop impact, etc.) and consequently has more resistance to erosion and degradation. Gener-



Figure 7 Photomicrograph of a vertically oriented thin section prepared from undisturbed samples collected in the surface layer (0–10 cm) of a sandy loam soil treated with farmyard manure. Plain polarized light. The presence of organic matter such as wall coating of elongated pores can be noticed.

ally, the other soil physical properties, such as porosity, water movement, and ease of root penetration will increase with increasing aggregation and aggregate stability.

The intensive crop systems of modern-day agriculture, based on continuous conventional tillage, complete vegetation removal, and copious administration of chemical fertilizers, can decrease the degree of aggregation at the surface and, on more vulnerable soils, can completely destroy surface soil structure. In these situations it is important to increase not only the aggregation but also the




Figure 8 Photomicrograph of a vertically oriented thin section prepared from undisturbed soil samples collected in the surface layer (0–10 cm) of silty clay soil not treated with organic manures (control soil). Plain polarized light. More compact soil structure is visible than in Fig. 6. The thin elongated pores are parallel to the soil surface, thus originating a rather compact platy structure. These pores are not continuous in a vertical sense and therefore not useful for water infiltration. The rounded pores are originated by entrapped air during the drying processes, and the presence of such pores is an indicator of soil structure degradation.

stability of the aggregates against the destructive forces of water. Increasing aggregation is very important on finer-textured soils such as clays, clay loams, silty clays, and silt loams. In these soils, a good water infiltration and drainage is possible if clay and silt particles are bound together into aggregates.

The addition of manures to soil has been found to be an effective management practice not only to increase the aggregation but also to increase the aggregate stability. Figure 9 shows the increase of aggregate stability in a silty clay soil following addition of pig slurry and farmyard manure. As was the case for macroporosity, the highest increase in aggregate stability was found in soils supplied with 300 MG ha⁻¹ of pig slurry in May. The winter applications of pig slurry were less efficacious in improving aggregate stability. The same trend was also observed in a sandy loam soil amended with sewage sludges, compost, and farmyard manure (Pagliai et al., 1981). Increases in aggregation and aggregate stability following long-term application of animal





Figure 9 Effect of surface application of pig slurry (PS) in February and in May and incorporation of farmyard manure (FYM) before ploughing on water stability index determined by wet-sieving. Water stability index values followed by the same letter are not significantly different at the 0.05 level. (M. Pagliai, 1993, personal communication.)

manure on sandy loam and silt loam soils were also reported by Darwish et al. (1995). Similar results were summarized by Clapp et al. (1986) following the addition of organic materials such as sewage sludge on different types of soils. The magnitude of the improvement of aggregate stability and of the other physical properties depends not only on the amount of organic matter but also on the type and quality of organic materials present in the manures added to the soil. The most efficacious in producing such an improvement is traditional farmyard manure, but the application of livestock effluents can also improve in the same way the aggregate stability even though their action is less long lasting than that of farmyard manure (Ekwue, 1992; Pagliai and Vittori Antisari, 1993). This is because of the rapid microbial degradation of these materials in soil.

IV. SOIL CRUSTING

Soil crusts are specific physical modifications in the topsoil that can be formed following natural events like raindrop impact, which cause the mechanical destruction of surface soil aggregates and the dispersion of clay materials. In the following drying processes, the formation of hard thin layers occurs. Surface crusts are very widespread especially in cultivated loam and silty loam soils of arid and semiarid regions. Their thickness usually ranges from 0.5 mm to 5 cm. When dry, these features are more compact, hard, and brittle than underlying


soil material and decrease both the size and the number of pores, reducing, in this way, water and air permeability. From the agronomic point of view the most important disadvantages of the soil crusts are the influence they have on seedling emergence and water infiltration. This latter leads, as a consequence, to an increase of surface runoff and soil erosion.



(a)

Figure 10 Photomicrograph of a vertically oriented thin section prepared from undisturbed soil samples collected in the surface layer (0–6 cm) of a silty clay soil. (a) Control soil; (b) soil treated with with 300 MG ha⁻¹ of pig slurry. Plain polarized light. A well developed surface crust is evident in the top of the control soil. It is also evident that in treated soil the addition of organic manures can reduce the surface crust formation.

Many results reported in the international literature (Pagliai, 1987a; Pagliai et al., 1985, 1987) show that the formation of soil crusts can be reduced or prevented by the landspreading of livestock effluents (Fig. 10) or the application of farmyard manure. This preventive action can be ascribed both to the organic matter that as said, increases the aggregate stability, and to the organic materials such as straw fragments that remain in the soil surface reducing the direct action of raindrop impact on surface soil aggregates. These organic materials could also break existing surface soil crusts reducing their compactness and improving water intake rates (Fig. 11).



(b)




Figure 11 Photomicrograph of a vertically oriented thin section prepared from undisturbed soil samples collected in the surface layer (0–6 cm) of a silty clay soil treated with 100 MG ha⁻¹ of pig slurry. Plain polarized light. The rest of the organic materials that break the layers of the surface crust can be noticed.

V. WATER RETENTION

The increase of microporosity in the range of storage pores (0.5–50 μm) in soil given organic material like animal manure leads, as a consequence, to an increase of water retention at both field capacity and wilting point and therefore the availability of water for plants (Rose, 1991; Schjøning et al., 1994; Giusquiani et al., 1995). The increase of water retention can be ascribed, first of all, to the improvement of the pore system in soil, which leads to better soil

structure conditions, but also to the water adsorption capacity of organic matter (Metzger and Yaron, 1987).

VI. WATER MOVEMENT

Studies on the effects of soil manure application on water movement are relatively few, owing to the difficulty and time consumption of such determinations. Moreover, results present considerable variability. However many papers report that generally in well structured soils the hydrological properties are good. Therefore the improvement of structural conditions following manure application can lead, as a consequence, to an improvement of water infiltration. Specific studies reported by Metzger and Yaron (1987) clearly show a close correlation between the increase of saturated hydraulic conductivity and the improvement of aggregate stability and porosity in soils amended with organic materials like animal manure and sewage sludge. In some cases it has been shown that the addition of organic materials with hydrorepellent substances can decrease the unsaturated hydraulic conductivity (Metzger and Yaron, 1987).

Water infiltration through the soil surface is strongly dependent on the hydraulic conductivity and increases following the addition of manures. This is an important finding, because a low water infiltration rate in the case of heavy rains leads, as a consequence, to flooding or runoff and erosion, depending on the slope. Kladivko and Nelson (1979b) described a reduction of runoff, and consequently of soil erosion following the administration of organic materials. Similar results were found by Pagliai et al. (1983a), which showed the importance of manure applications to improve aggregate stability, to reduce the formation of surface crusts, and therefore to increase water infiltration.

VII. SOIL STRENGTH, PENETRATION RESISTANCE, SOIL COMPACTION, BEARING CAPACITY

The effects of manure application on the other soil physical properties have been less investigated partly because such properties are strictly connected with the structural condition determined by the principal properties, discussed above, like porosity, aggregate stability, water retention, and water movement.

Generally, the addition of organic manures to soil significantly reduced penetration resistance when compared with control and chemical-fertilizer amended soils (Tester, 1990). The same trend was observed for the modulus of rupture, which decreased with organic matter application, especially in more cohesive soils, i.e., silt loam and clay loam soils (Darwish et al., 1995), thus indicating a reduction in soil cohesion and in the susceptibility of soil to compaction. De-

crease in soil strength with organic matter application was obtained in several studies carried out with soils amended with various types of organic matter additions (Ekwue, 1992). However, other studies have reported an increase of soil strength with organic matter addition (Schjønning et al., 1994) including a soil fertilized with farmyard manure which increased its cohesion with decreased water potential.

VIII. CONCLUSIONS

Considering soil physical parameters, such as porosity, soil structure, water retention, and water movement in soil, it can be concluded that generally the administration of manures improves the soil physical properties. Particularly, the addition of such materials causes an increase of "storage" (0.5–50 μm) and "transmission" (50–500 μm) i.e., pores that are necessary for the storage of water for plants and microorganisms and that allow water movement and root growth. Moreover, the application of manures improves the aggregate stability and reduces the formation of surface crusts. The maintenance of good soil structural conditions can be carried out not only with traditional farmyard manure but also with manures derived from livestock effluents. In this case continuous and moderate applications are necessary.

For a correct and efficient organic fertilization with manures it is fundamentally important to take into consideration the pedological environment in which such a fertilization is performed. The above-mentioned positive effects can at times be transformed into the dangerous phenomena of soil degradation. The salinity, for example, is one of the aspects that must not be undervalued in the additions of manures to soils.

REFERENCES

- Anderson, S. H., Gantzer, C. J., and Brown, J. R. (1990). Soil physical properties after 100 years of continuous cultivation. *Journal of Soil and Water Conservation* 45: 117–121.
- Brogan, J. C., ed. (1981). *Development in Plant and Soil Sciences*. Nitrogen Losses and Surface Runoff from Landspreading of Manures, Vol. 2 (J. C. Brogan, ed.). Martinus Nijhoff/Dr. W. Junk, London, p. 471.
- Clapp, C. E., Stark, S. A., Clay, D. E., and Larson, W. E. (1986). Sewage sludge organic matter and soil properties. In *The Role of Organic Matter in Modern Agriculture* (Y. Chen and Y. Avnimelech, eds.). Martinus Nijhoff, Dordrecht, pp. 209–253.
- Couteaux, M. M., Faurie, G., Palka, L., and Steimberg C. (1988). La relation prédateur proie (protozoaires–bactéries) dans le sols: rôle dans la régulation des populations et conséquences sur le cycles du carbone et de l'azote. *Rev. Ecol. Biol. Sol.* 25: 1–31.
- Darwish, O. H., Persaud, N., and Martens, D. C. (1995). Effect of long-term application of animal manure on physical properties of three soils. *Plant and Soil* 176: 289–295.

- Dexter, A. R. (1988). Advances in characterization of soil structure. *Soil Tillage Res.* 11: 199–238.
- Ekwue, E. I. (1992). Effect of organic and fertilizer treatments on soil physical properties and erodibility. *Soil Tillage Res* 22: 199–209.
- FitzPatrick, E. A., Makie, L. A., and Mullins, C. E. (1985). The use of plaster of Paris in the study of soil structure. *Soil Use Manag.* 1: 70–72.
- Guisquiani, P. L., Pagliai, M., Gigliotti, G., Businelli, D., and Benetti, A. (1995). Urban waste compost: effects on physical, chemical and biochemical soil properties. *J. Environ. Qual.* 24: 175–182.
- Greenland, D. J. (1977). Soil damage by intensive arable cultivation: temporary or permanent? *Philos. Trans. R. Soc. London* 281: 193–208.
- Greenland, D. J. (1981). Soil management and soil degradation. *J. Soil Sci.* 32: 301–322.
- Khaleel, R., Reddy, K. R., and Overcash, M. R. (1981). Changes in soil physical properties due to organic waste applications: a review. *J. Environ. Qual.* 10: 133–141.
- Kladvko, E. J., and Nelson, D. W. (1979a). Changes in soil properties from application of anaerobic sludge. *J. Water Pollution Control Federation* 51: 325–332.
- Kladvko, E. J., and Nelson, D. W. (1979b). Surface runoff from sludge-amended soils. *Jo. Water Pollution Control Federation* 51: 100–110.
- Lawrence, G. P. (1977). Measurement of pore size in fine textured soil: a review of existing techniques. *J. Soil Sci.* 28: 527–540.
- Metzger, L., and Yaron, B. (1987). Influence of sludge organic matter on soil physical properties. *Advances in Soil Science* 7: 141–163.
- Moran, C. J., McBratney, A. B., and Koppi, A. J. (1989). A rapid method for analysis of soil macropore structure. I. Specimen preparation and digital binary image production. *Soil Sci. Soc. Am. J.* 53: 921–928.
- Moran, C. J., and McBratney, A. B. (1992). Acquisition and analysis of three-component digital images of soil pore structure. II. Application to seed beds in a fallow management trial. *J. Soil Sci.* 43: 551–566.
- Murphy, C. P., Bullock, P., and Turner, R. H. (1977a). The measurement and characterization of voids in soil thin sections by image analysis: Part I. Principles and techniques. *J. Soil Sci.* 28: 498–508.
- Murphy, C. P., Bullock, P., and Biswell, K. J., (1977b). The measurement and characterization of voids in soil thin sections by image analysis: Part II. Applications. *J. Soil Sci.* 28: 509–518.
- Pagliai, M. (1987a). Effects of different management practices on soil structure and surface crusting. *Soil Micromorphology* (N. Fedoroff, L. M. Bresson, and M. A. Courty, eds.). AFES, Paris, pp. 415–421.
- Pagliai, M. (1987b). Micromorphometric and micromorphological investigations on the effect of compaction by pressures and deformations resulting from tillage and wheel traffic. *Soil Compaction and Regeneration* (G. Monnier and M. J. Goss, eds.). A. A. Balkema, Rotterdam, pp. 31–38.
- Pagliai, M., and De Nobili, M. (1993). Relationships between soil porosity, root development, and soil enzyme activity in cultivated soils. *Geoderma* 56: 243–256.
- Pagliai, M., and Vittori Antisari, L. (1993). Influence of waste organic matter on soil micro and macrostructure. *Bioresource Technology* 43: 205–213.

- Pagliai, M., Guidi, G., La Marca, M., Giachetti, M. and Lucamante, G. (1981). Effect of sewage sludges and composts on soil porosity and aggregation. *J. Environ. Qual.* 10: 556–561.
- Pagliai, M., Bisdom, E. B. A., and Ledin, S. (1983a). Changes in surface structure (crusting) after application of sewage sludges and pig slurry to cultivated agricultural soils in northern Italy. *Geoderma* 30: 35–53.
- Pagliai, M., La Marca, M., and Lucamante, G. (1983b). Micromorphometric and micro-morphological investigations of a clay loam soil in viticulture under zero and conventional tillage. *J. Soil Sci.* 34: 391–403.
- Pagliai, M., La Marca, M., Lucamante, G., and Genovese, L. (1984). Effects of zero and conventional tillage on the length and irregularity of elongated pores in a clay loam soil under viticulture. *Soil Tillage Res* 4: 433–444.
- Pagliai, M., La Marca, M., and Lucamante, G. (1985). Relationship between soil structure and time of landspreading of pig slurry. *Long-term Effects of Sewage Sludge and Farm Slurries Applications* (J. H. Williams, G. Guidi, and P. L'Hermite, eds.). Elsevier, London, pp. 45–56.
- Pagliai, M., La Marca, M., Lucamante, G. (1987). Changes in soil porosity in remoulded soils treated with poultry manure. *Soil Science* 144: 128–140.
- Pagliai, M., Pezzarossa, B., Mazzoncini, M., and Bonari, E. (1989). Effects of tillage on porosity and microstructure of a loam soil. *Soil Technology* 2: 345–358.
- Ringrose-Voase, A. J. (1991). Micromorphology of soil structure: description, quantification, application. *Aust. J. Soil Res.* 29: 777–813.
- Ringrose-Voase, A. J., and Bullock, P. (1984). The automatic recognition and measurement of soil pore types by image analysis and computer programs. *J. Soil Sci.* 35: 673–684.
- Rose, D. A. (1991). The effect of long-continued organic manuring on some physical properties of soils. *Advances in Soil Organic Matter Research*, Special Publication No. 90 (W. S. Wilson, ed.). Royal Society of Chemistry, Cambridge, pp. 197–205.
- Russel, E. W. (1978). Arable agriculture and soil deterioration. Transactions of the 11th International Congress of Soil Science, 19–27 June 1978 at University of Alberta, Edmonton, Vol. 3. Canadian Society of Soil Science, Alberta, pp. 216–227.
- Sartori, G., Ferrari, G. A., and Pagliai, M. (1985). Changes in soil porosity and surface shrinkage in a remoulded saline clay soil treated with compost. *Soil Science* 139: 523–530.
- Schjønning, P., Christensen, B. T., and Carstensen, B. (1994). Physical and chemical properties of a sandy loam receiving animal manure, mineral fertilizer, or no fertilizer for 90 years. *European J. Soil Sci.* 45: 257–268.
- Sequi, P., Cercignani, G., De Nobili, M., and Pagliai, M. (1985). A positive trend among two soil enzyme activities and a range of soil porosity under zero and conventional tillage. *Soil Biol. Biochem.* 17: 255–256.
- Shipitalo, M. J., and Protz, R. (1987). Comparison of morphology and porosity of a soil under conventional and zero tillage. *Can. J. Soil Sci.* 67: 445–456.
- Terry, R. E., Nelson, D. W., and Sommers, L. F. (1979). Decomposition of anaerobically digested sewage sludge as affected by soil environmental conditions. *J. Environ. Qual.* 8: 342–346.

- Tester, C. F., Sikora, C. J., Taylor, J. H., and Parr, J. F. (1977). Decomposition of sewage sludge compost in soil: 1. Carbon and nitrogen transformation. *J. Environ. Qual.* 6: 459–463.
- Tester, C. F. (1990). Organic amendment effects on physical and chemical properties of a sandy soil. *Soil Sci. Soc. Am. J.* 54: 827–831.
- Tippkötter, R. (1983). Morphology, spatial arrangement and origin of macropores in some hapludalfs, West Germany. *Geoderma* 29: 355–371.
- Utomo, W. M., and Dexter, A. R. (1982). Changes in soil aggregate water stability induced by wetting and drying cycles in non-saturated soil. *J. Soil Sci.* 33: 623–637.