



SMR.770/5

ADVANCED WORKSHOP ON ALGEBRAIC GEOMETRY
(15 - 26 August 1994)

Vector bundles and the Brill-Noether theory

S. Mukai

Department of Mathematics
School of Science
Nagoya University
464-01 Furo-cho, Chikusa-ku
Nagoya
Japan

These are preliminary lecture notes, intended only for distribution to participants

Vector bundles and the Brill-Noether theory

Shigeru MUKAI*

March 8, 1994

As a natural generalization of line bundles, vector bundles have two important roles in Algebraic Geometry. One is the moduli space. The moduli of vector bundles gives connections among different types of varieties, and sometimes yield new varieties which are difficult to describe by other means. The other is the linear system. In the same way as the classical construction of a map to a projective space, a vector bundle gives rise to a rational map to a Grassmannian if it is generically generated by its global sections. In this article, we shall describe some results for which vector bundles play such roles. They are obtained from an attempt to generalize the Brill-Noether theory of special divisors, reviewed in §2, to vector bundles. Our main subject is rank 2 vector bundles with canonical determinant on a curve C with as many global sections as possible: especially their moduli and the Grassmannian embeddings of C by them (§4).

1 Line bundles

Let X be a smooth algebraic variety over the complex number field \mathbb{C} . We consider the set of isomorphism classes of line bundles, or invertible sheaves, on X . This set enjoys two good properties, neither of which holds any more for vector bundles of higher rank. One is that it has a natural algebraic structure as a moduli space without any modification. The other is that it becomes a (commutative) group by the tensor product. In fact, the isomorphism classes are parametrized by the first cohomology group $H^1(\mathcal{O}_X^*)$ with coefficient in the (multiplicative) sheaf of nowhere vanishing holomorphic functions. $H^1(\mathcal{O}_X^*)$ endowed with the natural algebraic structure is called the *Picard variety* and denoted by $\text{Pic } X$. Let

$$\cdots \longrightarrow H^1(X, \mathbb{Z}) \longrightarrow H^1(\mathcal{O}_X) \longrightarrow H^1(\mathcal{O}_X^*) \xrightarrow{\delta} H^2(X, \mathbb{Z}) \longrightarrow \cdots \quad (1.1)$$

be the long exact sequence derived from the exponential exact sequence

$$0 \longrightarrow \mathbb{Z} \xrightarrow{2\pi i} \mathcal{O}_X \xrightarrow{\exp} \mathcal{O}_X^* \longrightarrow 1 \quad (1.2)$$

of sheaves on X . The connecting homomorphism δ associates the first Chern class $c_1(L)$ for each line bundle $[L] \in H^1(\mathcal{O}_X^*)$. For example, if X is a curve, then $\delta(L)$ is the degree of

*Based on the author's three talks given at JAMI in 1991, UCLA in 1992 and Durham University in 1993. Supported in part by a Grant under The Monbusho International Science Research Program: 04044081.

L under the natural identification $H^2(X, \mathbf{Z}) \simeq \mathbf{Z}$. By (1.1), the neutral component $\text{Pic}^0 X$ of $\text{Pic} X$ is isomorphic to the quotient group $H^1(\mathcal{O}_X)/H^1(X, \mathbf{Z})$, which is an abelian variety if X is a projective variety.

Let C be a *curve*, or a compact Riemann surface, of genus g . The Riemann-Roch theorem

$$\begin{cases} \chi(L) := h^0(L) - h^1(L) = \deg L + 1 - g \\ H^1(L) \simeq H^0(K_C L^{-1})^\vee, \end{cases} \quad (1.3)$$

is most fundamental for its study. The latter isomorphism is functorial in L and referred as the *Serre duality*. By (1.1), $\text{Pic} C$ is the disjoint union of $\text{Pic}^d C$, $d \in \mathbf{Z}$, where $\text{Pic}^d C$ is the set of isomorphism classes of line bundles of degree d . By (1.3), the number $h^0(L)$ of linearly independent global sections is constant on $\text{Pic}^d C$ unless $0 \leq d \leq 2g-2 = \deg K_C$. Conversely, when $0 \leq d \leq 2g-2$, $h^0(L)$ is equal to $d+1-g$ on a non-empty Zariski open subset of $\text{Pic}^d C$, but not constant since there exists a *special line bundle*, i.e., a line bundle L with $h^0(L)h^1(L) \neq 0$, of degree d . $\text{Pic}^d C$ is stratified by $h^0(L)$. We set

$$W_d^r(C) = \{[L] | h^0(L) \geq r+1\} \subset \text{Pic}^d C,$$

which is closed in the Zariski topology. The case $(d, r) = (g-1, 0)$ is most important. W_{g-1}^0 is a divisor and usually denoted by Θ . The self intersection number (Θ^g) is equal to $g!$, that is, Θ is a principal polarization of $\text{Pic}^{g-1} C$. This principally polarized abelian variety $(\text{Pic}^{g-1} C, \Theta)$ is called the *Jacobian* of C .

Often the isomorphism class of C is recovered from the variety $W_d^r(C)$ of special line bundles. The case of theta divisor Θ is classical:

Theorem 1.4 (Torelli) *Two curves are isomorphic to each other if their Jacobians are so.*

We refer [13] for various approaches to this important result. Let C be a non-hyperelliptic curve of genus 5. Then $W_4^1(C)$ is a curve[†] of genus 11. Another example is

Theorem 1.5 *The Jacobian of C is isomorphic to the Prym variety of $(W_4^1(C), \sigma)$, where σ is the involution of $\text{Pic}^4 C$ defined by $\sigma[L] = [K_C L^{-1}]$.*

See [2] for the proof in the case C is a complete intersection of three quadrics in \mathbf{P}^5 .

Another feature of special line bundles is their relation with projective embeddings. If a line bundle L is generated by its global sections, then we obtain a morphism

$$\Phi_{|L|} : C \longrightarrow \mathbf{P}^* H^0(L),$$

where $\mathbf{P}^* H^0(L)$ is the projectivization of the dual vector space of $H^0(L)$. The most interesting case is K_C , the canonical line bundle, which appears in (1.3). By the Riemann-Roch theorem, K_C is generated by global sections, and $\Phi_{|K_C|} : C \longrightarrow \mathbf{P}^* H^0(K_C) = \mathbf{P}^{g-1}$ is an embedding unless C is hyperelliptic. The image $C_{2g-2} \subset \mathbf{P}^{g-1}$ of $\Phi_{|K_C|}$ is called the *canonical model* of C . The following is a classical example:

Theorem 1.6 (Enriques-Petri) *The canonical model $C_{2g-2} \subset \mathbf{P}^{g-1}$ is an intersection of quadrics if and only if $W_3^1(C) = W_5^2(C) = \emptyset$.*

We refer [1] and [7] for further results of this kind. The latter discusses also an interesting use of vector bundles which we do not treat here.

[†]If C is trigonal or $W_4^1(C)$ contains a line bundle with $L^2 \simeq K_C$, then $W_4^1(C)$ is singular. But still the theorem holds true.

2 Brill-Noether theory

We study $W_d^r(C)$ more closely. First we note that it is not only a subset but a subscheme of $\text{Pic } C$. Take distinct points P_1, \dots, P_N of C and put $D = \sum_{i=1}^N P_i$. We choose N sufficiently large so that $H^1(L(D))$ vanishes for every $[L] \in \text{Pic}^d C$. The exact sequence

$$0 \longrightarrow L \longrightarrow L(D) \xrightarrow{\text{res}} \bigoplus_{i=1}^N L(D)|_{P_i} \longrightarrow 0 \quad (2.1)$$

of sheaves on C induces the exact sequence

$$0 \longrightarrow H^0(L) \longrightarrow H^0(L(D)) \xrightarrow{H^0(\text{res})} \bigoplus_{i=1}^N H^0(L(D)|_{P_i}) \longrightarrow H^1(L) \longrightarrow 0 \quad (2.2)$$

of vector spaces. There exists a homomorphism $R : \mathcal{E} \longrightarrow \mathcal{F}$ between two vector bundles[†] \mathcal{E} and \mathcal{F} on $\text{Pic}^d C$ whose fibre $R_{[L]}$ at $[L]$ is $H^0(\text{res})$ for every $[L] \in \text{Pic}^d C$. The difference of ranks of two vector bundles does not depend on D . In fact, we have

$$r(\mathcal{F}) - r(\mathcal{E}) = N - h^0(L(D)) = g - 1 - d$$

by (1.3). The following is easy to verify:

Lemma 2.3 *Let E and F be finite dimensional vector spaces, c a positive integer and set*

$$W = \{f \in \text{Hom}(E, F) \mid \dim \text{Ker } f \geq c\}.$$

Then we have

(1) *W is a closed subvariety of codimension $c(c + \delta)$ in the affine space $\text{Hom}(E, F)$, where $\delta = \dim F - \dim E$, and*

(2) *if $\dim \text{Ker } f = c$, then W is smooth at the point f and the normal space $N_{W/\text{Hom}, f}$ is isomorphic to $\text{Hom}(\text{Ker } f, \text{Coker } f)$.*

Since $W_d^r(C)$ is

$$\{\alpha \in \text{Pic}^d C \mid \dim \text{Ker } R_\alpha \geq r + 1\},$$

it is a closed subscheme of $\text{Pic}^d C$ and its codimension is at most $(r + 1)(g + r - d)$ by the lemma. It follows that

$$\dim W_d^r(C) \geq g - (r + 1)(g + r - d). \quad (2.4)$$

For a line bundle L on C , we put $\rho(L) = g - h^0(L)h^1(L)$ and call it the *Brill-Noether number*. When $[L] \in W_d^r(C)$ and $h^0(L) = r + 1$, then $\rho(L)$ is equal to the right hand side of the above inequality. Since the tangent space of $\text{Pic } C$ is isomorphic to $H^1(\mathcal{O}_C)$ by (1.1), the Zariski tangent space of $W_d^r(C)$ at $[L]$ is the kernel of the tangential map $H^1(\mathcal{O}_C) \longrightarrow \text{Hom}(H^0(L), H^1(L))$ by (2.2) and (2) of the lemma.

Now we describe the Zariski tangent space more directly. Let

$$\tau_L : H^1(\mathcal{O}_C) \longrightarrow \text{Hom}(H^0(L), H^1(L))$$

[†]Both \mathcal{E} and \mathcal{F} are the direct images of certain sheaves on $C \times \text{Pic } C$.

be the linear map induced by the cup product $H^1(\mathcal{O}_C) \times H^0(L) \longrightarrow H^1(L)$. By the Serre duality (1.3), the dual of τ_L is the multiplication map

$$H^0(L) \otimes H^0(K_C L^{-1}) \longrightarrow H^0(K_C), \quad (2.5)$$

called the *Petri map*.

Proposition 2.6 *Assume that $[L] \in W_d^r(C)$ and $h^0(L) = r + 1$. Then we have*

- (1) *the Zariski tangent space of $W_d^r(C)$ at $[L]$ is isomorphic to $\text{Ker } \tau_L$,*
- (2) *$\dim \text{Ker } \tau_L \geq \dim_{[L]} W_d^r(C) \geq \rho(L)$, and*
- (3) *the following three conditions are equivalent to each other:*
 - i) *$W_d^r(C)$ is smooth and of dimension $\rho(L)$ at $[L]$,*
 - ii) *τ_L is surjective, and*
 - iii) *the Petri map (2.5) is injective,*

Proof. Let $\alpha = \{a_{ij}\} \in H^1(\mathcal{O}_C^*)$ be the cohomology class corresponding to L , that is, $a_{ij} \in \mathcal{O}_{U_i \cap U_j}$ are the transition functions of L for a suitable open covering $\{U_i\}$ of C . Let ε be the dual number, i.e., $\varepsilon \neq 0$ but $\varepsilon^2 = 0$. A first order infinitesimal deformation \tilde{L} of L corresponds to a cohomology class $\tilde{\alpha} = \{\tilde{a}_{ij}\} \in H^1(\mathcal{O}_C[\varepsilon]^*)$ whose reduction modulo ε is α . $\tilde{\alpha}$ is of the form $\{a_{ij}(1 + b_{ij}\varepsilon)\}$ for $\beta = \{b_{ij}\} \in H^1(\mathcal{O}_C)$. Let $h \in H^0(L)$ be a global section of L . h is a collection $\{h_i\}$ of $h_i \in \mathcal{O}_{U_i}$ such that $h_i = a_{ij}h_j$. The differences $b_{ij}h_j\varepsilon$ of h_i and $\tilde{a}_{ij}h_j$ form a 1-cocycle, whose cohomology class is the cup product $(\beta^{\cup} h)\varepsilon$. Hence h extends to a global section of \tilde{L} if and only if $\beta^{\cup} h = 0$ in $H^1(L)$. Therefore, all global sections of L extend if and only if the cup product map $\cup \beta : H^0(L) \longrightarrow H^1(L)$ is zero, which shows (1). (2) follows from (1) and (2.4). (3) is straightforward from (2). \square

Let ρ be the right hand side of (2.4). We refer [1] for the following important results:

Theorem 2.7 (*Existence*) $W_d^r(C) \neq \emptyset$ if $\rho \geq 0$.
(Connectedness) $W_d^r(C)$ is connected if $\rho > 0$.

Let \mathcal{M}_g be the moduli space of curves of genus g .

Theorem 2.8 ([5], [6]) *If $[C] \in \mathcal{M}_g$ is general, then the Petri map (2.5) is injective for every (special) line bundle L on C .*

In particular, $W_d^r(C)$ is of dimension ρ if ρ is nonnegative, and empty otherwise. Thus the estimate (2.4) is best possible for the generic curve. When $\rho = 0$, the number of $W_d^r(C)$ is finite and was first computed by Castelnuovo. Let $G(a, a+b)$ be the Grassmannian of a -dimensional subspaces of \mathbb{C}^{a+b} and $G(a, a+b) \subset \mathbb{P}_* \wedge^a \mathbb{C}^{a+b}$ its Plücker embedding. The following is especially interesting (cf. (4.15)):

Theorem 2.9 *If $[C] \in \mathcal{M}_g$ is general and $\rho = 0$, then the number of $W_d^r(C)$ is equal to the degree of the g -dimensional Grassmannian $G(a, a+b) \subset \mathbb{P}_* \wedge^a \mathbb{C}^{a+b}$, where $a = r + 1$ and $b = g + r - d$.*

In fact, both $\#W_d^r(C)$ and $\deg G(a, a+b)$ are equal to $g! \prod_{1 \leq i \leq a < j \leq a+b} (j - i)^{-1}$.

3 Vector bundles on a curve

Let X be a smooth complete algebraic variety (over \mathbb{C}). By \mathcal{O}_X , we mean either the sheaf of holomorphic functions in the usual topology or the sheaf of regular functions in the Zariski topology. In the study of vector bundles, this is allowed by virtue of the GAGA principle, which says that the two categories of analytic and algebraic coherent sheaves on X are equivalent to each other. By a *vector bundle* E on X , we mean a locally free \mathcal{O}_X -module. There exists an open covering $\{U_i\}$ of X such that $E|_{U_i} \simeq \mathcal{O}_{U_i}^{\oplus r}$ for every i . This positive integer r is called the *rank* of E . The highest exterior product $\bigwedge^r E$ is a line bundle on X , which is denoted by $\det E$.

Assume that E is generated by global sections, that is, the evaluation homomorphism

$$ev_E : H^0(E) \otimes \mathcal{O}_X \longrightarrow E$$

is surjective. Then every fibre E_x of E is an r -dimensional quotient space of $H^0(E)$. Hence we obtain a map $\Phi_{|E|} : X \longrightarrow G(H^0(E), r)$ to the Grassmannian of r -dimensional quotient spaces of $H^0(E)$. This map is holomorphic since E is so. The exterior product

$$\bigwedge^r ev_E : \bigwedge^r H^0(E) \otimes \mathcal{O}_X \longrightarrow \bigwedge^r E$$

of ev_E induces a linear map

$$\bigwedge^r H^0(E) \longrightarrow H^0(\bigwedge^r E), \quad (3.1)$$

which we denote by λ_E . The exterior product $\bigwedge^r E_x$ of fibres E_x are quotient spaces of both $H^0(\bigwedge^r E)$ and $\bigwedge^r H^0(E)$. The former determines a point in the projective space $\mathbb{P}^* H^0(\bigwedge^r E)$ and the latter the Plücker coordinate of $[E_x] \in G(H^0(E), r)$. Hence we obtain the following commutative diagram:

$$\begin{array}{ccccc} & X & \xrightarrow{\Phi_{|E|}} & G(H^0(E), r) & \\ \Phi_{|\bigwedge^r E|} \downarrow & \downarrow & & \cap & \text{Plücker embedding} \\ \mathbb{P}^* H^0(\bigwedge^r E) & \xrightarrow{\mathbb{P}^* \lambda_E} & \mathbb{P}^* \bigwedge^r H^0(E). & & \end{array} \quad (3.2)$$

Thus λ_E connects the projective and Grassmannian embeddings. This map is important in other consideration, too. See (3.5), (4.7) and (4.15).

Let $A_{ij} \in GL(r, \mathcal{O}_{U_i \cap U_j})$ be the transition matrix function of a vector bundle E , that is, the matrix expression of the composite of the two isomorphisms $\mathcal{O}_{U_j}^{\oplus r} \simeq E|_{U_j}$ and $E|_{U_i} \simeq \mathcal{O}_{U_i}^{\oplus r}$ over $U_i \cap U_j$. The collection $\{A_{ij}\}$ of all such (matrix) functions is a 1-cocycle with coefficient in the sheaf $GL(r, \mathcal{O}_X)$ of non-commutative groups. Hence the rank r vector bundles on X are parametrized by the first cohomology set $H^1(GL(r, \mathcal{O}_X))$. The determinant homomorphism $\det : GL(r, \mathcal{O}_X) \longrightarrow \mathcal{O}_X^*$ induces the map

$$H^1(\det) : H^1(GL(r, \mathcal{O}_X)) \longrightarrow H^1(\mathcal{O}_X^*) = \text{Pic } X,$$

whose fibre at $[L] \in \text{Pic } X$ is denoted by $B_X(r, L)$.

Moduli problem: a) Give a natural algebraic structure to a suitable *open* subset of $B_X(r, L)$ and construct its *geometric* compactification.

b) What properties of (X, L) are inherited by the moduli space constructed in a)?

$B_X(r, L)$ does not have a nice description such as $\text{Pic } X$ in (1.1). But its Zariski tangent space is easy to identify. Let E be a vector bundle and $\{A_{ij}\}$ the 1-cocycle of transition functions. A 1-cochain $\{A_{ij}(I_r + B_{ij}\varepsilon)\}$ with values in $GL(r, \mathcal{O}_X[\varepsilon])$ is a cocycle if and only if

$$A_{jk}^{-1} B_{ij} A_{jk} + B_{jk} = B_{ik}$$

holds on $U_i \cap U_j \cap U_k$ for every i, j and k , where ε is the dual number. This is the same as saying that $\{B_{ij}\}$ is a 1-cocycle with values in the sheaf $\mathcal{E}nd E \simeq E^\vee \otimes E$ of (local) endomorphisms of E . By this correspondence, the first order infinitesimal deformations of E are parametrized by the cohomology group $H^1(\mathcal{E}nd E)$. Since

$$\det(A_{ij}(I_r + B_{ij}\varepsilon)) = (\det A_{ij})(I_r + \text{tr } B_{ij}\varepsilon),$$

the Zariski tangent space of $B_X(r, L)$ is isomorphic to the kernel of

$$H^1(\text{tr}) : H^1(\mathcal{E}nd E) \longrightarrow H^1(\mathcal{O}_X),$$

which is the tangential map of $H^1(\det)$. Let $sl(E)$ be the sheaf of traceless endomorphisms of E . Then $\mathcal{E}nd E$ is the direct sum of two vector bundles $sl(E)$ and \mathcal{O}_X . Therefore, the Zariski tangent space is isomorphic to $H^1(sl(E))$.

Let C be a curve of genus g . The Riemann-Roch theorem (1.3) is generalized to

$$\begin{cases} \chi(E) := h^0(E) - h^1(E) = \deg E + r(1 - g) \\ H^1(E) \simeq H^0(K_C E^\vee)^\vee \text{ (Serre duality),} \end{cases} \quad (3.3)$$

where $\deg E$ is the degree of $\det E$. For simplicity, we restrict ourselves to the case $r = 2$. The answer to a) of the moduli problem is the notion of stability:

Definition 3.4 ([11]) A rank 2 vector bundle E on C is *stable* (resp. *semi-stable*) if $\deg \xi < \frac{1}{2} \deg E$ (resp. \leq) for every line subbundle ξ of E .

Let L be a line bundle on C and E a member of $B_C(2, L)$. Fix an ample line bundle $\mathcal{O}_C(1)$ on C . $E(n)$ belongs to $B_C(2, L(2n))$ and we obtain the linear map

$$\lambda_{E(n)} : \bigwedge^2 H^0(E(n)) \longrightarrow H^0(L(2n)) \quad (3.5)$$

as in (3.1). The above condition $\deg \xi < \frac{1}{2} \deg E$ is equivalent to the asymptotic stability of the linear map $\lambda_{E(n)}$ with respect to the action of the special linear group $SL(H^0(E(n)))$ ([4]). By the geometric invariant theory, the (coarse) moduli space $M_C(2, L)$ of stable 2-bundles with determinant L exists as a quasi-projective algebraic variety. Moreover, it becomes a projective algebraic variety $\overline{M}_C(2, L)$ by adding certain equivalence classes of semi-stable 2-bundles.

Every line bundle M induces an isomorphism $M_C(2, L) \simeq M_C(2, LM^2)$, $E \mapsto E \otimes M$. Hence there are only two isomorphism classes of the moduli space of 2-bundles: $M_C(2, \text{odd})$ and $M_C(2, \text{even})$. Both are smooth and of dimension $3g - 3 = \dim H^1(sl(E))$. Since every semi-stable 2-bundle of odd degree is stable, $M_C(2, \text{odd})$ is a projective variety. Among many known global properties of $M_C(2, \text{odd})$, we state two. One is

Theorem 3.6 ([19]) $M_C(2, \text{odd})$ is a Fano manifold of index two.

A smooth projective variety X is called *Fano* if the anti-canonical line bundle $K_X^{-1} = \det T_X$ is ample. The largest integer which divides $c_1(X)$ in $H^2(X, \mathbb{Z})$ is called the index. In the case of $M_C(2, \text{odd})$, the Picard group is free cyclic and the anti-canonical line bundle is the square of the positive generator.

Example ([16], [17]) Let C be a curve of genus 2 defined by the equation $y^2 = (x - \lambda_1)(x - \lambda_2) \cdots (x - \lambda_6)$. Then the moduli space $M_C(2, \text{odd})$ is the complete intersection

$$\sum_{i=1}^6 x_i^2 = \sum_{i=1}^6 \lambda_i x_i^2 = 0$$

in \mathbb{P}^5 . The anti-canonical line bundle is the square of the restriction of tautological line bundle.

The other is a Torelli type theorem:

Theorem 3.7 ([15]) *The intermediate Jacobian $H^2(\Omega_M^1)/H^3(M, \mathbb{Z})$ of the moduli space $M_C(2, \text{odd})$ is isomorphic to the Jacobian of C as polarized abelian variety.*

4 Toward a Brill-Noether type theory for 2-bundles

$M_C(2, \mathcal{O}_C)$ and $M_C(2, K_C)$ are two natural representatives of $M_C(2, \text{even})$. The following is fundamental for the former:

Theorem 4.1 ([16], [3]) *There is a natural bijection between the following two sets:*

- 1) $M_C(2, \mathcal{O}_C)$, the set of isomorphism classes of stable 2-bundles with trivial determinant, and
- 2) $\text{Hom}^{\text{irr}}(\pi_1(C), SU(2))/SU(2)$, the set of conjugacy classes of irreducible 2-dimensional special unitary representations of the fundamental group of C .

We take the latter to develop a Brill-Noether type theory. See [20] and [18] for another direction of the development. The moduli space $M_C(2, K)$ of stable 2-bundles with canonical determinant is stratified by the number $h^0(E)$ of linearly independent global sections. As an analogy of W_d^r , we set

$$M_C(2, K, n) = \{[E] | h^0(E) \geq n + 2\} \subset M_C(2, K).$$

We denote by $\overline{M}_C(2, K, n)$ the union of $M_C(2, K, n)$ and the set of isomorphism classes of semi-stable vector bundles $[\xi \oplus K_C \xi^{-1}] \in \overline{M}_C(2, K)$ with $[\xi] \in W_{g-1}^{n/2}(C)$.

By the same argument[§] as in §2, $M_C(2, K, n)$ is a closed subscheme of $M_C(2, K)$. A similar consideration gives the estimate $\dim M_C(2, K, n) \geq 3g - 3 - (n + 2)^2$. But this is not sharp. The proper one is

[§]The universal family does not exist on $C \times M_C(2, K)$. But this does not cause a problem for the study of such local properties of the moduli.

Theorem 4.2 $\dim M_C(2, K, n) \geq 3g - 3 - \frac{1}{2}(n+2)(n+3)$.

Before giving the proof, we recall some notion of symplectic geometry. Let W be a 2ν -dimensional vector space with a non-degenerate skew-symmetric bilinear form $\langle, \rangle: W \times W \rightarrow \mathbb{C}$. A ν -dimensional subspace V of W is a *Lagrangian* of W if the bilinear form \langle, \rangle is identically zero on $V \times V$. We denote the set of Lagrangeans by $\mathcal{L}(W)$, which is a subset of the ν^2 -dimensional Grassmannian $G(\nu, W)$. Fix a Lagrangean U_∞ and set $Z = \{[U] | U \cap U_\infty = 0\}$ in $G(\nu, W)$. When $[U_0] \in Z$ is fixed, Z is a ν^2 -dimensional affine space by the bijection

$$\text{Hom}(U_0, U_\infty) \ni f \mapsto \Gamma_f \in Z, \quad (4.3)$$

where $\Gamma_f \subset U_0 \times U_\infty$ is the graph of f . Assume that U_0 is also a Lagrangean. Then U_0 and U_∞ are each other's dual by the pairing \langle, \rangle . Γ_f is a Lagrangean of W if and only if $f \in \text{Hom}(U_0, U_\infty) \simeq U_\infty \otimes U_0^*$ is symmetric. Hence $\mathcal{L}(W)$ is a smooth subvariety of dimension $\frac{1}{2}\nu(\nu+1)$ in the Grassmannian $G(\nu, W)$.

Proposition 4.4 *Fix a Lagrangean $[V_0] \in \mathcal{L}(W)$ and let c be a positive integer. Then the Schubert subvariety*

$$\mathcal{L}(W)_c = \{[V] \in \mathcal{L}(W) | \dim V \cap V_0 \geq c\}$$

is of codimension $\frac{1}{2}c(c+1)$ in $\mathcal{L}(W)$.

Proof. For $[V] \in \mathcal{L}(W)$, choose a Lagrangean V_∞ so that $V \cap V_\infty = V_0 \cap V_\infty = 0$. Then V corresponds to a symmetric matrix of size ν via (4.3) and $\dim V \cap V_0$ is equal to the co-rank of the symmetric matrix. Hence we have our assertion. \square

Proof of Theorem 4.2: Let E be a 2-bundle with canonical determinant and D an effective divisor on C . Since E is self-Serre adjoint, i.e., $E^\vee K_C \simeq E$, the two vector spaces $H^0(E)$ and $H^1(E)$ are each other's dual by (3.3). Similarly, so are $H^0(E(-D))$ and $H^1(E(D))$. Hence, by the Riemann-Roch theorem (3.3), we have

$$h^0(E(D)) - h^0(E(-D)) = \chi(E(D)) = 2N, \quad (4.5)$$

where $N = \deg D$. Now we denote the quotient sheaf $E(D)/E(-D)$ by A , which is supported by a finite set and of length $4N$. We consider the composite of the pairing $A \times A \rightarrow K_C(2D)/K_C$, induced by $\wedge^2 E \simeq K_C$, and the residue map

$$r: K_C(2D)/K_C \rightarrow \mathbb{C}, \quad \omega \mapsto \sum_{P \in \text{Supp } D} \text{Res}_P \omega.$$

This induces a non-degenerate skew-symmetric pairing \langle, \rangle on the vector space $H^0(A)$ of dimension $4N$. Since r is identically zero on the image of $H^0(K_C(2D))$ (Residue Theorem), so is \langle, \rangle on the image V of $H^0(E(D)) \rightarrow H^0(A)$. By the exact sequence

$$0 \rightarrow E(-D) \rightarrow E(D) \rightarrow A \rightarrow 0,$$

the image V is isomorphic to the quotient space $H^0(E(D))/H^0(E(-D))$. Hence V is a Lagrangean of the symplectic vector space $H^0(A)$ by (4.5). It is obvious that $V_0 = H^0(E/E(-D))$ is also a Lagrangean. Now we choose D so that $H^0(E(-D)) = 0$ for every $[E] \in M_C(2, K)$. Then $H^0(E)$ is the intersection of two Lagrangeans $V = H^0(E(D))$ and V_0 of $H^0(A)$. Hence we have our inequality by the above proposition. \square

Remark 4.6 The above proof works also for a vector bundle E of even rank with a non-degenerate skew-symmetric pairing $E \times E \rightarrow K_C$. The same trick was used in [12] to show the parity preservation of $h^0(E)$ when E has a non-degenerate quadratic form $q : E \rightarrow K_C$.

Let E be a rank 2 vector bundle on C . For $f \in \text{Hom}(H^0(E), H^1(E))$, let $T(f)$ be the linear map

$$\bigwedge^2 H^0(E) \ni h_1 \wedge h_2 \mapsto h_1 \cup f(h_2) - h_2 \cup f(h_1) \in H^1(\bigwedge^2 E),$$

where $\cup : H^0(E) \times H^1(E) \rightarrow H^1(\bigwedge^2 E)$ is the cup product. It is easy to check that the following diagram is commutative:

$$\begin{array}{ccc} H^1(\mathcal{E}nd E) & \longrightarrow & \text{Hom}(H^0(E), H^1(E)) \\ H^1(\text{tr}) \downarrow & & \downarrow \\ H^1(\mathcal{O}_C) & \longrightarrow & \text{Hom}(\bigwedge^2 H^0(E), H^1(\bigwedge^2 E)), \end{array} \quad T \quad (4.7)$$

where the lower horizontal linear map is the composite of

$$\tau_{\det E} : H^1(\mathcal{O}_C) \rightarrow \text{Hom}(H^0(\bigwedge^2 E), H^1(\bigwedge^2 E)),$$

defined in §2, and $\text{Hom}(\lambda_E, H^1(\bigwedge^2 E))$.

Theorem 4.8 Assume that $[E] \in M_C(2, K, n)$ and $h^0(E) = n + 2$. Then we have

- (1) the Zariski cotangent space of $M_C(2, K, n)$ at $[E]$ is isomorphic to the cokernel of $S^2 H^0(E) \rightarrow H^0(S^2 E)$,
- (2) $\dim \text{Coke}[S^2 H^0(E) \rightarrow H^0(S^2 E)] \geq \dim_{[E]} M_C(2, K, n) \geq \sigma(E)$, and
- (3) $M_C(2, K, n)$ is smooth and of dimension $\sigma(E)$ at $[E]$ if and only if $S^2 H^0(E) \rightarrow H^0(S^2 E)$ is injective.

Proof. As we saw in §3, the tangent space of $M_C(2, K)$ at $[E]$ is the kernel $H^1(\mathfrak{sl}(E))$ of the trace map $H^1(\mathcal{E}nd E) \rightarrow H^1(\mathcal{O}_C)$. Let \tilde{E} be a first order infinitesimal deformation of E corresponding to $B = \{B_{ij}\} \in H^1(\mathcal{E}nd E)$. By the same argument as in the proof of Proposition 2.6, a global section $h \in H^0(E)$ extends that of \tilde{E} if and only if the cup product $h \cup B \in H^1(E)$ vanishes. Hence, all global sections of E extend if and only if the cup product map $\cup B : H^0(E) \rightarrow H^1(E)$ is zero. Therefore, the Zariski tangent space of $M_C(2, K, n)$ is the kernel of $H^1(\mathfrak{sl}(E)) \rightarrow \text{Ker } T \subset \text{Hom}(H^0(E), H^1(E))$. Since $\bigwedge^2 E \simeq K_C$, the dual of (4.7) is reduced to the commutative diagram

$$\begin{array}{ccc} H^0(E \otimes E) & \longleftarrow & H^0(E) \otimes H^0(E) \\ \uparrow & & \uparrow \\ H^0(\bigwedge^2 E) & \xleftarrow{\lambda_E} & \bigwedge^2 H^0(E), \end{array}$$

by the Serre duality (3.3), which shows (1). (2) follows from (1) and Theorem 4.2. (3) is straightforward from (1) and (2). \square

Remark 4.9 The obstructions for $M_C(2, K, n)$ to be smooth at $[E]$ lie in the cokernel of $H^0(S^2 E)^\vee \rightarrow S^2 H^0(E)^\vee$. This fact gives another proof of Theorem 4.2 and 4.8.

Let σ be the right hand side of Theorem 4.2. Theorem 2.7, 2.8 and 2.9 lead us to the following problems:

Problem 4.10 (Existence) Is $\overline{M}_C(2, K, n)$ non-empty when $\sigma \geq 0$?
(Connectedness) Is $\overline{M}_C(2, K, n)$ connected when $\sigma > 0$?

Problem 4.11 Assume that $[C] \in \mathcal{M}_g$ is general.

- (1) Is $S^2 H^0(E) \longrightarrow H^0(S^2 E)$ injective for every $E \in M_C(2, K)$?
- (2) Is $M_C(2, K, n)$ of dimension σ when $\sigma \geq 0$?
- (3) Compute the number of $M_C(2, K, n)$ when $\sigma = 0$. More generally, describe the cohomology class of $\overline{M}_C(2, K, n)$ in $H^*(\overline{M}_C(2, K), \mathbf{Z})$ when $\sigma > 0$.

Another direction is

Problem 4.12 Study the Grassmannian map associated to a member of $M_C(2, K, n)$ and its relation with the canonical model $C_{2g-2} \subset \mathbf{P}^{g-1}$.

We give some sample results in these directions. They are closely related with our classification of Fano 3-folds via vector bundles (cf. [8]). We first consider the three cases $(g, n+2) = (7, 5), (9, 6)$ and $(11, 7)$. The Brill-Noether number σ is equal to 3, 3 and 2, respectively.

Theorem 4.13 *Let C be a curve of genus 7 with $W_4^1(C) = \emptyset$. Then we have*

- (1) $M_C(2, K, 3)$ is smooth, complete and of dimension 3,
- (2) $M_C(2, K, 3)$ is a Fano 3-fold of genus 7, i.e., $(-K_M)^3 = 12$, and with Picard number one, and
- (3) the intermediate Jacobian $H^2(\Omega_M^1)/H^3(M, \mathbf{Z})$ of $M_C(2, K, 3)$ is isomorphic to the Jacobian of C as polarized abelian variety.

Conversely, every smooth Fano 3-fold of genus 7 with Picard number one is obtained in this manner from a non-tetragonal curve of genus 7.

Similarly, $\overline{M}_C(2, K, 4)$ is a quartic 3-folds in \mathbf{P}^4 with 21 singular points at the boundary if C is a general curve of genus 9, and $M_C(2, K, 5)$ is a (polarized) K3 surface of genus 11 if C is a general curve of genus 11.

In the case $(g, n+2) = (8, 6)$, the number σ is equal to zero.

Theorem 4.14 ([9], [10]) *If C is a curve of genus 8 with $W_7^2(C) = \emptyset$, then $M_C(2, K, 4)$ consists of the unique isomorphism class of stable 2-bundles E . The linear map λ_E in (3.1) is surjective and the following diagram, essentially (3.2), is Cartesian:*

$$\begin{array}{ccccc} & C & \xrightarrow{\Phi_{|E|}} & G(H^0(E), 2) & \\ \text{canonical embedding} & \cap & & \cap & \text{Plücker embedding} \\ \mathbf{P}^\bullet H^0(K_C) & \xrightarrow{\mathbf{P}^\bullet \lambda_E} & \mathbf{P}^\bullet \wedge^2 H^0(E). & & \end{array}$$

In particular, C is a complete linear section of the 8-dimensional Grassmannian, that is,

$$[C \subset \mathbf{P}^7] = [G(6, 2) \subset \mathbf{P}^{14}] \cap H_1 \cap \cdots \cap H_7$$

for seven hyperplanes $H_i, 1 \leq i \leq 7$.

Let C and E be as in the theorem and consider the intersection of $G(2, H^0(E))$ and $\mathbf{P}_* \text{Ker } \lambda_E$, where $G(2, H^0(E))$ is the Grassmannian of 2-dimensional subspaces of $H^0(E)$ embedded into $\mathbf{P}_* \wedge^2 H^0(E)$ by the Plücker coordinates. If a subspace $[U] \in G(2, H^0(E))$ belongs to $\mathbf{P}_* \text{Ker } \lambda_E$, then the evaluation homomorphism $ev^U : U \otimes \mathcal{O}_C \rightarrow E$ is not injective and its kernel is a line bundle. Moreover, the inverse of $\text{Ker } ev^U$ belongs to $W_5^1(C)$, and if C is general, then the map

$$G(2, H^0(E)) \cap \mathbf{P}_* \text{Ker } \lambda_E \longrightarrow W_5^1(C), \quad [U] \mapsto (\text{Ker } ev^U)^{-1} \quad (4.15)$$

is an isomorphism between two reduced 0-dimensional schemes, which shows Theorem 2.9 in the case $(a, b) = (2, 4)$. This idea leads us to a computation-free proof of Theorem 2.9, which we will discuss elsewhere.

References

- [1] Arbarello, E., Cornalba, M., Griffiths, P.A. and J. Harris: *Geometry of Algebraic Curves I*, Springer-Verlag, 1985.
- [2] Beauville, A.: Variétés de Prym et jacobiniennes intermédiaires, *Ann. Sci. Ec. Norm Sup.* (4), **10** (1977), 309-391.
- [3] Donaldson, S.K.: A new proof of a theorem of Narasimhan and Seshadri, *J. Diff. Geom.* **18** (1983), 269-2277.
- [4] Gieseker, D.: On the moduli of vector bundles on an algebraic surface, *Ann. of Math.* **106** (1977), 45-60.
- [5] ——— : Stable curves and special divisors: Petri's conjecture, *Invent. Math.*, **66** (1982), 251-275.
- [6] Lazarsfeld, R.: Brill-Noether-Petri without degenerations, *J. Diff. Geometry* **23** (1986), 299-307.
- [7] ——— : A sampling of vector bundle techniques in the study of linear system, in '*Lectures on Riemann surfaces*', World Scientific, Singapore, 1989, pp. 500-559.
- [8] Mukai, S.: Fano 3-folds, in '*Complex Projective Geometry*', London Math. Soc. Lect. Note Ser. 179, Cambridge University Press, 1992, pp. 255-263.
- [9] ——— : Curves and symmetric spaces, *Proc. Japan Acad.* **68**(1992), 7-10.
- [10] ——— : Curves and Grassmannians, to appear in '*Algebraic Geometry and Related Topics*', the proceeding of a symposium held at Inha Univ., Inchon, Korea, 1992.
- [11] Mumford, D.: Projective invariants of projective structures and applications, *Int'l. Cong. Math. Stockholm*, 1962, pp. 526-530.
- [12] ——— : Theta characteristic of an algebraic curve, *Ann. Sci. Ec. Norm Sup.* (4), **4** (1971), 181-192.

- [13] — : Curves and their Jacobians, The University of Michigan Press, 1975.
- [14] — and J. Fogarty: *Geometric Invariant Theory*, second enlarged edition, Springer-Verlag, 1982.
- [15] — and P.E. Newstead: Periods of a moduli space of bundles on a curve, *Amer. J. Math.* **90** (1968), 1201-1208.
- [16] Narasimhan, M.S. and C.S. Seshadri: Stable and unitary vector bundles on a compact Riemann surface, *Ann. of Math.* **82**(1965), 540-564.
- [17] Newstead, P.E.: Stable bundles of rank 2 and odd degree over a curve of genus 2, *Topology* **7**(1968), 205-215.
- [18] — : Brill-Noether Problems List Update, University of Liverpool, 1992.
- [19] Ramanan, S.: The moduli space of vector bundles over an algebraic curve, *Math. Ann.* **200** (1973), 69-84.
- [20] Teixidor, M.: Brill-Noether theory for stable vector bundles, *Duke Math. J.*, **62**, (1991), 385-400.

Department of Mathematics
School of Science
Nagoya University
464-01 Furō-chō, Chikusa-ku
Nagoya, Japan

