



INTERNATIONAL ATOMIC ENERGY AGENCY
UNITED NATIONS EDUCATIONAL, SCIENTIFIC AND CULTURAL ORGANIZATION
INTERNATIONAL CENTRE FOR THEORETICAL PHYSICS
I.C.T.P., P.O. BOX 586, 34100 TRIESTE, ITALY, CABLE: CENTRATOM TRIESTE



SMR.853 - 49

ANTONIO BORSELLINO COLLEGE ON NEUROPHYSICS

(15 May - 9 June 1995)

**"Grasping Objects: The Cortical Mechanisms
of Visuomotor Transformation"**

Giacomo Rizzolatti
Istituto di Fisiologia Umana
Universita' degli Studi di Parma
43100 Parma
Italy

**These are preliminary lecture notes, intended only for distribution to
participants.**

Grasping Objects: The Cortical Mechanisms of Visuomotor Transformation.

M. Jeannerod, M.A. Arbib, G. Rizzolatti, H. Sakata*

* M. Jeannerod is at Vision et Motricite, INSERM U94, 69500 Bron, France. M.A. Arbib is at the Center for Neural Engineering, University of Southern California, Los Angeles CA 90089-2520. G. Rizzolatti is at the Istituto di Fisiologia Umana, Universita di Parma, 43100 Parma, Italy. H. Sakata is at the Department of Physiology, Nihon University, Oyaguchi Itabashi, Tokyo, Japan.

This article stemmed from a collaboration between the authors within the framework of a Research Grant awarded by the Human Frontier Science Program (Principal Investigator, MJ).

Running title: **Grasping movements**

Key words: Grasping movements, Object-oriented behavior, Motor control, Prefrontal cortex, Parietal cortex, Visuomotor transformation, Schema theory,

Summary

Grasping requires coding of the object intrinsic properties (size and shape) and the transformation of these properties into a pattern of distal (finger, wrist) movements. Computational models address this behavior through the interaction of perceptual and motor schemas. In monkeys, the transformation of object intrinsic properties into specific grips takes place in a circuit formed by the inferior parietal lobule (area AIP) and the inferior premotor area (area F5). Neurons in both these areas code size, shape and orientation of objects and specific types of grip necessary to grasp them. Grasping movement are coded more globally in area AIP, whereas they are more segmented in F5. In humans, neuropsychological studies of parietal patients confirm that object primitives for grasping are analyzed in the parietal lobe, and also demonstrate that the "pragmatic" analysis of objects is separated from the "semantic" one performed in the temporal lobe.

In our everyday life we interact continually with objects. We reach for them, we grasp them, we manipulate them. All these actions are apparently very simple. Yet, this is not so. The mechanisms underlying them are complex and require multiple visuomotor transformations. This article examines one of these object-oriented actions, grasping. We first conduct a functional analysis of hand-object interactions in terms of behavioral units. We then discuss anatomical and single neuron data on the cortical circuits that in the monkey play a central role in grasping. Finally, we review neuropsychological data on object encoding following brain damage and draw conclusions on the different modes of information processing involved in object-oriented behavior.

The hand as a grasping apparatus

Grasping is a highly evolved type of behavior, based on the specialization of the hand and its neural apparatus in primates and man. Precision grip with true opposition of the pulpal surfaces of the thumb and index finger is considered as the hallmark of dextrous hands. Film and motion analysis of grasping show that the motor configuration that is formed by the hand in contact with the object represents the end result of a motor sequence which starts well ahead of the action of grasping itself. The fingers begin to shape during transport of the hand. This process of preshaping first involves a progressive opening of the

grip with straightening of the fingers, followed by a closure of the grip until it matches object size¹ (Fig. 1 a, b, d). The point in time where grip size is the largest (maximum grip size) is a clearly identifiable landmark which occurs well before the fingers come in contact with the object^{2, 3, 4} (Fig 1d). The question of why grip aperture is larger than required by object size is still a matter of debate. The critical point is that amplitude of maximum grip size linearly covaries with object size⁵. Monkeys also perform a similar preshaping with extra-opening and anticipatory closure of the fingers⁶.

Arbib and his colleagues^{7,8} proposed an approach in which control programs combine perceptual and motor schemas to determine the interactions between the hand and the environment (see Insert for the basic notions of Schema Theory). For the purpose of prehension, schemas like "preshape", "enclose", "orient", etc., are postulated. These schemas are carried out by specific grasping units (the "virtual fingers"). In a precision grip with pad opposition⁹, for example, the thumb is one virtual finger (VF1), the finger(s) that oppose the thumb is another one (VF2), and the unused finger(s) a third one (VF3). In grasping a small object, VF2 will be composed of the index finger only. In whole hand prehension with palm opposition, VF1 will be the palm and VF2 will (usually) include the four fingers other than the thumb.

The role of vision in grasping is not only to activate the proper schemas and specify the composition of the virtual fingers, but also to determine the relative positions of the hand and the object to be grasped, so that the forces during the lift of the object can be applied in exactly opposite directions. Accurate positioning of the fingerpads on the object surface is a prerequisite for subsequent handling and manipulation. This requires defining an "opposition space" corresponding to the grasp axis embedded in the object¹⁰. Then, the hand will be transported (the "approach" schema) and the wrist will rotate it (the "rotate" schema) in order to approximate the correct position. The formation of the grasp prior to contact with the object is thus the critical factor that governs the movements of the other segments of the upper limb during the reach. Although reach and grasp can be described as separate subsystems¹¹, studies of reaching in isolation from grasping ignore many of the key aspects of its control. The kinematic redundancy of the whole limb, and not only its distal segments, is exploited in building the appropriate opposition space¹².

Coactivation of reach and grasp raises the question of how schemas actuated by different limb segments are temporally coordinated by the program. The first coordinated control program for reach and grasp⁷ postulated that completion of the activity for grasping an object involved two motor schemas,

one for the slow phase of the reach and the second for the enclose phase of the hand movement. However, subsequent experiments showed this model to be inadequate. Paulignan et al¹³ suddenly displaced the target-object at the onset of a reach-to-grasp movement. In this condition, the untrained subject is perfectly able to correct for this visual "perturbation" and to accurately grasp the displaced object. This correction, however, results in prolonging the duration of the reach by about 100 ms. Meanwhile, the opening of the grip is interrupted, grip size decreases and increases again until it reaches its peak aperture at a later time, when the hand gets close to the displaced object (Figure 1c, e). To address these data, Hoff & Arbib¹⁴ proposed a model with a two-way interaction between the transport and grasp schemas. They postulated the existence of an additional, coordinating schema which receives from each of the constituent schemas an estimate of the time it needs to move from its current state to the desired final state. Whichever schema is going to take longer (in this case, the reach) is given the full time it needs, while the others will be slowed down (Figure 2). The time needed by each schema is regulated by optimality criteria which are embedded in feedback controllers which respond to disturbances with some latency.

The schema hypothesis provides a framework for segmenting grasping into elementary action units and for relating these units to the neural substrate. It also explains how grasping interacts with other functions of the upper limb, such as reaching. Ultimately, as will be shown in the last section, it may help in understanding pathological disorders of grasping.

Neural mechanisms involved in the control of visually guided grasping.

Correct execution of grasping requires the integrity of primary motor cortex (Brodmann area 4 or Field F1¹⁵). Lesion of this area in primates, as well as damage to the pyramidal tract, produces a profound deficit in the control of individual fingers and, as a consequence, a disruption of normal grasping^{16, 17}. Direct access of visual information needed for hand shaping, however, is very limited in F1, where visually responsive neurons are rare and have visual properties (brisk, transient responses to abrupt stimulus presentation¹⁸) that hardly fit those one would expect for grip formation. The visuomotor transformations required for grasping movements, therefore, have to occur upstream in motor control, in areas more closely connected to the visual system. In this section, we will present evidence for the existence of a specialized visuomotor system encoding object primitives and generating the corresponding

hand configurations. This system, which involves frontal and parietal areas, is depicted in Figure 3.

Premotor area F5. F5 forms the rostral part of inferior area 6. Its connections with F1 are within the hand field of this area^{19, 20, 21}. Intracortical microstimulation and single neuron studies showed that F5 is specifically related to distal movements^{22, 23}.

Rizzolatti and his colleagues^{23, 24} recorded single neurons from F5 in behaving monkeys tested during object oriented motor actions. These experiments showed that most neurons located in the upper part of F5 (the arm field) are related to grasping and other object-related motor actions (holding, tearing, manipulating). "Grasping" neurons discharge in relation with finger and hand movements during the action of grasping an object. The temporal relation of this discharge with grip movements changes from neuron to neuron. Some fire during the last part of grasping, that is, during finger flexion. Others start firing with finger extension and continue during finger flexion. Others are activated in advance of finger movements and often cease discharging only when the object is grasped.

An important property of most F5 neurons is their selectivity for different types of hand prehension. Eighty five per cent of grasping neurons show selectivity for one of three basic types of grip: precision grip (the most represented type), finger prehension and whole hand prehension²³. There is specificity for different finger configurations, even within the same grip type. Thus, the prehension of a sphere, which requires the opposition of all fingers, is encoded by different neurons than the prehension of a cylinder, for which a palm opposition grip is used²⁴ (Fig 4a, b).

The schema approach can be used here to conceptualize these physiological data. In F5, the schemas are represented by populations of neurons which code different motor acts. Various types of schemas can be distinguished. Some define general categories of action, e.g., grasp, hold, tear. Others indicate how the objects are to be grasped, held, torn, etc. In this case, each schema specifies the effectors appropriate for the action; "index finger and thumb" (precision grip), "all fingers" (manipulation), "all fingers but the thumb" (palm opposition), etc. Finally a third group of schemas would be concerned with the temporal segmentation of the actions (the coordinating schemas). Thus, the motor schemas in F5 form a basic "vocabulary"²⁵ from which many dextrous movements may be constructed as coordinated control programs.

The presence in F5 of this vocabulary has some important implications. First, since information is concentrated in relatively few elements, the number of

variables to be controlled is much less than if the movements were described in terms of motoneurons or muscles. This solution for reducing the high number of degrees of freedom of hand movements comes close to that proposed theoretically with the virtual fingers. Second, the retrieval of the appropriate movement is simplified. Both for internally generated actions and for those emitted in response to an external stimulus, only one schema or a small ensemble of schemas have to be selected or coordinated. In particular, the retrieval of a movement in response to a visual object is reduced to the task of matching its size and orientation with the appropriate schema. Third, the presence of a vocabulary of motor schemas should greatly facilitate the learning of associations, including arbitrary associations between stimuli and schemas (e.g., "if red, grasp; if green, don't"). Lesion studies showed that motor association learning is markedly impaired after damage to rostral premotor areas F5 and F7²⁶.

How is the motor vocabulary of F5 addressed by external stimuli? The simplest way to examine this issue is to present different types of visual stimuli and to establish whether the neuron will fire in the absence of movements and, if yes, in response to which stimuli. By using this approach, visual responses were observed in about 20-30% of F5 neurons. Two types of responses can be distinguished. Neurons of the first class respond to presentation of graspable objects. Often, there is a relation between the type of prehension coded by the cell and the size of the stimulus effective in triggering the neurons. This is particularly clear for the precision grip neurons which are activated only by small visual objects. Neurons of the second class ("mirror neurons") respond when the monkey sees movements, similar to those coded by the neuron, but executed by the experimenter or another monkey. For example, many mirror neurons fire when the monkey grasps a piece of food and also when the experimenter or the other monkey does so. They do not fire, however, when the experimenter makes a grasping movement without food, or when the food is grasped with a tool, etc.²⁷ (Fig 4c). Thus, the vocabulary of F5 can be addressed in two ways: by objects and by events. In both cases the eliciting stimuli address specifically the F5 neurons coding the grip congruent with them.

Parietal areas. Parietal cortex is known to be concerned with the visual control of hand movement from the effects of posterior parietal lesions in man (see below) and animals. Monkeys with lesions in the inferior parietal lobule typically present misreaching with the contralesional arm²⁸. In addition, their contralesional hand fails to shape and makes awkward grasps²⁹.

Neurons related to active arm movements were first recorded in the inferior parietal lobe by Mountcastle and his colleagues. They classified those neurons into two classes: "arm projection" and "hand manipulation" neurons³⁰. More recently, hand movement-related neurons were found to be concentrated in a small zone within the rostral part of the posterior bank of intraparietal sulcus, corresponding to an area (AIP) closely connected with area F5 of the premotor cortex³¹ (Fig 3). Neurons from this area were recorded in monkeys trained to manipulate various types of switches (some are shown in Fig 5), which elicited from the animal different motor configurations of the hand. Most of them were selectively activated during grasping one or two of these objects among the four routinely used ones. Neuron activity was not influenced by changing the position of the object in space, which shows that they were related to distal hand and finger movements rather than to proximal movements of the arm.

In order to determine the role of visual factors in activating these neurons, Sakata and his colleagues let the monkey perform the same task in the dark, guided only by a small spot of light on the object^{32, 33}. Thus, the task-related neurons were classified into three groups according to the difference between the activity in the light and in the dark: "motor dominant" neurons (Fig 5a) did not show any significant difference in activity between these two conditions; "visual and motor" neurons (Fig 5b, c) were less active in the dark; "visual dominant" neurons (Fig 5d) were exclusively active in the light.

Many of these visually responsive neurons were activated by the sight of objects during fixation without grasping ("object" type, Fig 5b, d). In most "visual and motor" neurons the visually effective object and the type of grip coded (assessed in the dark) coincided. Another type of neuron was not activated during the fixation of objects ("non-object" type, Fig 5C) but seemed to require other visual stimuli, such as the view of the moving hand to be activated. "Non-object" type responses were usually elicited after the initiation of the hand movement, and were likely to be concerned with the interaction of the hand with the object. Finally, using a broader variety of graspable objects, including primitive shapes such as spheres, cubes, cones, cylinders, rings and plates of different sizes, Sakata et al.³³ found many neurons in the rostral IPS posterior bank (area AIP) which were selectively activated during grasping or fixation of one or two of these objects. Some of them were also sensitive to the size or the orientation of the objects.

"Visual and motor" and "visual dominant" neurons in this region are likely to be interconnected. In addition, a group of neurons sensitive to the 3-D orientation of the longitudinal axis of visual stimuli were recently found in the

caudal part of the IPS posterior bank³⁴. It is therefore likely that the 3-D characteristics of the object are processed outside AIP and the output of such processing is sent to AIP. Thus, the parietal visual neurons encode the 3-D features of objects in a way that is suitable to guide the movements for grasping them. They may be regarded as a neural implementation of perceptual schemas.

If one compares the properties of parietal neurons with those of F5, striking similarities, but also important differences emerge. Visual responses to 3-D objects are found more frequently in parietal cortex than in F5. By contrast, "mirror neurons" responding to the view of hand action of other individuals were not found in AIP. As for the motor properties, parietal "motor dominant" neurons also code elementary motor acts, such as precision grip, whole hand prehension, wrist rotation, etc. However, most of the parietal neurons appear to represent the entire action, since they start to discharge with the hand shaping and continue to fire while the monkey is holding the object. This property contrasts with those of F5 neurons, which were commonly related to a particular segment of the action. Indeed, in primary motor cortex, on which F5 heavily projects, neurons code even more fragmental movements³⁵.

Transient inactivation of AIP, by injecting a GABA agonist (muscimol) in the rostral IPS posterior bank, produces a subtle change in the performance of visually guided movements during grasping tasks. In some cases, grasping errors are observed only in difficult tasks requiring a precision grip or during sticking out the index finger to insert it in a groove. Lack of preshaping can be observed as well during easier tasks such as grasping a small cube or sphere. In addition, there is a clearcut dissociation of the muscimol effects on grasping and reaching³⁶. Whereas the alteration of preshaping is consistently obtained after injection in the rostral part of the posterior bank of the sulcus, misreaching occurs after injection within its more caudal part. These results support the view that the parietal neurons related to manipulation play a specific role in the visual guidance of grasping.

A neuropsychological perspective on grasping movements.

The foregoing data can now be integrated within a broader framework concerning the way object-related visual information is processed. A highly influential conception³⁷ attributed different modes of visual processing to diverging corticocortical pathways. One of these pathways, the ventral route, links striate cortex to prestriate areas and inferotemporal cortex. Its interruption abolishes object discrimination without affecting perception of spatial relations

between objects. The other pathway, the dorsal route, diverges from the previous one by linking the prestriate areas to the posterior part of the parietal lobe. Its interruption produces visual spatial disorientation.

Recent observations, however, have prompted a reappraisal of the respective functions of the two cortical pathways. Posterior parietal cortex would be critical for organizing object-oriented action. This role would be complementary with that of occipitotemporal structures specialized for object identification and recognition³⁸. This view can be dramatically illustrated by two clinical cases. The first case is that of DF, a 35 year old woman, observed by Goodale et al³⁹. Following a bilateral lesion of occipito-temporal cortex (the ventral route), DF was unable to recognize objects. She was also unable to demonstrate with her fingers the size of visually inspected objects. In contrast, when instructed to take objects by performing prehension movements, she was quite accurate and her maximum grip size correlated normally with object size. An opposite picture is provided by the second case, that of AT, also a 35 year old woman, with a lesion of the occipitoparietal region, which was likely to have interrupted the dorsal route of visual processing⁴⁰. Whereas object recognition was normal, as was also the action of demonstrating object size with the fingers, preshape of the hand during object directed movements was incorrect. Correlation between object size and maximum grip size was lacking, with the consequence that objects could not be grasped between the fingertips: instead, the patient made awkward palmar grasps^{40, 41}. The schema framework offers a compelling explanation for this deficit. Because the grasp schemas were destroyed by the lesion or disconnected from visual input, the grip aperture did not stop at the required size, grip closure was delayed and the transport was prolonged in order to remain coordinated with the grasp.

This double dissociation suggests that object attributes are processed differently according to the task in which a subject is involved. To serve object-oriented action, these attributes (spatial as well as intrinsic) are subjected to a "pragmatic" mode of processing, the function of which is to extract parameters relevant to action and to generate the corresponding motor commands. During identification, another, "semantic" mode operates, through which object attributes are bound together to produce a unique percept. Although these modes of processing correspond to widely different cognitive functions^{42, 43}, they can be integrated and the semantic system can influence the pragmatic system. AT cannot preshape for size neutral objects like plastic cylinders, yet, when faced with a familiar object whose size is a semantic property, like a lipstick, she can

grasp it with reasonable accuracy⁴⁰. This interaction reflects the role of the abundant anatomical interconnections between the two cortical systems⁴⁴.

References

- 1 Jeannerod, M. (1981) in *Attention and Performance IX* (J. Long and A. Baddeley Eds), Erlbaum, Hillsdale, 153-168
- 2 Jeannerod, M. (1984) *J. Motor Behav.* 16, 235-254
- 3 Wing, A.M., Turton, A. and Fraser, C. (1986) *J. Motor Behav.* 18, 245-260
- 4 Gentilucci, M., Castiello, U., Corradini, M.L. et al (1991) *Neuropsychologia* 29, 361-378
- 5 Marteniuk, R.G., Leavitt, J.L., MacKenzie et al (1990) *Hum. Movement Sci.*, 9, 149-176
- 6 Gentilucci, M., Fogassi, L., Gallese, V. et al (in preparation)
- 7 Arbib, M.A. (1981) in *Handbook of Physiology* Sect. 1, Vol. 2, V.B. Brooks (Ed) Am. Physiol. Soc. Bethesda, Maryland, 1449-1480
- 8 Arbib, M.A., Iberall, T. and Lyons, D. (1985) in: *Hand function and the neocortex* A.W. Goodwin and I Darian-Smith (Eds), Springer, Berlin, 111
- 9 Napier, J.R. (1960) *Proc. Zool. Soc. London* 134, 647-657
- 10 MacKenzie, C.L. and Iberall, T. (1994) *The grasping hand* North Holland, Amsterdam
- 11 Jeannerod, M. (1988) *The Neural and Behavioral Organization of Goal-directed Movements* Clarendon, Oxford
- 12 Stelmach, G.E., Castiello, U. and Jeannerod, M. (1994) *J. Motor Behav.* 26, 178-186
- 13 Paulignan, Y., MacKenzie, C.L., Marteniuk, R. et al (1991) *Exp. Brain Res. X*
- 14 Hoff, B. and Arbib, M.A. (1993) *J. Motor Behav.* 25, 175-192
- 15 Matelli, M., Luppino, G. and Rizzolatti, G. (1985) *Behav. Brain Res.* 18, 125-137
- 16 Fulton, J.F. (1949) *Physiology of the nervous system* Oxford University Press, New-York
- 17 Passingham, R., Perry, H. and Wilkinson, F. (1978) *Brain Res.* 145, 410-414
- 18 Wannier, T.M.J., Maier, M.A. and Hepp-Reymond, M.C. (1989) *Neurosci Lett.* 98, 63-68.
- 19 Muakkassa, K.F. and Strick, P.L. (1979) *Brain Res.* 177, 176-182.
- 20 Matsumura, M. and Kubota, K. (1979) *Neurosci. Let.* 11, 241-246.
- 21 Matelli, M. Camarda, R., Glickstein, M. et al (1986) *J. Comp. Neurol.* 251, 281-298.
- 22 Kurata, K. and J. Tanji (1986) *J. Neurosci.* 6, 403-411.

23 Rizzolatti, G., Camarda, R., Fogassi, M. et al (1988) *Exp. Brain Res.* 71, 491-507

24 Gallese, V., di Pellegrino, G., Fadiga, L. et al (1992) *ENA Satellite Symposium.*, Ohlstadt.

25 Rizzolatti, G. and Gentilucci, M. (1988) in *Neurobiology of Neocortex* P. Rakic and W. Singer, (Eds) Wiley, Chichester, pp 269-284.

26 Passingham, R. (1993) *The frontal lobe and voluntary action* Oxford University press, Oxford.

27 di Pellegrino, G., Fadiga, L., Fogassi, L. et al (1992) *Exp. Brain Res.* 91, 176-180.

28 Haaxma, H. and Kuypers, H.G.J.M. (1975) *Brain* 98, 239-260

29 Faugier-Grimaud, S., Frenois, C. and Stein, D.G. (1978) *Neuropsychologia* 16, 151-168

30 Mountcastle, V.B., Lynch, J.C., Georgopoulos, A.P. et al (1975) *J. Neurophysiol.* 38, 871-908

31 Taira, M., Mine, S., Georgopoulos, A.P. et al (1990) *Exp. Brain Res.* 83, 29-36

32 Sakata, H., Taira, M., Mine, S. et al (1992) *Exp. Brain Res. suppl.* 22, 185

33 Sakata, H., Murata, A., Luppino, G. et al (1992) *Soc. Neurosci. Abstr.* 18, 504

34 Kusunoki, M., Tanaka, Y., Ohtsuka, H. et al (1993) *Soc. Neurosci. Abstr.* 19, 770.

35 Lemon, R.N., Mantel, G.W.H. and Muir, R.B. (1986) *J. Physiol.* 318, 497-527

36 Gallese, V., Murata, A., Kaseda, M. et al (1994) *Neuroreport* 5, 1525-1529

37 Ungerleider, L.G., Mishkin, M. (1982) in *The analysis of visual behavior* D.J. Ingle, M.A. Goodale and R.J.W. Mansfield (Eds) MIT Press, Cambridge, pp 549-586

38 Goodale, M. A. and Milner, A.D. (1992) *TINS* 15, 20-25

39 Goodale, M. A., Milner, A.D., Jakobson, L.S. et al (1991) *Nature* 349, 154-156

40 Jeannerod, M., Decety, J. and Michel, F. (1994) *Neuropsychologia* 32, 369-380

41 Jeannerod, M. (1986) *Behav. Brain Res.* 19, 99-116

42 Castiello, U., Paulignan, Y. and Jeannerod, M. (1991) *Brain* 114, 2639-2655

43 Castiello, U. and Jeannerod, M. (1994) *Neuroreport* 2, 797-800.

44 Morel, A. and Bullier, J. (1990) *Visual neurosci.* 4, 555-578.

45 Arbib, M.A. and Liaw, J.S. (1995) *Artificial Intelligence* 61, in press

Figure Legends

Figure 1.

Kinematics of grasping. A. The hand preshapes during transport to target-object (superimposed views of a single movement at a rate of 5 Hz.) B. Averaged 2-D recording of spatial paths of wrist (green), tip of thumb (red) and tip of index finger (blue) during 10 reach-to-grasp movements. C. Averaged spatial paths when a perturbation occurred (object was briskly displaced rightward at onset of movement). Small horizontal and vertical bars on tracings represent the values of one standard deviation from mean trajectory. D and E. Time (in milliseconds) plots of wrist velocity (green line, scale on right in cm/s) and grip size (distance between fingertips, scale on left in mm, red line) during one of the trials represented in B and C, respectively. Data from Ref 13.

Figure 2.

An overview of a model of the motor schemas, and their coordination through timing, for reaching and grasping¹⁴. dashed lines carry activation signals to a schema, as in a computer flow diagram, while solid lines carry signals coding relevant variables, as in a block diagram for a control system. Thus, the entire diagram is an example of a schema defined as a coordinated control program of finer schemas. The overall schema combines three motor schemas (rectangles) and one coordinating schema (with rounded corners) Whichever motor schema needs more time (arm: Transport, or hand: Preshape + Enclose) sets the total duration specified by The Time-Based Coordination model. The schemas then provide the optimal arm and hand trajectories for the specified duration. Parameters of the movement may be adjusted on-line (but with about 100 ms delay) in response to unexpected perturbations.

Figure 3.

Lateral and mesial views of monkey cerebral cortex. Visuomotor stream for grasping is indicated by large arrows. F5 also receives somatosensory input from area SII and somatosensory and visual input from area 7b (circled areas). Frontal agranular cortical areas are classified according to Matelli et al¹⁵. Cortical areas controlling grasping are connected with basal ganglia and cerebellar circuits. These circuits, although involved in grasping, are not shown on the figure. Abbreviations: AIP, Anterior intraparietal area; AIs, Inferior arcuate sulcus; ASs, superior arcuate sulcus; Cs, central sulcus; Cgs,

cingulate sulcus; IPs, intraparietal sulcus; LIP, lateral intraparietal area; Ls, lateral sulcus; MIP, medial intraparietal area; Ps, principal sulcus; STs, superior temporal sulcus; VIP, ventral intraparietal area. Note that IPs and Ls have been opened to show hidden areas.

Figure 4.

Examples of F5 neurons. A and B: grasping neurons. The monkey is seated in front of a dark box. The trial started when the monkey pressed a bar. The box was then illuminated and a geometric solid, located inside it, became visible. After a variable time interval the door of the box opened automatically and the monkey was allowed to release the bar and reach for the object. Time plots and of neuronal discharge (rasters and histograms) and the distance between the thumb and index finger (recorded with a computerized movement analyzer) are presented. The traces are aligned with the onset of hand movement (vertical bar). Open circles indicate the moment when the door opened. The presented objects were (from left to right) a small sphere, a large sphere, and a horizontally positioned cylinder. Abscissae: time; ordinates: spikes/bin; binwidth: 20 ms. C: grasping neuron with mirror properties. First two panels: an experimenter grasps a raisin in front of the monkey (first discharge), moves it towards the monkey (no discharge), the monkey grasps it (second discharge). Note the difference between hand and tool grasping (pliers). Right panel: same neuron. Monkey grasps an object in the dark.

Figure 5.

Types of hand manipulation related neurons in monkey anterior intraparietal area (AIP). Activity of cells during hand manipulation in light and in dark, as well as during visual fixation of objects is shown with rasters and histograms. A: "Motor dominant" neuron that preferred open pull knob. The cell was equally active during manipulation in light and in dark, but was not active during object fixation. B: An object-type of "visual and motor" neuron that preferred push button. The cell was less active during manipulation in dark than in light, and was partly activated during object fixation. C: A non-object type of "visual and motor" neuron that preferred pull knob in groove. The cell was less active during manipulation in dark, but was not activated during object fixation. D: "Visual dominant" neuron that preferred upright pull lever. The cell was not activated during hand movement in dark but was fully activated during fixation of the object in light. At bottom of figure, Key indicates the period of pressing the anchor key before moving to the object. Obj indicates the period of holding the object to keep the switch on. Data from Ref 31.

INSERT.

Neuroscience has a well-established terminology for levels of structural analysis (e.g., brain area, layer, column...) but pays little attention to the need for a functional terminology. Schema theory⁷ provides a rigorous analysis of behavior which requires no prior commitment to hypotheses on neural localization. Schemas are units for this analysis. Perceptual schemas serve perceptual encoding, while motor schemas provide control units for movement. Crucially, schemas can be combined to form coordinated control programs which control the phasing in and out of patterns of schema coactivation and the passing of control parameters from perceptual to motor schemas. The notion of schema is recursive - a schema may later be analyzed as a coordinated control program of finer schemas, and so on until such time as a secure foundation of neural localization is attained.

The activity level of an instance of a perceptual schema represents a "confidence level" that the object represented by the schema is indeed present; while that of a motor schema may signal its "degree of readiness" to control a part of an action. Mutually consistent schema instances are strengthened and reach high activity levels to constitute the overall solution of a problem, whereas instances which do not reach the evolving consensus lose activity, and thus are not part of this solution. A corollary to this view is that the instances related to a given object oriented action are distributed. A given schema, defined functionally, may be distributed across more than one brain region; conversely, a given brain region may be involved in many schemas. Hypotheses about localization of schemas in the brain may be tested by observation of the effects of lesions or functional imaging, and a given brain region can then be modelled by seeing if its known neural circuitry can indeed be shown to implement the posited schemas. An example of this approach is given here.

In providing an account of the development (or evolution) of schemas, we find that new schemas often arise as modulators of existing schemas rather than as new systems with independent functional roles⁴⁵. Thus, schemas for control of dextrous hand movements serve to modulate less specific schemas for reaching with an undifferentiated grasp and to adapt them to the shape or the use of an object.

M.A. Arbib




Fig. 1




Fig. 2





Fig. 3



Grasping neuron with mirror property

experimenter grasps (hand) - gives - monkey grasps (tool) - experimenter grasps (tool) - gives - monkey grasps in the dark




Fig. 5