

the  
**abduS salam**  
international centre for theoretical physics

SMR.1508 - 30

*SUMMER SCHOOL ON PARTICLE PHYSICS*

16 June - 4 July 2003

LECTURES ON SUPERSYMMETRY

Lecture IV & V

G. WEIGLEIN  
Dept. of Physics. IPPP  
University of Durham  
Durham  
U.K.



# Lectures on Supersymmetry

*Georg Weiglein*

*IPPP Durham*

Trieste, 06/2003

Lectures 4 and 5:

- Higgs mass bound in SUSY theories
- The LSP as a dark matter candidate
- Electroweak precision tests: SM vs. MSSM
- Prospects for SUSY searches at the next generation of colliders

## Higgs mass bounds in SUSY theories

MSSM predicts upper bound on  $m_h$ :

tree-level bound:  $m_h < M_Z$ , excluded by LEP Higgs searches!

Large radiative corrections:

Yukawa couplings:  $\frac{e m_t}{2M_W s_W}$ ,  $\frac{e m_t^2}{M_W s_W}$ ,  $\dots$

$\Rightarrow$  Dominant one-loop corrections:  $G_\mu m_t^4 \ln \left( \frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2} \right)$

Present status of  $m_h$  prediction in the MSSM:

Complete one-loop and 'almost complete' two-loop result available

## Upper bound on $m_h$ in the MSSM:

“Unconstrained MSSM”:

$M_A$ ,  $\tan \beta$ , 5 parameters in  $\tilde{t}$ - $\tilde{b}$  sector,  $\mu$ ,  $m_{\tilde{g}}$ ,  $M_2$

Diagrammatic result: *FeynHiggs*

[*S. Heinemeyer, W. Hollik, G. W. '98, '00, '02*]

<http://www.feynhiggs.de>


$$m_h \lesssim 135 \text{ GeV}$$

for  $m_t = 175 \text{ GeV}$

no theoretical uncertainties from unknown higher orders included

Upper bound on  $m_h$  saturated for large  $M_A$ , large  $\tan\beta$ , significant mixing in  $\tilde{t}$  sector

[S. Heinemeyer, W. Hollik, G. W. '01]



Upper bound  $m_h \lesssim 135$  GeV reduced by

$\approx 6, 11, 8$  GeV in **mSUGRA**, **GMSB**, **AMSB** scenarios

[*S. Ambrosanio, A. Dedes, S. Heinemeyer, S. Su, G. W. '01*]

Upper bound on  $m_h$  in extensions of MSSM:  $m_h \lesssim 200$  GeV

[*G. Kane, C. Kolda, J. Wells '93*] [*J. Espinosa, M. Quirós '93, '98*]

**$\Rightarrow$  SUSY requires light Higgs boson:**

definite and robust prediction of SUSY models

testable at next generation of colliders

## Remaining theoretical uncertainties in prediction for $m_h$ in the MSSM:

[G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich, G. W. '02]

– From unknown higher-order corrections:

$$\Rightarrow \Delta m_h \approx 3 \text{ GeV}$$

– From uncertainties in input parameters

$$m_t, \dots, M_A, \tan \beta, m_{\tilde{t}_1}, m_{\tilde{t}_2}, \theta_{\tilde{t}}, m_{\tilde{g}}, \dots$$

$$\Delta m_t \approx 5 \text{ GeV} \Rightarrow \Delta m_h \approx 5 \text{ GeV}$$

## Higgs couplings, production cross sections

Also affected by large SUSY loop corrections


FD two-loop corrections implemented in results for  $h \rightarrow f\bar{f}$ ,  $e^+e^- \rightarrow hZ, hA$

[S. Heinemeyer, W. Hollik, G. W. '00]

[S. Heinemeyer, W. Hollik, J. Rosiek, G. W. '01]



$hf\bar{f}$  coupling:



$$A(h \rightarrow f\bar{f}) = \sqrt{Z_h} \left( \Gamma_h - \frac{\hat{\Sigma}_{hH}(m_h^2)}{m_h^2 - m_H^2 + \hat{\Sigma}_{HH}(m_h^2)} \Gamma_H \right)$$

$\Rightarrow$  Effective  $hf\bar{f}$  coupling can vanish for large  $\hat{\Sigma}_{hH}$

Glauino vertex corrections to  $h \rightarrow q\bar{q}$ :

$\Rightarrow$  ratio  $\Gamma(h \rightarrow \tau^+\tau^-)/\Gamma(h \rightarrow b\bar{b})$  can significantly differ from SM value for large  $\tan\beta$


Effective  $hf\bar{f}$  coupling can go to zero for large  $\hat{\Sigma}_{hH}$

⇒ “Pathological regions”

[W. Loinaz, J. Wells '98] [M. Carena, S. Mrenna, C. Wagner '99]

⇒ Suppression of  $BR(h \rightarrow b\bar{b})$ ,  
 $BR(h \rightarrow \tau\tau)$ , ...

[S. Heinemeyer, W. Hollik, G. W. '00]



## $\mathcal{CP}$ violation in the Higgs sector:

MSSM Higgs sector is  $\mathcal{CP}$ -conserving at tree level

Complex parameters enter via loop corrections:

- $\mu$ : Higgsino mass parameter
- $A_{t,b,\tau}$ : trilinear couplings  $\Rightarrow X_{t,b,\tau} = A_{t,b} - \mu^* \{\cot \beta, \tan \beta\}$  complex
- $M_{1,2}$ : gaugino mass parameter (one phase can be eliminated)
- $m_{\tilde{g}}$ : gluino mass

$\Rightarrow$  can induce  $\mathcal{CP}$ -violating effects

$\Rightarrow$  Mixing between neutral Higgs bosons  $h_1, h_2, h_3$

## Inclusion of higher-order corrections:

Propagator / mass matrix with higher-order corrections:

$$\begin{pmatrix} q^2 - M_A^2 + \hat{\Sigma}_{AA}(q^2) & \hat{\Sigma}_{AH}(q^2) & \hat{\Sigma}_{Ah}(q^2) \\ \hat{\Sigma}_{HA}(q^2) & q^2 - m_H^2 + \hat{\Sigma}_{HH}(q^2) & \hat{\Sigma}_{Hh}(q^2) \\ \hat{\Sigma}_{hA}(q^2) & \hat{\Sigma}_{hH}(q^2) & q^2 - m_h^2 + \hat{\Sigma}_{hh}(q^2) \end{pmatrix}$$


$\hat{\Sigma}_{ij}(q^2)$  ( $i, j = h, H, A$ ) : renormalized Higgs self-energies

$\hat{\Sigma}_{Ah}, \hat{\Sigma}_{AH} \neq 0 \Rightarrow \mathcal{CPV}$ ,  $\mathcal{CP}$ -even and  $\mathcal{CP}$ -odd fields can mix

Result:  $(A, H, h) \rightarrow (h_3, h_2, h_1)$  with  $m_{h_3} > m_{h_2} > m_{h_1}$

Example:  $\phi_{A_t}$  dependence of  $m_{h_1}, m_{h_2}$  ( $\phi_\mu = 0, |A_t| = 2 M_{\text{SUSY}}$ ):

[M. Frank, S. Heinemeyer, W. Hollik, G. W. '03]




$t/\tilde{t}, b/\tilde{b}, q^2 = 0$ , (s)fermion,  $q^2 = 0$ , full MSSM,  $q^2 = 0$ , full MSSM,  $q^2 \neq 0$

Complex phases can have large effects on Higgs couplings

Example:  $g_{hVV}$  for  $h_1, h_2, h_3$ :

[M. Frank, S. Heinemeyer, W. Hollik, G. W. '03]




full:  $h_1$ , dashed:  $h_2$ , dotted:  $h_3$


Parameters:  $M_{\text{SUSY}} = 500 \text{ GeV}$ ,  $M_2 = 500 \text{ GeV}$ ,  $\mu = 2000 \text{ GeV}$ ,  $|A_t| = 1000 \text{ GeV}$ ,  
 $M_{H^\pm} = 150 \text{ GeV}$ ,  $\tan \beta = 5$

Search for neutral MSSM Higgs bosons at LEP:

$$e^+e^- \rightarrow Zh, ZH$$




$$e^+e^- \rightarrow Ah, AH$$



Constraints from the Higgs search at LEP [*LEP Higgs Working Group '01*]

Experimental search vs. upper  $m_h$ -bound (*FeynHiggs* 1.0)

$m_h^{\max}$ -scenario ( $m_t = 174.3$  GeV,  $M_{\text{SUSY}} = 1$  TeV):



$m_h > 91.0$  GeV  
(expected: 94.6 GeV), 95% C.L.

$M_A > 91.9$  GeV  
(expected: 95.0 GeV)



Parameter region where experimental lower bound on  $m_h$  is significantly lower than SM bound,  $M_H > 114.4$  GeV, corresponds to  $\sin^2(\beta - \alpha_{\text{eff}}) \ll 1$

“Excluded”  $\tan \beta$  region:


$$0.5 < \tan \beta < 2.4$$

**Note:** this exclusion bound assumes

$m_t, M_{\text{SUSY}}$  fixed,  $m_t = 174.3$  GeV,  $M_{\text{SUSY}} = 1$  TeV

no theoretical uncertainties included

Effect of new corrections and  $m_t \rightarrow m_t + \sigma_{m_t}$ ,  $M_{\text{SUSY}} = 1 \text{ TeV} \rightarrow 2 \text{ TeV}$   
 [G. Degrandi, S. Heinemeyer, W. Hollik, P. Slavich, G. W. '02]




⇒ no  $\tan \beta$  exclusion for  $m_t \rightarrow m_t + \sigma_{m_t}$ ,  $M_{\text{SUSY}} = 1 \text{ TeV} \rightarrow 2 \text{ TeV}$

⇒ Low  $\tan \beta$  region not fully excluded by LEP!

# Experimental search vs. upper $m_h$ -bound (*FeynHiggs*)

[*LEP Higgs Working Group '01*]

no-mixing scenario:



## The LSP as a dark matter candidate

Astrophysical data (cosmic microwave background, ...)  $\Rightarrow$  existence of non-baryonic cold dark matter in the Universe

SUSY with R parity conservation  $\Rightarrow$  LSP relic density falls naturally in favoured range if  $m_{\text{LSP}} \lesssim 1 \text{ TeV}$

Most recent result (WMAP)

[C. Bennet et al. '03], [D. Spergel et al. '03]

$\Rightarrow$  cold dark matter density:  $\Omega_{\text{CDM}} h^2 = 0.1126^{+0.0161}_{-0.0181}$  at 95% C.L.  
( $h$ : expansion rate)



$\Rightarrow$  Constraints on the SUSY parameter space

Example: mSUGRA scenario with  $\mu > 0$ ,  $A_0 = 0$ ,  $\tan \beta = 10, 50$

Parameter region allowed by the CDM constraint in the  $m_{1/2}$ - $m_0$  plane with and without the new WMAP data

(also shown: favoured region from  $(g_\mu - 2)$ , region excluded by  $b \rightarrow s\gamma$ )

[J. Ellis, K. Olive, Y. Santoso, V. Spanos '03]




⇒ mSUGRA scenario consistent with all experimental constraints

## Direct search for dark matter:

E.g.: scattering off nuclei  $\Rightarrow$  measurement of nuclear recoil

Present limits in the plane of  $m_{\text{WIMP}}$  and the WIMP–nucleon cross section  $\sigma_{\text{SI}}$  from different experiments




⇒ Present limits on  $\sigma_{SI}$  (“Stage 1” dark matter experiments):  
 $\approx 10^{-5} - 10^{-6}$  pb

Anticipated sensitivity of possible future dark matter experiments  
 (“Stage 3”):  $\sigma_{SI} \approx 10^{-9} - 10^{-10}$  pb

Accessible parameter region (yellow) in mSUGRA scenario ( $\tan\beta = 30$ ,  
 $\mu > 0$ ,  $A_0 = 0$ ) with Stage 3 sensitivity vs. CDM allowed region and LHC  
 sensitivity

[H. Baer, C. Bal'azs,  
 A. Belyaev, J. O'Farrill '03]



## Electroweak precision tests: SM vs. MSSM

### Electroweak precision measurements:

$M_Z$ [GeV]	=	$91.1875 \pm 0.0021$	0.002%
$G_\mu$ [GeV <sup>-2</sup> ]	=	$1.16637(1) 10^{-5}$	0.0009%
$m_t$ [GeV]	=	$174.3 \pm 5.1$	2.9%
$M_W$ [GeV]	=	$80.426 \pm 0.034$	0.04%
$\sin^2 \theta_{\text{eff}}^{\text{lept}}$	=	$0.23148 \pm 0.00017$	0.07%
$\Gamma_Z$ [GeV]	=	$2.4952 \pm 0.0023$	0.09%
...			

Quantum effects of the theory: loop corrections:  $\sim \mathcal{O}(1\%)$

SM:  $M_H$  is free parameter

precise measurement of  $M_W$ ,  $\sin^2 \theta_{\text{eff}}$ , ...  $\Rightarrow$  constraints on  $M_H$

MSSM:  $m_h$  is predicted

precise meas. of  $M_W$ ,  $\sin^2 \theta_{\text{eff}}$ ,  $m_h$ , ...  $\Rightarrow$  constr. on  $m_{\tilde{t}}$ ,  $\theta_{\tilde{t}}$ ,  $m_{\tilde{b}}$ ,  $\theta_{\tilde{b}}$ , ...




## Comparison of ew precision data with theory:

EW precision data:  
 $M_Z, M_W, \sin^2 \theta_{\text{eff}}^{\text{lept}}, \dots$

Theory:  
SM, MSSM, ...



Test of theory at quantum level:



Improve indirect constraints on unknown parameters:  $M_H, m_{\tilde{t}}, \dots$

effects of “new physics”?

Indirect determination of  $m_t$  from precision data:

$$m_t = 180.3_{-9.2}^{+11.7} \text{ GeV}$$

Direct measurement:

$$m_t = 174.3 \pm 5.1 \text{ GeV}$$

Leading corrections to precision observables:

$$\begin{aligned} &\sim m_t^2 \\ &\sim \ln M_H \end{aligned}$$

⇒ Very high accuracy of measurements and theoretical predictions needed


Theoretical uncertainties:

- unknown higher-order corrections
- experimental error of input parameters:  $m_t$ ,  $\Delta\alpha_{\text{had}}$ , ...

Global fit of the SM to all data:

Basic assumption: SM provides correct description of experimental data

Comparison of SM prediction with the data:  
[LEPEWWG '03]



Overall fit probability (quality of the fit): 4.4%

Comparison of current experimental errors with anticipated precision at  
 Run II of the Tevatron ( $p\bar{p}$  collider,  $E_{\text{CM}} \approx 2$  TeV;  $\geq 2001$ ),  
 LHC ( $pp$  collider,  $E_{\text{CM}} \approx 14$  TeV;  $\gtrsim 2007$ ),  
 LC ( $e^+e^-$  collider,  $E_{\text{CM}} \approx 500\text{--}1000$  GeV,  $\gtrsim 201x?$ )  
 with and without low-energy running mode (GigaZ)

	now	Tev. Run IIA	Run IIB	LHC	LC	GigaZ
$\delta \sin^2 \theta_{\text{eff}} (\times 10^5)$	17	78	29	14–20	—	1.3
$\delta M_W$ [MeV]	34	27	16	15	10	7
$\delta m_t$ [GeV]	5.1	2.7	1.4	1.0	0.1	0.1
$\delta m_h$ [MeV]	—	—	$\mathcal{O}(2000)$	100	50	50

Additional sources for sizable radiative corrections in the MSSM:

- Mass and couplings of light  $\mathcal{CP}$ -even Higgs:

Large Yukawa corrections:  $\sim G_\mu m_t^4 \ln\left(\frac{m_{\tilde{t}_1} m_{\tilde{t}_2}}{m_t^2}\right), \dots$

- t /  $\tilde{t}$  loops, b /  $\tilde{b}$  loops (for large  $\tan\beta$ )
- Corr. to relation between bottom mass and bottom Yukawa coupling:

$$y_b = \frac{\sqrt{2}}{v \cos\beta} \frac{m_b}{1 + \Delta_b},$$

$$\Delta_b = \mu \tan\beta (\alpha_s I(\dots) + \alpha_t I(\dots))$$


$\Rightarrow$  Coupling non-perturbative for  $\Delta_b \rightarrow -1$

- Loop contributions from light SUSY particles
- $\dots$

Prediction for  $M_W$  in the **SM** and the **MSSM**:

[A. Djouadi, P. Gambino, S. Heinemeyer,  
W. Hollik, C. Jünger, G. W. '97]

[S. Heinemeyer, W. Hollik, G. W. '98]




**SM:**  $M_H$  varied

**MSSM:** SUSY parameters varied

# Prediction for $M_W$ in the SM and the MSSM:

[A. Djouadi, P. Gambino, S. Heinemeyer,  
W. Hollik, C. Jünger, G. W. '97]

[S. Heinemeyer, W. Hollik, G. W. '98]




SM:  $M_H$  varied


MSSM: SUSY parameters varied

Experimental result for anomalous magnetic moment of the muon vs. SM prediction:

[*Muon (g - 2) Collaboration '02*]



$\Rightarrow a_\mu(\text{exp}) - a_\mu(\text{SM}): 1-3 \sigma$  deviation (sizable hadronic uncertainties)



Compatible with effect of SUSY contributions to  $a_\mu$ :




mSUGRA predictions for  $(g_\mu - 2)$ :

[W. de Boer, C. Sander '02]

$$a_\mu = \frac{1}{2}(g_\mu - 2)$$

$$a_\mu - a_\mu^{\text{SM}} = (338 \pm 112) \times 10^{-11}$$



$\Rightarrow$  Preference for  $\mu > 0$

note: influenced by hadronic uncertainties of SM result

No convincing new physics model for “explaining” deviations from SM in  $\sin^2 \theta_{\text{eff}}$  and  $\nu$ -nucleon scattering from NuTeV

Note, however:

if NuTeV result interpreted as deviation in  $Z\nu\bar{\nu}$  coupling:

( $\nu$  neutral current rate)/prediction:

$0.995 \pm 0.003$     LEP1 lineshape (invisible Z width)

$0.988 \pm 0.004$     NuTeV

Need some patience until experimental clarification:

$A_{\text{LR}}, A_{\text{FB}}$ :            LC with GigaZ option?

$\nu$ -nucleon scatt.:         $\nu$  factory?


If hadronic data on  $A_{\text{FB}}$  and NuTeV data on  $\nu$ -nucleon scattering were removed from fit:

much better fit quality, probability  $\approx 70\%$

stronger preference for light Higgs:  $M_{\text{H}} \lesssim 150$  GeV, 95% C.L.

The effective weak mixing angle  $\sin^2 \theta_{\text{eff}}$ :

[LEPEWWG '03]




Prediction for  $M_W$ ,  $\sin^2 \theta_{\text{eff}}$  in **SM** and **MSSM**:

[S. Heinemeyer, G. W. '01]

Scenario where measurement  
of  $A_{\text{FB}}^b$  is discarded:

$$\sin^2 \theta_{\text{eff}}^{\text{SLD}} = 0.23098 \pm 0.00026$$

⇒ good agreement with  
MSSM prediction




⇒ High sensitivity to deviations both from the SM and the MSSM

# MSSM prediction for neutrino–nucleon scattering

(measured at NuTeV):




[*S. Davidson, S. Forte, P. Gambino,  
N. Rius, A. Strumia '02*]



⇒ Discrepancy cannot be explained in MSSM (light slepton corrections)

# Global fit: SM / unconstrained MSSM / mSUGRA


[W. de Boer, C. Sander '02]

	SM:	$\chi^2/\text{d.o.f} = 27.1/16$
	MSSM:	$\chi^2/\text{d.o.f} = 16.4/12$
	CMSSM:	$\chi^2/\text{d.o.f} = 23.2/17$

⇒ MSSM contributions can improve agreement with data for  $g_\mu - 2$  (for large  $\tan\beta$ ) and  $M_W$

no significant improvement for  $A_{\text{FB}}^b$  and neutrino–nucleon cross section

Similar quality of global fit (slightly better in MSSM than in SM)



## Prospects for SUSY searches at the next generation of colliders

Limits on SUSY particles from LEP, Tevatron Run I:  $\mathcal{O}(100 \text{ GeV})$

But: some SUSY particles can be much lighter if they have small coupling to Z boson

E.g.:

- no strict lower bound on lightest neutralino if “GUT relation” between  $M_1$  and  $M_2$  is relaxed
- A light sbottom with  $m_{\tilde{b}_1} \approx 5 \text{ GeV}$  is in agreement with electroweak precision tests, Higgs searches, ...

*[M. Carena, S. Heinemeyer, C. Wagner, G. W. '01]*

– ...

## SUSY searches at the Tevatron, Run II:

compared to Run I:  $\approx 100\times$  higher luminosity, slightly increased energy (1.8  $\rightarrow$  2 TeV)

Limited mass window in which discovery of SUSY particles above Run I is possible

Best prospects for:

- ‘Trilepton signal’:  $\tilde{\chi}_2^0 \tilde{\chi}_1^+ \rightarrow \ell^+ \ell^- \tilde{\chi}_1^0 \ell^+ \nu \chi_1^0$
- $\tilde{t}$ ,  $\tilde{b}$  searches
- light SUSY Higgs  $h$  in region of large  $\tan \beta$

## SUSY searches at the LHC:


Dominated by production of **colored** particles: **gluino, squarks**

Very large mass range in the searches for **jets + missing energy**

$\Rightarrow$  gluino, squarks accessible up to 2–3 TeV



Discovery reach contours in  $m_0$ - $m_{1/2}$  plane (mSUGRA scenario) for various final states with  $100 \text{ fb}^{-1}$ : [CMS '99]



⇒ discovery of SUSY particles expected if low-energy SUSY is realized

Production of SUSY particles at the LHC will in general result in complicated final states, e.g.

$$\tilde{g} \rightarrow \bar{q}\tilde{q} \rightarrow \bar{q}q\tilde{\chi}_2^0 \rightarrow \bar{q}q\tilde{\tau}\tau \rightarrow \bar{q}q\tau\tau\tilde{\chi}_1^0$$

Production of uncolored particles via cascade decays often dominates over direct production

Many states are produced at once



**⇒ Main background for SUSY is SUSY itself!**

Searches for MSSM Higgs bosons:

good prospects for detecting light Higgs  $h$

$H/A$  discovery possible in significant part of parameter space

MSSM Higgs discovery contours in  $M_A$ - $\tan\beta$  plane ( $m_h^{\max}$  benchmark scenario): [ATLAS '99] [CMS '03]



In order to establish SUSY experimentally:

Need to demonstrate that:

- every particle has superpartner
- their spins differ by  $1/2$
- their gauge quantum numbers are the same
- their couplings are identical
- mass relations hold

...

⇒ Precise measurements of masses, branching ratios, cross sections, angular distributions, ... mandatory for

- establishing SUSY experimentally
- disentangling patterns of SUSY breaking

Requires clean experimental environment, high luminosity, beam polarization, . . .

⇒ High luminosity LC necessary, complementary to hadron machines

SUSY searches at the LC:

Clean signatures, small backgrounds

**Thresholds** for pair production of SUSY particles

⇒ precise determination of mass and spin of SUSY particles, mixing angles, complex phases, . . .

Limited by kinematic reach

Good prospects for production of uncolored particles


⇒ LHC / LC complementarity

## Examples for SUSY physics at the LC:

Determination of mass and spin of  $\tilde{\mu}_R$  from production at threshold:

[TESLA TDR '01]


$$\Rightarrow \frac{\Delta m_{\tilde{\mu}_R}}{m_{\tilde{\mu}_R}} < 1 \times 10^{-3}$$



$\Rightarrow$  test of  $J = 0$  hypothesis

Determination of **phase  $\phi_1$**  in neutralino sector from measurement of  $e^+e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_2^0$  and  $e^+e^- \rightarrow \tilde{\chi}_1^+ \tilde{\chi}_1^-$ ,  $\mathcal{L} = 500 \text{ fb}^{-1}$ :


$m_{\tilde{\chi}_1^0} = 96.12 \pm 0.05 \text{ GeV}$ ,  $m_{\tilde{\chi}_2^0} = 177.13 \pm 0.05 \text{ GeV}$ ,  $m_{\tilde{\chi}_1^\pm} = 174.90 \pm 0.05 \text{ GeV}$ ,  $\Delta\sigma(\tilde{\chi}_1^0 \tilde{\chi}_2^0) = \pm 2\%$  [K. Desch, G. Moortgat-Pick '02]



$\Rightarrow \phi_1 = 30^\circ \pm 3^\circ$

Determination of  $m_{\tilde{t}_1}$ ,  $\theta_{\tilde{t}}$  from  $\sigma(e^+e^- \rightarrow \tilde{t}_1\tilde{t}_1)$  with polarized beams:  
 [R. Keränen, H. Nowak, A. Sopczak '00]

stop into c neutralino 80/60 pol



$$\Rightarrow \frac{\Delta m_{\tilde{t}_1}}{m_{\tilde{t}_1}} \approx 0.5\%, \quad \frac{\Delta \cos\theta_{\tilde{t}}}{\cos\theta_{\tilde{t}}} \approx 1.5\%$$



## Complementarity of LHC and LC:

⇒ Results obtained at one collider can be used for improving experimental analyses at the other

⇒ investigated in “LHC / LC Study Group”

[www.ippp.dur.ac.uk/~georg/lhclc](http://www.ippp.dur.ac.uk/~georg/lhclc)

Collaborative effort of Hadron Collider (HC) and Linear Collider (LC) community


Started in spring 2002, currently about 190 working group members from ATLAS, CMS, LC working groups, theory + Tevatron contact person

## Example: SUSY parameters at LHC and LC

### Reconstruction of sparticle masses at the LHC

[*B. Gjelsten, E. Lytken, D. Miller, P. Osland, G. Polesello, M. Chiorboli, A. Tricoli*]



Complicated decay chains for squarks and gluinos




Examples worked out for SPS1a from ATLAS and CMS  
main tool: dilepton “edge” from  $\tilde{\chi}_2^0 \rightarrow l^+ l^- \tilde{\chi}_1^0$

## Sbottom/squark and gluino reconstruction:

Edge in same flavour-opposite sign lepton distribution (left), invariant mass distributions with kinematical endpoints (right)



Strong correlation between slepton mass and LSP mass, LSP mass can be constrained at LHC at the 10% level only:





⇒ Take LSP mass as input from LC

Select events close to the edge and combine with b-jet / q-jet

$$\vec{p}(\tilde{\chi}_2^0) = \left(1 - \frac{m(\tilde{\chi}_1^0)}{m(\ell\ell)}\right) \vec{p}_{\ell\ell}$$


⇒ Get sbottom/squark mass if LSP mass is known

Squark peak (left) and gluino reconstruction from  $(\chi_2^0 bb)$  invariant mass distribution (right):



$m_{\tilde{g}}$  as function of the LSP mass:

$$\Rightarrow \Delta m_{\tilde{g}} \approx \Delta m_{LSP}$$



Accuracies for the case of the LHC alone (left) and with the LC measurement of the LSP mass with 0.2% accuracy (middle) and 1.0% accuracy (right column) in GeV:


	LHC	LHC+LC (0.2%)	LHC+LC (1.0%)
$\Delta m_{\tilde{\chi}_1^0}$	9.2	0.2	1.0
$\Delta m_{\tilde{t}_R}$	9.2	0.5	1.0
$\Delta m_{\tilde{\chi}_2^0}$	9.0	0.3	1.0
$\Delta m_{\tilde{b}_1}$	23.1	16.9	17.0
$\Delta m_{\tilde{q}_L}$	15.0	5.1	5.3

⇒ LC input improves accuracy significantly

One step further:

Determination of the mass of the heaviest neutralino at the LHC using LC input from the neutralino/chargino sector:

[J. Kalinowski, G. Moortgat-Pick, M. Nojiri, G. Polesello]



⇒ Need besides LSP mass also masses of sleptons and charginos from LC in order to correctly identify  $\tilde{\chi}_4^0$

⇒ Feeding  $m(\tilde{\chi}_4^0)$  back into LC analysis improves accuracy of parameter determination at the LC

Even further:

## Full reconstruction of stop/sbottom parameters with LHC $\otimes$ LC

[J. Hisano, K. Kawagoe, M. Nojiri]


Complete set of electroweak SUSY parameters (from LC) and branching ratios used to exploit LHC rate measurements

Stop/sbottom sector determined by 5 parameters, e.g.  $m_{\tilde{b}_1}, m_{\tilde{b}_2}, m_{\tilde{t}_1}, \theta_{\tilde{b}}, \theta_{\tilde{t}}$

- Take  $m_{\tilde{b}_1}, m_{\tilde{b}_2}$  from previous study
- $\Rightarrow$  need three more observables:
  - $tb$  invariant mass distribution
  - rate of “edge-events” in  $m_{tb}$  distribution (chargino chain)
  - rate of events in  $llb$  distribution ( $\tilde{\chi}_2^0$  chain)



$m_{tb}$  distribution for SPS1a:





Involved couplings are stop–sbottom–W, top–sbottom–chargino and stop–bottom–chargino

⇒ If chargino couplings + BR's are known then observed rates are sensitive to sbottom/stop mixing parameters

Determination of  $\theta_{\tilde{b}}$  from  $\text{BR}(\tilde{g} \rightarrow b\tilde{b}_2 \rightarrow bb\tilde{\chi}_2^0)/\text{BR}(\tilde{g} \rightarrow b\tilde{b}_1 \rightarrow bb\tilde{\chi}_2^0)$  in SPS1a, sbottom masses and parameters of chargino/neutralino sector are assumed to be known (left)

Determination of  $m_{\tilde{t}_1}$ ,  $\theta_{\tilde{t}}$  assuming also  $\theta_{\tilde{b}}$  is known (right)



$\Rightarrow \mathcal{O}(50\%)$  determination of mixing angles,  $\Delta m_{\tilde{t}_1}/m_{\tilde{t}_1} < 5\%$

Results used for:

mSUGRA fit to LHC  $\oplus$  LC data in SPS1a


[B. Allanach, S. Kraml, W. Porod]

LHC / LC complementarity in mSUGRA fits

[D. Tovey]

Discrimination between different SUSY-breaking scenarios

[B. Allanach, D. Grellscheid, F. Quevedo]




$\Rightarrow$  Need %-level accuracy to distinguish between different models

## Extrapolation to physics at high scales

from combination of LHC and LC results, precise measurement of masses of SUSY particles, couplings

E.g.: Test of gaugino mass unification


[G. Blair, W. Porod, P. Zerwas '01]



# Higgs searches at the Tevatron and the LHC

## mSUGRA scenario with CDM constraints

$$\mu > 0, \tan \beta = 10, 50: \left[ \sigma(gg \rightarrow h) \times \text{BR}(h \rightarrow \gamma\gamma) \right]_{\text{CMSSM}} / \left[ \sigma(gg \rightarrow h) \times \text{BR}(h \rightarrow \gamma\gamma) \right]_{\text{SM}} :$$



- ⇒ no significant suppression of  $\sigma(gg \rightarrow h) \times \text{BR}(h \rightarrow \gamma\gamma)$  compared to SM
- ⇒ Discovery of lightest Higgs boson within about one year at LHC possible  
[J. Ellis, S. Heinemeyer, K. Olive, G. W. '01]

Similar results in GMSB and AMSB scenarios [A. Dedes, S. Heinemeyer, S. Su, G. W. '03]

## Suggested benchmarks for Higgs searches at the Tevatron and the LHC

[M. Carena, S. Heinemeyer, C. Wagner, G. W. '02]

Scenarios for general MSSM, no specific SUSY-breaking scenario assumed, no indirect constraints,  $M_A$ ,  $\tan\beta$  varied


- $m_h^{\max}$ -scenario:  $X_t = 2 M_{\text{SUSY}}$  (FD),  $M_{\text{SUSY}} = 1$  TeV  
 $\Rightarrow$  maximal  $m_h(\tan\beta)$  for fixed  $m_t$ ,  $M_{\text{SUSY}}$
- no-mixing scenario:  $X_t = 0$ ,  $M_{\text{SUSY}} = 2$  TeV
- gluophobic Higgs scenario:  $M_{\text{SUSY}} = 350$  GeV,  $X_t = -750$  GeV (FD)  
 $\Rightarrow$  suppression of  $gg \rightarrow h$
- small  $\alpha_{\text{eff}}$  scenario:  
 $M_{\text{SUSY}} = 800$  GeV,  $\mu = 2.5 M_{\text{SUSY}}$ ,  $X_t = -1100$  GeV (FD)  
 $\Rightarrow$  suppression of  $h \rightarrow b\bar{b}$ ,  $h \rightarrow \tau\tau$

Small  $\alpha_{\text{eff}}$  scenario:

$\sigma(q\bar{q} \rightarrow Vh) \times \text{BR}(h \rightarrow b\bar{b})$  at the Tevatron:

[M. Carena, S. Heinemeyer, C. Wagner, G. W. '02]

small  $\alpha_{\text{eff}}$ :  $\sigma(V^* \rightarrow Vh) \times \text{BR}(h \rightarrow b\bar{b})$




⇒ Significant suppression  
for  $M_A \lesssim 250$  GeV

## Gluophobic Higgs scenario:

$\sigma(gg \rightarrow h) \times \text{BR}(h \rightarrow \gamma\gamma)$  at the LHC:

[*M. Carena, S. Heinemeyer, C. Wagner, G. W. '02*]

gluophobic Higgs:  $\sigma(gg \rightarrow h) \times \text{BR}(h \rightarrow \gamma\gamma)$



⇒ Large suppression in whole  $M_A$ - $\tan\beta$  plane



# Precision physics in the MSSM Higgs sector



Precise measurement of Higgs branching ratios

⇒ Sensitivity to deviations SM / MSSM

E.g.: Prediction for  $\sigma(e^+e^- \rightarrow Zh) \times \text{BR}(h \rightarrow WW^*)$  in parameter region allowed by cosmology: comparison mSUGRA – SM:

[J. Ellis, S. Heinemeyer, K. Olive, G. W. '02]

$\mu > 0$ ,  $\tan \beta = 10, 50$ :



⇒ In allowed parameter space: sizable deviations from SM predictions for precision observables in the Higgs sector possible


## Combination of LHC data on heavy Higgs states with LC data on the light $\mathcal{CP}$ -even Higgs

[*K. Desch, S. Heinemeyer, G.W.*]

Assume: LHC information on  $M_A$ ,  $\tan \beta \oplus$  (LHC  $\otimes$  LC) information on stop/sbottom masses  $\oplus$  LHC / LC measurement of  $m_h$ :



$M_A$ : 15% accuracy,  $m_{\tilde{t}_1}, m_{\tilde{t}_2}, m_{\tilde{b}_1}, m_{\tilde{b}_2}$ : 5% accuracy,  $\tan \beta > 15$

$\text{BR}(h \rightarrow b\bar{b})$ : 2.5% accuracy,  $\text{BR}(h \rightarrow WW^*)$ : 5% accuracy



⇒ Comparison of MSSM prediction based on assumed inputs with BR's measured at the LC yields very sensitive test of the model

⇒ Indirect determination of trilinear coupling  $A_t$ :



Precise measurement of  $m_t$  at the LC crucial,  $\delta m_t \lesssim 100 \text{ MeV}$

$\Delta m_t^{\text{LC}}$  vs.  $\Delta m_t^{\text{LHC}}$  ⇒ accuracy of  $A_t$  determination improved by factor 3

Necessary improvements in accuracy of theoretical prediction in order to match experimental precision at LHC,  $\delta m_h^{\text{exp}} \approx 0.2 \text{ GeV}$ :

- Uncertainty from experimental errors of input parameters:

⇒ Complementarity example:

In order to match

experimental precision at LHC,  $\delta m_h^{\text{exp}} \approx 0.2 \text{ GeV}$

need

LC precision on  $m_t$ ,  $\delta m_t^{\text{exp}} \lesssim 0.2 \text{ GeV}$

- Uncertainty from unknown higher-order corrections:

⇒ Need improvement by more than a factor 10!

## Summary of Lectures 4 and 5:

- SUSY requires light Higgs boson:  
definite and robust prediction of SUSY theories, testable at next generation of colliders
- mSUGRA in agreement with exp. constraints from Higgs search, CDM,  $(g_\mu - 2)$ ,  $B \rightarrow X_s \gamma$
- EW precision tests: MSSM yields equally complete description as SM  
**Global fit:** better agreement of MSSM with data for  $g_\mu - 2$ ,  $M_W$  than SM, no improvement for  $A_{FB}^b$ , NuTeV, **similar fit quality**

- SUSY searches (same for other kinds of new physics) at next generation of colliders:

Tevatron and LHC: big discovery potential

LC: high-precision physics

⇒ need both to

- establish SUSY experimentally
- disentangle patterns of SUSY breaking

- Complementarity LHC / LC can be exploited for improving analyses at both machines
- Precision physics in Higgs sector: ⇒ Very sensitive test of ew theory

## Outlook:

- Low-energy Supersymmetry continues to be our best bet for physics beyond the Standard Model

- Data rules:

We need experimental information from Tevatron, LHC, LC,  $\nu$  experiments, dark matter searches, low-energy experiments, . . . to verify / falsify models of new physics understanding

- The experiments in the next years will bring a decisive test of our ideas about low-energy SUSY

⇒ Very exciting prospects for the coming years

Expect the unexpected!