

SUMMER SCHOOL ON PARTICLE PHYSICS

16 June - 4 July 2003

GRAND UNIFICATION AND FERMION MASSES

S. M. BARR
Bartol Research Institute
University of Delaware
Delaware, DE
U.S.A.

Grand Unification and Fermion Masses

(ICTP July 1-4, 2003)

Hints in favor of GUTs:

(1) Remarkable fit of SM quark and lepton multiplets into multiplets of $SU(5)$, $SO(10)$

(2) Unification of gauge couplings
at scale $\approx 2 \times 10^{16}$ GeV = M_G
(in the MSSM).

(3) Neutrino masses

(a) ν_R predicted by Pati-Salam, $SO(10)$

(b) M_R as inferred from see-saw
formula $\approx M_G$.

(4) Hints from q, l masses, mixings

(a) $m_b^0 \approx m_\tau^0$, Similar q, l hierarchies

(b) $\theta_{CKM} \ll 1$ ($\theta_{CKM} = 0$ in minimal $SO(10)$)

(c) Relationships among V_{cb} , m_s/m_b ,
 $V_{\mu 3}$ etc (see lopsided models)

Areas of research

- Gauge hierarchy problem,
 $\frac{2}{3}$ splitting problem,
 $d=5$ proton decay operators
- Leptogenesis/baryogenesis
- Neutrino masses and mixings
- Quark and lepton masses and mixings (the flavor problem)
- Flavor changing processes
 $(\mu \rightarrow e\gamma, \tau \rightarrow \mu\gamma \text{ etc})$
- Other low energy tests?
- Higher dimensions ($d=5, 6$ brane models)
- Strings?

(iii)

My goal in these lectures is to explain the basic

- Schemes of unification

(Pati-Salam, $SU(5)$, $SO(10)$, flipped $SU(5)$, ...)

- issues and problems

(gauge hierarchy, doublet-triplet splitting, $d=5$ proton decay operators, quark and lepton masses.)

and ● practical group theory

Outline

I. $SU(5)$ unification

long { II. $SO(10)$ and other groups
for unification
(Pati-Salam, flipped $SU(5)$, ...)

III. The gauge hierarchy problem

short { IV. Fermion Masses

LECTURE I SU(5) UNIFICATION

A. Basic Group Theory Facts

Group $G = \{g\}$

- closure $g_1 g_2 = g_3$
- identity $eg = ge = g$
- inverse $gg^{-1} = g^{-1}g = e$
- assoc. $g_1(g_2g_3) = (g_1g_2)g_3$

Lie groups

"Represent" elements by unitary matrices

$$U(g) = e^{i \sum_a \theta^a \lambda^a}$$

↑ generators
↑ parameters associated with element g.

generators λ^a are hermitian matrices, satisfying the group algebra

$$[\lambda^a, \lambda^b] = i f^{abc} \lambda^c$$

↑ structure constants

Number of generators = dim. of group.

If U is a $d \times d$ matrix representing an element of the group, it can be thought of as acting on a d -component object

$$[U] \begin{pmatrix} T_1' \\ T_2' \\ \vdots \\ T_d' \end{pmatrix} = \begin{pmatrix} T_1' \\ T_2' \\ \vdots \\ T_d' \end{pmatrix}$$

Strictly speaking, the U matrices form a "representation" of the group elements. But physicists often talk of the d -component multiplet as being a "representation".

Recall: always exists an adjoint representation 1.2

$d \times d$ matrices, where $d = \text{dimension of group}$
 $= \# \text{ of generators}$

generators in the adjoint representation are a set of d hermitian $d \times d$ matrices

→ These act on d -component multiplets, which are called "adjoint representations" or "adjoint multiplets"

Proof: Jacobi identity

$$0 = [[\lambda^a, \lambda^b], \lambda^c] + [[\lambda^c, \lambda^a], \lambda^b] + [[\lambda^b, \lambda^c], \lambda^a]$$

$$0 = f^{abd} f^{dce} + f^{cad} f^{dbe} + f^{bcd} f^{dae}$$

$$\text{Call } f^{abd} \equiv i(\lambda^a)^{bd}$$

$$0 = i f^{abd} (\lambda^d)^{ce} - (\lambda^a)^{cd} (\lambda^b)^{de} + (\lambda^b)^{cd} (\lambda^a)^{de}$$

which is group algebra!

B) SU(N) GROUP THEORY

$$\left\{ \begin{array}{l} N \\ \bar{N} \end{array} \right. \begin{array}{l} T^\alpha \rightarrow T^{\alpha'} = U^{\alpha'}_\alpha T^\alpha \quad \underline{\alpha=1, \dots, N} \\ T_\alpha \rightarrow T_{\alpha'} = U_{\alpha'}^\alpha T_\alpha \end{array}$$

These multiplets are called "fundamental rep.", "anti fundamental rep."

$$T_\alpha = (T^\alpha)^* \quad , \quad U_{\alpha'}^\alpha = (U^{\alpha'}_\alpha)^*$$

$$\sum_{\alpha=1}^N T^\alpha T_\alpha = \text{invariant} \Rightarrow U U^\dagger = I = U^\dagger U$$

$U = N \times N$ unitary matrices, $\det U = 1$

$N \times N$, unitary, $\det = 1$ matrices represent the elements of group $SU(N)$

Such matrices can be written in terms of $N^2 - 1$ parameters

$$U_{\alpha}^{\alpha'} = \left(e^{i \sum_{a=1}^{N^2-1} \theta^a \lambda^a} \right)_{\alpha}^{\alpha'} \quad \begin{matrix} a=1, \dots, N^2-1 \\ \alpha, \alpha'=1, \dots, N \end{matrix}$$

$N^2 - 1$ generators. In this fundamental representation they are $N \times N$ traceless hermitian matrices.

[I will denote "adjoint indices" by a, b, c, \dots
"fundamental indices" of $SU(N)$ by $\alpha, \beta, \gamma, \dots$]

Adjoint rep.

For $SU(N)$ there exist $(N^2 - 1)$ -component multiplets T^a that get acted upon by $(N^2 - 1) \times (N^2 - 1)$ unitary matrices. [Adjoint rep.]

$$T^{\alpha}_{\beta} \equiv \sum_{a=1}^{N^2-1} T^a (\lambda^a)^{\alpha}_{\beta}$$

Two ways of writing this multiplet
 $\alpha, \beta = 1, \dots, N$; $a = 1, \dots, N^2 - 1$

$$T^{\alpha}_{\beta} \rightarrow T^{\alpha'}_{\beta'} = U^{\alpha'}_{\alpha} \times U_{\beta'}^{\beta} T^{\alpha}_{\beta}$$

$$T \rightarrow U T U^{\dagger}$$

[NOTE: CAN ALSO DENOTE λ^a BY λ^{α}_{β}
eg. IN $SU(3)$ $\lambda^6 + i\lambda^7 = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 1 \\ 0 & 0 & 0 \end{pmatrix} \equiv \lambda^2_3$]

Let a Higgs field be in such an adjoint representation (i.e. multiplet) of an $SU(N)$ gauge sym.

$$H^\alpha_\beta = \sum_{a=1}^{N^2-1} H^a (\lambda^a)^\alpha_\beta$$

$$\begin{aligned} \langle H \rangle \rightarrow U \langle H \rangle U^\dagger &= e^{i \sum_b \theta^b \lambda^b} \langle H \rangle e^{-i \sum_c \theta^c \lambda^c} \\ &= \langle H \rangle + i \sum_b \theta^b [\lambda^b, \langle H \rangle] + \dots \end{aligned}$$

$\langle H \rangle$ is left invariant by "unbroken generators"

$$\Rightarrow \begin{cases} \text{Unbroken generators: } [\lambda^a, \langle H \rangle] = 0 \\ \text{Broken generators: } [\lambda^a, \langle H \rangle] \neq 0. \end{cases}$$

c) $SU(5)$ multiplets and symmetry breaking

$$\left\{ \begin{array}{l} 5 \\ \bar{5} \end{array} \right. \psi^\alpha = \begin{pmatrix} \psi^1 \\ \psi^2 \\ \hline \psi^3 \\ \psi^4 \\ \psi^5 \end{pmatrix} \left\{ \begin{array}{l} \psi^i \quad i=1,2 \quad SU(2)_L \\ \psi^a \quad a=3,4,5 \quad SU(3)_C \end{array} \right.$$

24 generators = 5×5 traceless, hermitian matrices (in fundamental rep.)

$$(\lambda^a)^\alpha_\beta = \begin{bmatrix} SU(2) & 0 \\ 0 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & SU(3) \end{bmatrix}, \begin{bmatrix} 0 & m \\ m^\dagger & 0 \end{bmatrix}, \begin{bmatrix} \frac{1}{2} & \frac{1}{2} \\ & -\frac{1}{3} & \frac{1}{3} & \frac{1}{3} \end{bmatrix} \equiv \begin{matrix} (1, 3, 0) & (8, 1, 0) & (3, 2, \frac{1}{6}) + \text{h.c.} & (1, 1, 0) \end{matrix}$$

Now, let there be a Higgs in an adjoint multiplet of $SU(5)$, and let its VEV be

$$\langle \Omega \rangle = \begin{bmatrix} 1/2 & & & & \\ & 1/2 & & & \\ & & -1/3 & & \\ & & & -1/3 & \\ & & & & -1/3 \end{bmatrix} \omega = \left(\frac{Y}{2} \right) \omega$$

$\left\langle \lambda^a \right\rangle$

[Note: VEVs of adjoint Higgs multiplets always point in direction of some generator (or linear combination of gens.)]

Which generators are unbroken then?

$$[\lambda^a, \langle \Omega \rangle] = 0$$

Unbroken generators are those of

$$SU(3)_c \times SU(2)_L \times U(1)_Y = G_{SM}$$

Embedding of SM multiplets in $SU(5)$ \longrightarrow

$$\bar{5}_L = \psi_L = \begin{pmatrix} \psi_i \\ \psi_a \end{pmatrix} = \begin{pmatrix} (1, 2, -1/2) \\ (\bar{3}, 1, 1/3) \end{pmatrix} \equiv \begin{pmatrix} \bar{2} \\ 3 \end{pmatrix} = \begin{pmatrix} \begin{pmatrix} e_L^- \\ \nu_L \end{pmatrix} \\ d_L^c \end{pmatrix} \text{ etc}$$

\uparrow my notation

$$10_L = \psi_{[\alpha\beta]} = \begin{bmatrix} \psi_{[12]} & \psi_{ia} \\ \psi_{ai} & \psi_{ab} = \epsilon^{abcd} \psi_d \end{bmatrix} = \begin{bmatrix} (1, 1, +1) & (\bar{3}, 2, 1/6) \\ & (\bar{3}, 1, -2/3) \end{bmatrix} = \begin{bmatrix} e_L^+ & (u_L^c) \\ & d_L^c \end{bmatrix}$$

$$24_g = A^a = \begin{bmatrix} A^i_j & A^i_a \\ A^i_a & A^a_b \end{bmatrix} = \begin{bmatrix} (1, 3, 0) & (\bar{3}, 2, 5/6) \\ (3, 2, -5/6) & (8, 1, 0) \\ & + (1, 1, 0) \end{bmatrix}$$

$A^a(A^b)^c = A^c{}^a{}_b$

Key point: $\psi_L(r) \xleftrightarrow{C} \psi_R(\bar{r})$ 1.6

SM family (1st family)

$$(1, 2, -\frac{1}{2})_L = \begin{pmatrix} \nu_L \\ e_L^- \end{pmatrix} \equiv L$$

$$(1, 1, -1)_R = e_R^-$$

$$(1, 1, 0)_R = N_R$$

$$(3, 2, \frac{1}{6})_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix} \equiv Q$$

$$(3, 1, \frac{2}{3})_R = u_R$$

$$(3, 1, -\frac{1}{3})_R = d_R$$

$$\xleftrightarrow{C} (1, 1, +1)_L = e_L^+$$

$$\xleftrightarrow{C} (1, 1, 0)_R = N_L^c$$

$$\xleftrightarrow{C} (\bar{3}, 1, -\frac{2}{3}) = u_L^c$$

$$\xleftrightarrow{C} (\bar{3}, 1, +\frac{1}{3}) = d_L^c$$

Two ways to write a mass term (or Yukawa term)

$$m \psi_R(r_2) \overleftarrow{\text{DIRAC BAR}} \psi_L(r_1) = m [\psi_R(r_2)]^\dagger \gamma^0 \psi_L(r_1)$$

or \parallel



$$m \psi_L(\bar{r}_2)^T C \psi_L(r_1)$$

More convenient for GUTs (also SUSY!)

eg electron mass:

$$m(\overline{e_R^-}) e_L^- = m(e_L^+)^T C e_L^-$$

often do not bother to show



Arrows follow flow of "left-handedness"

Think of chirality as a charge $\chi(L) = -1$, $\chi(R) = +1$

$$\Delta\chi = 2$$

(D) Fermion Masses

$$\left\{ \begin{aligned} \text{FAMILY} &= 10_L + \bar{5}_L + (1_L) \\ &\quad \parallel \\ &(\psi^{12}, \psi^{1a}, \psi^{ab})_L + (\psi_i, \psi_a)_L + \psi_L \\ &\quad \parallel \\ &(e_L^+, (u)_L, u_L^c) + ((\nu)_L^-, d_L^c) + N_L^c \end{aligned} \right.$$

$$\left\{ \begin{aligned} \text{HIGGS ARE IN } &5_H + \bar{5}_H \\ &\quad \parallel \\ &(H^i, H^a) + (H_i, H_a) \\ &(H_u, H_{uc}) + (H_d, H_{dc}) \\ \text{[In non-SUSY: } &\bar{5}_H = (5_H)^* \text{ i.e. } H_\alpha = (H^\alpha)^* \\ \text{In SUSY } &5_H \text{ and } \bar{5}_H \text{ are distinct, L-H superfields.]} \end{aligned} \right.$$

YUKAWA TERMS (minimal SU(5))

$$\left\{ \begin{aligned} (Y_u)_{AB} (10_{LA} 10_{LB}) \langle 5_H \rangle &= (Y_u)_{AB} \epsilon_{\alpha\beta\gamma\delta} (\psi_A^{\alpha\beta} \psi_B^{\gamma\delta} \langle H^\epsilon \rangle) \\ &= (Y_u)_{AB} (\psi_A^{12} \psi_B^{bc} \langle H^2 \rangle) = (Y_u)_{AB} u_A u_B^c \langle H_u \rangle \end{aligned} \right.$$

NOTE: $Y_u = Y_u^T \Rightarrow \boxed{M_u = M_u^T \propto Y_u}$

$$\left\{ \begin{aligned} (Y_D)_{AB} (10_{LA} \bar{5}_{LB}) \langle \bar{5}_H \rangle &= (Y_D)_{AB} (\psi_A^{\alpha\beta} \psi_{B\alpha} \langle H_\beta \rangle) \\ &= (Y_D)_{AB} (\psi_A^{12} \psi_{B1} + \psi_A^{22} \psi_{B2}) \langle H_2 \rangle \\ &= (Y_D)_{AB} (e_A^+ e_B^- + d_A d_B^c) \langle H_d \rangle \end{aligned} \right.$$

NOTE: $\boxed{M_D = M_D^T \propto Y_D}$

$$M_D = M_L^T \quad (\text{in minimal SU(5)}) \quad 1.8$$

$$\Rightarrow \begin{cases} m_b^0 = m_\tau^0 & (\text{Good within 20\% in SUSY GUTs}) \\ m_s^0 = m_\mu^0 & (m_s^0 \lesssim \frac{1}{3} m_\mu^0, \text{ from SUSY GUTs RGE}) \\ m_d^0 = m_e^0 & (m_d^0 \approx 3 m_e^0, \text{ " " " "}) \end{cases}$$

If there exist N_L^c (Right-handed ν 's):


$$\begin{cases} (Y_N)_{AB} (\bar{5}_{LA} 1_{LB}) \langle S_H \rangle = (Y_N)_{AB} (\psi_{A1} \psi_B) \langle H^c \rangle \\ = (Y_N)_{AB} (\psi_{A2} \psi_B) \langle H^2 \rangle = (Y_N)_{AB} \nu_A N_B^c \langle H_u \rangle \\ \Rightarrow \boxed{M_N \propto Y_N} \text{ (}\nu \text{ Dirac mass matrix)} \approx V_u \end{cases}$$

$$\begin{cases} (M_R)_{AB} (1_{LA} 1_{LB}) = (M_R)_{AB} (N_A^c N_B^c) \\ \boxed{M_R} \text{ (}\nu_R \text{ Majorana mass matrix)} \approx M_G \end{cases}$$

$$\Rightarrow (\nu \ N^c)^T \begin{pmatrix} 0 & M_N \\ M_N^T & M_R \end{pmatrix} \begin{pmatrix} \nu \\ N^c \end{pmatrix} \rightarrow (\nu \ N^c)^T \begin{pmatrix} M_\nu & 0 \\ 0 & M_R \end{pmatrix} \begin{pmatrix} \nu \\ N^c \end{pmatrix}$$

$$\boxed{M_\nu \approx -M_N M_R^{-1} M_N^T} \quad \text{SEE-SAW FORMULA}$$

DIAGRAMMATICALLY:



$$\begin{aligned} M_\nu &\approx M_N \frac{1}{E + M_R} M_R^* \frac{1}{E + M_R} M_N^T \\ &\approx M_N M_R^{-1} M_N^T \end{aligned}$$


So, in minimal SU(5) there are four independent mass matrices for q and l :

$$M_D = M_L^T, M_u, M_N, M_R$$




$\underbrace{\hspace{10em}}_{M_\nu}$

(E) PROTON DECAY OPERATORS

TWO KINDS OF DIAGRAMS FOR PDK IN NON-SUSY SU(5):
GAUGE BOSON MEDIATED, HIGGS BOSON MEDIATED



HIGGS BOSON MEDIATED



The Higgs-mediated proton decay is caused by exchange of colored Higgs

These must be superheavy

⇒ "DOUBLET-TRIPLET SPLITTING PROBLEM"

$$5_H = H^\alpha = \begin{pmatrix} H^i \\ H^a \end{pmatrix} \begin{matrix} \leftarrow \text{Weak-scale mass} \\ \leftarrow \text{GUT scale mass} \end{matrix}$$

THIS IS AN ASPECT OF THE INFAMOUS "GAUGE HIERARCHY PROBLEM"

- THREE ASPECTS:
- (a) 2/3 SPLITTING PROBLEM
 - (b) FINE-TUNING PROBLEM
(stability under radiative corrections)
 - (c) SMALL NUMBER PROBLEM
(why $M_W/M_G \sim 10^{-14}$)
- (d) (In SUSY) "μ PROBLEM"

Why do all of these operators conserve B-L?

In minimal SU(5) model:

$$(10_L 10_L) 5_H + (10_L \bar{5}_L) \bar{5}_H + (\bar{5}_L 1_L) 5_H + M^2 \bar{5}_H 5_H$$

$$X = \begin{matrix} +1 & +1 & -2 & +1 & -3 & +2 & -3 & +5 & -2 & +2 & -2 \end{matrix}$$

Can assign a global charge X that is conserved by couplings

Now $\langle 5_H \rangle, \langle \bar{5}_H \rangle$ break $\underbrace{U(1)_{I_{3L}}}_{\text{local}} \times \underbrace{U(1)_Y}_{\text{local}} \times \underbrace{U(1)_X}_{\text{global}}$

by 't Hooft mechanism there remain

$$\underbrace{U(1)_Q}_{\text{local}} \times \underbrace{U(1)_{B-L}}_{\text{global}} \quad Q = I_{3L} + Y/2$$

$$B-L = \frac{4}{5} \frac{Y}{2} + \frac{1}{5} X$$

In SO(10), as we shall see, $U(1)_X$ is contained in SO(10) and thus a local sym.
 \Rightarrow B-L is a generator of SO(10).

NOTE In SU(5), if there are right-handed neutrinos, they can have Majorana masses that break L and conserve B, thus ~~(B-L)~~.

$$M_R \begin{pmatrix} 1_L & 1_L \\ +5 & +5 \end{pmatrix} \Rightarrow \Delta X = -10 \Rightarrow \Delta(B-L) = -2$$

IMPORTANT FOR "LEPTOGENESIS".

PROTON DECAY OPERATORS IN NON SUSY SU(5)

gauge
and
higgs
mediated

$$\left\{ \begin{array}{l} 10_L^+ 10_L^+ 10_L 10_L \rightarrow (e_L^+ u_L^c)^t u_L d_L = e_R^- u_R u_L d_L \\ 10_L^+ \bar{5}_L^+ 10_L \bar{5}_L \rightarrow (u_L^c d_L^c)^t Q_L L_L = u_R d_R Q_L L_L \end{array} \right.$$

only
higgs
mediated

$$\left\{ \begin{array}{l} 10_L 10_L 10_L \bar{5}_L \rightarrow e_L^+ u_L^c u_L^c d_L^c = (e_R^- u_R u_R d_L)^t \\ \text{and} \\ Q_L Q_L Q_L L_L \end{array} \right.$$

IN SUSY SU(5)

$$\mathcal{L} \supset \int d^2\theta d^2\bar{\theta} (10_L^+ 10_L^+ 10_L 10_L + 10_L^+ \bar{5}_L^+ 10_L \bar{5}_L) / M_G^2$$

Since both Φ and Φ^+ are involved, these are D-terms with $\int d^2\theta d^2\bar{\theta}$ thus by dimensional analysis $1/M_G^2$.

\Rightarrow "d=6" proton decay operators
caused by exchange of gauge and higgs bosons

But also allowed: "d=5" proton decay operators

$$\mathcal{L} \supset \int d^2\theta (10_L 10_L 10_L \bar{5}_L) / M_G$$

Term in superpotential W as only Φ so
by dimensional analysis only suppressed by $1/M_G$.

NOTE: From HIGGS NO EXCHANGE. DANGEROUS!


Also could have (disastrous) $d = 4$ operators" 1.13
 such as $\int d^2\theta (10_L \bar{5}_L \bar{5}'_L)$, but these can be
 forbidden by imposing "matter parity" (\approx R parity):

Z_2 under which $\begin{cases} \text{"matter"} (\equiv q + l) \text{ are odd} \\ \text{higgs, gauge} \text{ are even.} \end{cases}$

(Such a Z_2 can arise automatically in $SO(10)$ as a subgroup of $SU(16)$)


$d = 5$ p decay comes from (colored) Higgsino exchange

$10_L 10_L 10_L \bar{5}_L$ from
 M_G



partner of Higgs boson exchange
 graph shown earlier

Why not $d = 5$ from exchange of gaugino?




No "chirality flip" on gaugino line (since $m_{24_L 24_L}$,
 being a gaugino mass $\propto M_{\text{sysy}} \ll M_G$).

$$\Rightarrow \frac{1}{p + M_G} = \frac{p - M_G}{p^2 + M_G^2} \sim \frac{1}{M_G^2}$$

LOOK AT HIGGSINO EXCHANGE DIAGRAM AGAIN.

NEED A CHIRALITY FLIP


SUPERFIELD DIAGRAM:



NOTE: IN SUSY SU(5) $\bar{5}_H$, 5_H ARE NOT CONJUGATES OF EACH OTHER, BUT BOTH LEFT-HANDED CHIRAL SUPERFIELDS.

SO "X" IN DIAGRAM IS FROM $\mu \bar{5}_H 5_H$

$$W \supset (10_L 10_L) 5_H + (10_L \bar{5}_L) \bar{5}_H + (\bar{5}_L 1_L) 5_H + \mu \bar{5}_H 5_H$$




$\rightarrow \mu_3 H_{dc} H_{uc}$

$A_{p\text{ decay}} \propto \frac{\mu_3}{M_{H_{uc}} M_{H_{dc}}}$

CRUCIAL POINT

1.15




WHY NOT DISASTROUS IF $\sim \frac{1}{M_G}$?

ADDITIONAL SMALL FACTORS (RELATIVE TO GAUGE MED)

(a) Couplings are Yukawa couplings of the light quarks and leptons $\sim \frac{m_{q,l}}{v}$

(b) These diagrams involve scalar quarks and leptons which must be turned into quarks and leptons by gaugino \Rightarrow loop



Calculation is complicated and involves hadronic matrix element.


FOR SUSY SU(5) (minimal) [See H. Murayama and A. Pierce, hep-ph/0108104]

$$3.5 \times 10^{14} \leq M_{H_c} \leq 3.6 \times 10^{15} \text{ GeV} \quad (90\% \text{ c.l. from R.G.E.})$$

$$\text{BUT } M_{H_c} \geq 7.6 \times 10^{16} \text{ GeV} \quad (\text{from superK limit on } p \rightarrow K^+ \bar{\nu})$$

PROBLEM OF HIGGSINO-MEDIATED P DECAY (or $d=5$ p decay)

LECTURE II : $SO(10)$ AND OTHER GROUPS FOR UNIFICATION ^{2.1}



(A) MOST IMPORTANT GROUPS FOR UNIFICATION

PATI-SALAM $(SU(4)_c \times SU(2)_L \times SU(2)_R)$

First unification scheme, not "grand unified" but does have $g-l$ unification.

- Implies Right-handed neutrinos exist.
- Can have Left-Right symmetry.

SU(5) (GEORGI-GLASHOW)

- Smallest "grand unified" group
[Only rank-4 unification group
⇒ one stage of symmetry breaking
⇒ most predictive for p decay]

SO(10)

Contains both $SU(5)$ and Pati-Salam groups

- Implies Right-handed neutrinos exist
- Has a Left-Right symmetry
- Is grand unified
- Complete $g-l$ unification of a family
- Most predictive for g, l masses
- Automatic anomaly freedom.

(B) PATI-SALAM $SU(4)_c \times SU(2)_L \times SU(2)_R$

$$\begin{array}{ccc}
 \begin{array}{c} \updownarrow SU(2)_L \\ \left(\begin{array}{cccc} u & u & u & \nu \\ d & d & d & e^- \end{array} \right)_L \\ \leftarrow SU(4)_c \rightarrow \end{array} & & \begin{array}{c} \updownarrow SU(2)_R \\ \left(\begin{array}{cccc} u & u & u & N \\ d & d & d & e^- \end{array} \right)_R \\ \leftarrow SU(4)_c \rightarrow \end{array}
 \end{array}$$

$$\begin{array}{c}
 \updownarrow \\
 \left(\begin{array}{cccc} u^c & u^c & u^c & N^c \\ d^c & d^c & d^c & e^+ \end{array} \right)_L \\
 \updownarrow \\
 (\bar{4}, 1, 2)_L
 \end{array}$$

||
 $(4, 2, 1)_L$

NOTE: $B-L = \lambda_{15} = \begin{pmatrix} 1/3 & 1/3 & 1/3 & -1 \\ & & & \end{pmatrix}$ of $SU(4)_c$

$$\begin{aligned}
 Q &= I_{3L} + I_{3R} + \frac{1}{2}(B-L) \\
 Q &= I_{3L} + Y/2
 \end{aligned}$$

Higgs of MSSM $\subset (1, 2, 2)$ "bidoublet"

$$\begin{array}{ccc}
 \begin{array}{c} \updownarrow SU(2)_L \\ \left[\begin{array}{cc} H_u & H_d \end{array} \right] = \begin{pmatrix} v_u & 0 \\ 0 & v_d \end{pmatrix} \\ \leftarrow SU(2)_R \rightarrow \end{array} & & \begin{array}{l} H_u = (1, 2, 1/2) \\ H_d = (1, 2, -1/2) \end{array}
 \end{array}$$

Yukawa terms in minimal Pati Salam $Y_{AB} (4, 2, 1)_{LA} (\bar{4}, 1, 2)_{LB} (1, 2, 2)_H$

$$\Rightarrow M_N = M_u \propto M_D = M_L \quad (\text{not necess. symmetric})$$

(C) $SO(2N)$ GROUP THEORY:

2.4

VECTORS AND TENSORS

A VECTOR HAS $2N$ REAL COMPONENTS, SO WE DO NOT DISTINGUISH UPPER AND LOWER INDICES. CALL VECTOR INDICES i, j, k , etc.

$$T^i = \begin{pmatrix} T^1 \\ T^2 \\ \vdots \\ T^{2N} \end{pmatrix}$$

TRANSFORMATIONS THAT LEAVE $\sum T^i T^i$ INVARIANT ARE REAL UNITARY (\equiv ORTHOGONAL) $2N \times 2N$ MATRICES.

$$O^T O = I$$

$$O = e^{i \sum_a \theta^a \lambda^a}$$

$\Rightarrow \lambda^a$ are purely imaginary, hermitian (= anti-symmetric) matrices.

$\Rightarrow a = 1, \frac{1}{2}(2N)(2N-1) \Rightarrow SO(10)$ has 45 generators.
 $i = 1, 2N$

ADJOINT REP:

$$T^{ij} = \sum_a T^a (\lambda^a)^{ij}$$

(USEFUL TO WRITE THE ADJOINT INDEX a AS A PAIR OF VECTOR INDICES (ANTISYMMETRIZED). i.e. LABEL

GENERATORS $\lambda^{[ij]}$ = GEN. OF ROTATION IN ij PLANE

$$(\lambda^{[ij]})^{kl} = -i [\delta^{ik} \delta^{jl} - \delta^{il} \delta^{jk}]$$

FOR EXAMPLE:

2.5

$$\lambda^{[12]} = \left[\begin{array}{cc|cc} 0 & -i & & \\ +i & 0 & & \\ \hline & & 0 & \\ & & & 0 \end{array} \right] \Rightarrow e^{i\theta \lambda^{[12]}} = \left[\begin{array}{cc|cc} \cos\theta & \sin\theta & & \\ -\sin\theta & \cos\theta & & \\ \hline & & 1 & \\ & & & 1 \end{array} \right]$$

IT IS EASY TO SEE THAT $SO(2N) \supset U(N)$
 $= SU(N) \times U(1)$.

$$-i = \begin{pmatrix} T^1 \\ T^2 \\ \vdots \\ T^{2N} \end{pmatrix} \rightarrow \begin{pmatrix} T^1 + iT^2 \\ T^3 + iT^4 \\ \vdots \\ T^{2N-1} + iT^{2N} \end{pmatrix} \oplus \begin{pmatrix} T^1 - iT^2 \\ T^3 - iT^4 \\ \vdots \\ T^{2N-1} - iT^{2N} \end{pmatrix}$$

\parallel \parallel
 T^α T_α
 $\alpha = 1, \dots, N$ $\alpha = 1, \dots, N$

So T^α of $SU(N)$ = $\frac{1}{2}(T^{(2\alpha-1)} + iT^{(2\alpha)})$ of $SO(2N)$ $\equiv \frac{1}{2} T^{(2\alpha-1) + i(2\alpha)}$
 T_α = $\frac{1}{2}(T^{(2\alpha-1)} - iT^{(2\alpha)})$ $\equiv \frac{1}{2} T^{(2\alpha-1) - i(2\alpha)}$

$$\sum_{i=1}^{2N} T^i T^i = 4 \sum_{\alpha=1}^N T^\alpha T_\alpha$$

So $U(N)$ TRANSFORMATIONS THAT KEEP $\sum T^\alpha T_\alpha$ INVARIANT ARE ALSO $SO(2N)$ TRANSFORMATIONS.

For example, Cartan generators of $SO(2N)$ form

$$-i \begin{bmatrix} a_1 & & & \\ -a_1 & & & \\ & a_2 & & \\ & -a_2 & & \\ & & \dots & \\ & & & a_N \\ & & & -a_N \end{bmatrix} \text{ corresponds to } \begin{bmatrix} a_1 & & & \\ & a_2 & & \\ & & \dots & \\ & & & a_N \end{bmatrix} \text{ of } U(N)$$

eg. For $SO(2) = U(1)$

$$-i \begin{bmatrix} & 1 \\ -1 & \end{bmatrix} \text{ equals } (1) \text{ of } U(1)$$

$$e^{i\theta} \begin{pmatrix} i & \\ & -i \end{pmatrix} = \begin{pmatrix} \cos\theta & \sin\theta \\ -\sin\theta & \cos\theta \end{pmatrix} \text{ equals } e^{i\theta} \text{ of } U(1)$$

We have seen $\left\{ \begin{array}{l} \text{vector rep } T^i \\ \text{adjoint rep } T^{[ij]} \end{array} \right. = \mathfrak{a}$

Of course there are an infinitely many types of tensor reps.

eg. $T^{(ij)}$ = traceless symmetric rank 2 tensor

$T^{[ijk]}$ etc.

(D) $SO(2N)$ GROUP THEORY:

2.7

SPINOR REPRESENTATIONS

ALGEBRA OF $SO(2N)$

$$[\lambda^{[ij]}, \lambda^{[kl]}] = i \left(\delta^{ik} \lambda^{[jl]} - \delta^{il} \lambda^{[jk]} - \delta^{jk} \lambda^{[il]} + \delta^{jl} \lambda^{[ik]} \right)$$

eg. $[\lambda^{[23]}, \lambda^{[31]}] = -i \lambda^{[21]} = i \lambda^{[12]}$

i.e. $[\lambda^1, \lambda^2] = i \lambda^3$

WE CAN FIND $2^N \times 2^N$ MATRICES THAT SATISFY THIS ALGEBRA.

FIRST, CONSTRUCT $2^N \times 2^N$ MATRICES

$$(\gamma^i)^\lambda_{\lambda'} \quad \lambda, \lambda' = 1, \dots, 2^N$$
$$i = 1, \dots, 2N$$

THAT SATISFY "CLIFFORD ALGEBRA"

$$\{\gamma^i, \gamma^j\} = 2 \delta^{ij} I$$

THEN DEFINE

$$\sigma^{[ij]} \equiv \frac{i}{2} [\gamma^i, \gamma^j]$$

THESE OBEY $SO(2N)$ GROUP ALGEBRA (LEFT AS EXERCISE)

THEN ELEMENTS OF GROUP CAN BE REPRESENTED BY $2^N \times 2^N$ MATRICES

2.8

$$U = e^{i \sum_j \theta^j \sigma^j}$$

THESE ARE SPINOR REPRESENTATION. THESE MATRICES ACT ON 2^N -COMPONENT "SPINORS" (CALLED BY PHYSICISTS SPINOR REPRESENTATIONS).

THESE SPINORS CAN BE WRITTEN :

$$\psi^\lambda = |s_1 s_2 s_3 \dots s_N\rangle$$

where $s_\alpha = \pm 1$.

∴ WILL WRITE

$$\gamma^{2k-1} = \underbrace{I \otimes I \otimes \dots \otimes I}_{k-1} \otimes \underbrace{\tau_1 \otimes \tau_3 \otimes \dots \otimes \tau_3}_{N-k}$$

$$\gamma^{2k} = -\underbrace{I \otimes I \otimes \dots \otimes I}_{k-1} \otimes \tau_2 \otimes \underbrace{\tau_3 \otimes \dots \otimes \tau_3}_{N-k}$$

THESE ARE $2N$ HERMITIAN $2^N \times 2^N$ MATRICES. CAN CHECK THAT THEY SATISFY CLIFFORD ALGEBRA.

NOTE: γ^1, γ^2 FLIP FIRST "SPIN" s_1
 γ^3, γ^4 FLIP SECOND "SPIN" s_2
 \vdots
 $\gamma^{2N-1}, \gamma^{2N}$ FLIP N^{TH} "SPIN" s_N

CHECK CLIFFORD ALGEBRA SATISFIED.

2.9

$$\begin{cases} \{\gamma^i, \gamma^i\} = 2\delta^{ii}I = 2I \Leftrightarrow (\gamma^i)^2 = I \\ (I \otimes I \otimes \dots \otimes \tau_2 \otimes \tau_3 \otimes \dots \otimes \tau_3)^2 = I \checkmark \end{cases}$$

$$\begin{cases} \left\{ \begin{array}{l} \frac{I}{\tau_i} \frac{\tau_3}{\tau_j} \\ \frac{I}{\tau_j} \frac{\tau_3}{\tau_i} \end{array} \right\} \\ = \frac{I}{\tau_i} \frac{\tau_3}{\tau_j} \{\tau_3, \tau_j\} \frac{I}{\tau_i} = 0 \checkmark \end{cases}$$

$$\begin{cases} \left\{ \begin{array}{l} \frac{I}{\tau_i} \frac{\tau_3}{\tau_j} \\ \frac{I}{\tau_j} \frac{\tau_3}{\tau_i} \end{array} \right\} \quad \begin{array}{l} i \neq j \\ ij=1,2 \end{array} \\ = \frac{I}{\tau_i} \{\tau_i, \tau_j\} \frac{I}{\tau_j} = 0 \end{cases}$$

LATER WE WILL LOOK AT $\sigma^{ij} = \frac{i}{2} [\gamma^i, \gamma^j]$.

BUT NOTICE THAT σ^{ij} FLIPS TWO SPINS
(UNLESS $ij = (2k-1)(2k)$, WHEN FLIPS ZERO SPINS)

$$\begin{aligned} \text{THUS } |s_1 \dots s_N\rangle &= |s_1 \dots s_N\rangle_{\tau_s = +1} \oplus |s_1 \dots s_N\rangle_{\tau_s = -1} \\ &= \Psi_+ \oplus \Psi_- \\ &= \text{TWO IRREDUCIBLE SPINORS.} \end{aligned}$$

CONJUGATION: $U \rightarrow U^* \Rightarrow e^{i\theta^a \lambda^a} \rightarrow e^{-i\theta^a \lambda^{a*}}$

$\Rightarrow \lambda^a \rightarrow -\lambda^{a*} \Rightarrow \left(\frac{i}{2} [\gamma^i, \gamma^j]\right) \rightarrow \left(\frac{i}{2} [\gamma^{i*}, \gamma^{j*}]\right)$

$\Rightarrow \gamma^i \rightarrow \gamma^{i*} \Rightarrow \tau_1 \rightarrow \tau_1, \tau_2 \rightarrow -\tau_2$

$\Rightarrow \tau^\pm \rightarrow \tau^\mp \Rightarrow \text{SPIN UP} \leftrightarrow \text{SPIN DOWN}$

$\Rightarrow \underline{N = \text{even}} \quad \overline{\Psi}_+ = \Psi_+, \quad \overline{\Psi}_- = \Psi_-$

$\underline{N = \text{odd}} \quad \overline{\Psi}_+ = \Psi_-, \quad \overline{\Psi}_- = \Psi_+$

TABLE OF SPINORS FOR SO(2N)

<u>Equivalent GP.</u>	<u>SO(2N)</u>	<u>Ψ_+</u>	<u>Ψ_-</u>	<u>vector</u>
$U(1)$	SO(2)	1	$\bar{1}$	$2 = 1 + \bar{1}$
$U(2) = SU(2)$	SO(4)	$2^{(2,1)}$	$2'^{(1,2)}$	$4^{(2,2)}$
$SU(4)$	SO(6)	$4 = T^{\alpha}$	$\bar{4} = T_{\alpha}$	$6 = T^{[\alpha\beta]}$
	SO(8)	8	$8'$	8
	SO(10)	16	$\bar{16}$	10
	SO(12)	32	$32'$	12
	SO(14)	64	$\bar{64}$	14
	SO(16)	128	$128'$	16
	SO(18)	256	$\bar{256}$	18

D.2 Family Unification (See Wilczek + Zee "Families from spinors")

$$SO(16) \supset SO(10) \times SO(6)$$

$$128_L \longrightarrow (16, 4)_L + (\overline{16}, \overline{4})_L$$

4 families

4 mirror families

$$\overline{16}_L \sim 16_R$$

(V-A) Weak int.

(V+A) Weak int.

128 is self-conjugate (i.e. a "real representation")

So can have $M(128_L, 128_L)$

"Naturally", 128_L would be superheavy unless some global symmetry prevents this mass term.
(or local)

$$SO(18) \supset SO(10) \times SO(8)$$

$$256_L \longrightarrow (16, 8)_L + (\overline{16}, 8')_L$$

256 not self-conjugate, so $M(256, 256)$ not possible

Problem with such family unification schemes (also with E_8) is how to make V+A families heavy. Can be solved in higher dimensions. [KS Babu, S.M.B., and B. Kyae hep-ph/0202178]

D.3 How SO(10) CONTAINS SU(5) MODEL

2.13

We know that a family

$$= 16 \text{ of } SO(10)$$

$$= (4, 2, 1) + (\bar{4}, 1, 2) \text{ of Pati-Salam}$$

$$= 10 + \bar{5} + 1 \text{ of } SU(5)$$

We can see that under $SO(10) \rightarrow SU(5)$
 $16 \rightarrow 10 + \bar{5} + 1$
directly, by looking more closely at spinor representations of $SO(2N)$.

SO(2N)

$$\lambda^{[ij]} \equiv \sigma^{ij} = \frac{i}{2} [\gamma^i, \gamma^j]$$

Recall that $U(N) = SU(N) \times U(1)$ embedded in $SO(2N)$ as follows:

$$T^\alpha = \frac{1}{2} T^{(2\alpha-1) + i(2\alpha)} = \frac{1}{2} (T^{(2\alpha-1)} + i T^{(2\alpha)})$$

$$\underbrace{T_\alpha}_{U(N)} = \frac{1}{2} \underbrace{T^{(2\alpha-1) - i(2\alpha)}}_{SO(2N)} = \frac{1}{2} (T^{(2\alpha-1)} - i T^{(2\alpha)})$$

SO $SU(N) \times U(1)$ GENERATORS CAN BE WRITTEN^{2.14}
AS $SO(2N)$ GENERATORS THUS:

$$\begin{aligned} \lambda_{\beta}^{\alpha} &= -\frac{i}{4} \sigma[(2\alpha-1)+i(2\alpha), (2\beta-1)-i(2\beta)] \\ &= -\frac{1}{8} [\gamma^{2\alpha-1} + i\gamma^{2\alpha}, \gamma^{2\beta-1} - i\gamma^{2\beta}] \\ &= +\frac{1}{8} \left[\underbrace{I \otimes \dots \otimes I}_{\alpha^{\text{th}} \text{ place} = 2\gamma^{-}} \otimes (\tau_1 - i\tau_2) \otimes \tau_3 \otimes \dots \otimes \tau_3, \right. \\ &\quad \left. I \otimes \dots \otimes I \otimes (\tau_1 + i\tau_2) \otimes \tau_3 \otimes \dots \otimes \tau_3 \right] \\ &\quad \underbrace{\hspace{10em}}_{\beta^{\text{th}} \text{ place} = 2\gamma^{+}} \end{aligned}$$

\Rightarrow for $\alpha < \beta$

$$\begin{aligned} \lambda_{\beta}^{\alpha} &= +\frac{1}{8} (I \otimes \dots \otimes I \otimes \underbrace{2\tau^{+}}_{\alpha^{\text{th}}} \otimes \tau_3 \otimes \dots \otimes \underbrace{[\tau_3, 2\tau^{+}]}_{\beta^{\text{th}}} \otimes I \otimes \dots \otimes I) \\ &= (I \otimes \dots \otimes I \otimes \underbrace{\tau^{-}}_{\alpha^{\text{th}}} \otimes \tau_3 \otimes \dots \otimes \underbrace{\tau^{+}}_{\beta^{\text{th}}} \otimes I \otimes \dots \otimes I) \end{aligned}$$

SO λ_{β}^{α} raises β^{th} spin and lowers α^{th} spin.

For $\alpha = \beta$

$$\lambda_{\alpha}^{\alpha} = \frac{1}{2} (I \otimes \dots \otimes I \otimes \underbrace{\tau_3}_{\alpha^{\text{th}}} \otimes I \otimes \dots \otimes I) = -\frac{S}{2}$$

(no sum)

$$\begin{pmatrix} a_1 \\ a_2 \\ \dots \\ a_N \end{pmatrix} = \sum_{\alpha=1}^N a_{\alpha} \lambda_{\alpha}^{\alpha} = -\frac{1}{2} \sum_{\alpha=1}^N a_{\alpha} S_{\alpha}$$

CONSIDER SPINOR $\Psi_+ = 16$ OF $SO(10)$. 2.16
(NO 2.15)
 $= |s_1 s_2 s_3 s_4 s_5\rangle_{\prod s_i = +1}$

THIS CONTAINS

$$\left[\begin{array}{l} \{ |++++\rangle \} \longrightarrow 1 \text{ component} \\ \{ |+++--\rangle, \text{ permutations} \} \longrightarrow 10 \text{ components} \\ \{ |+----\rangle, \text{ permutations} \} \longrightarrow 5 \text{ components} \end{array} \right.$$

SINCE $SU(5) \times U(1)$ GENERATORS DO NOT CHANGE NUMBER OF + SIGNS \Rightarrow THESE THREE TYPES DO NOT MIX WITH EACH OTHER UNDER $SU(5) \times U(1)$:
 IRREDUCIBLE REPS.

$$\lambda'_2 | \underbrace{+----}_{T_1} \rangle = | \underbrace{-+----}_{T_2} \rangle$$

$$\underbrace{\sum_{\alpha} a_{\alpha} \lambda'_{\alpha}}_{SU(5) \text{ GEN}} | +---- \rangle = -\frac{1}{2} (a_1 - a_2 - a_3 - a_4 - a_5) | \rangle = -a_1 | \rangle \quad (\text{since } \sum a_{\alpha} = 0 \text{ for } su(5))$$

$$\Rightarrow \begin{aligned} \{ | +---- \rangle, \text{ permutations} \} &= \bar{5} \text{ of } su(5) \\ \{ | +++-- \rangle, \text{ permutations} \} &= 10 \text{ " } \\ \{ | +++++ \rangle \} &= 1 \end{aligned}$$

$$\sum_{\alpha} a_{\alpha} \lambda'_{\alpha} | +++-- \rangle = -\frac{1}{2} (a_1 + a_2 + a_3 - a_4 - a_5) = -(a_4 + a_5) | \rangle = -(a_4 + a_5) | \rangle \Rightarrow | \rangle = T^{45} = T_{45}$$

CONSIDER THE $U(1)$ GENERATOR

$$X \equiv -2 \sum_{\alpha=1}^5 \lambda^\alpha \alpha = -2 \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} = \sum_{\alpha=1}^5 S_\alpha$$

↑
a common normalization

$$\left. \begin{aligned} X |++++\rangle &= 5 |++++\rangle \\ X |+++--\rangle &= 1 |+++--\rangle \\ X |+----\rangle &= -3 |+----\rangle \end{aligned} \right\} \Rightarrow 16 = 1^5 + 10^1 + 5^{-3}$$

D.4 ANOTHER LOOK AT B-L

PATI - SALAM: $SO(4) \times SO(6) \subset SO(10)$

$$[SU(2)_L \times SU(2)_R] \times SU(4)_C$$

$$\cup \quad \cup$$

$$[SU(2)_L \times U(1)_{I_{3R}}] \times [SU(3)_C \times U(1)_{B-L}]$$

LOOK AT $U(3) = SU(3)_C \times U(1)_{B-L}$ GENERATOR

EMBEDDED IN $SO(6)$ AS DESCRIBED ABOVE:

$$B-L = \frac{-2}{3} \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} = \frac{-2}{3} (\lambda^3_3 + \lambda^4_4 + \lambda^5_5) = \frac{1}{3} (S_3 + S_4 + S_5)$$

$$\Rightarrow \begin{cases} (B-L) |+++ \rangle = \frac{1}{3}(1+1+1) = +1 & 1 \text{ of color} \\ (B-L) |+- \rangle = \frac{1}{3}(1-1-1) = -\frac{1}{3} & \bar{3} \text{ of color} \end{cases}$$

$$\text{So } \Psi_+ \text{ of } SO(6) = \bar{4} \text{ (fundamental of } SU(4)_C)$$

$$= \underbrace{\bar{3}^{-1/3}}_{\text{antiquark}} + \underbrace{1^+}_{\text{antilepton}}$$

AS A $U(5)$ GENERATOR: $B-L = -\frac{2}{3} \begin{pmatrix} 0 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix}$

The Higgs of $SU(5)$ model are

$$\text{in } \bar{5} + 5 = \begin{pmatrix} H_d \\ H_{dc} \end{pmatrix} + \begin{pmatrix} H_u \\ H_{uc} \end{pmatrix}$$

These combine into a 10 of $SO(10)$.

$$X = -2 \begin{pmatrix} 1 & & & & \\ & 1 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} \Rightarrow 10 = \bar{5}^{+2} + 5^{-2}$$

$$B-L = -\frac{2}{3} \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & 1 & & \\ & & & 1 & \\ & & & & 1 \end{pmatrix} \Rightarrow \begin{array}{l} H_d, H_u \text{ have } B-L = 0 \\ H_{dc} \text{ has } B-L = \frac{2}{3} \\ H_{uc} \text{ has } B-L = -\frac{2}{3} \end{array}$$

This is important for later discussion of $\frac{2}{3}$ splitting.

$$\text{If } M_{\text{Higgs}} \propto B-L$$

$$\Rightarrow \begin{cases} M(H_u, H_d) = 0 \\ M(H_{uc}, H_{dc}) \neq 0 \end{cases}$$

2.5 COUPLING SPINORS TO VECTORS AND TENSORS

Define $\gamma^{[ij \dots k]} \equiv (\gamma^i \gamma^j \dots \gamma^k)$ antisymmetrized.

γ^i FLIPS ONE "SPIN." $\Rightarrow \gamma^{[ij \dots k]}$ with even (odd) number of indices flips even (odd) number of spins.

$C \equiv$ charge conjugation matrix. C flips all spins.

We can construct an $SO(2N)$ invariant as follows:

$$\Psi^T C (\gamma^{[ij \dots k]}) \Psi \quad T^{[ij \dots k]}$$

[Note: if $\Psi \rightarrow e^{i\theta \sigma} \Psi$, then $(\Psi^T C) \rightarrow (\Psi^T C) e^{-i\theta \sigma}$.

Also only totally antisymmetrized product of γ^i need to be considered, since symmetric product $\gamma^{(i} \gamma^{j)} = \delta^{ij} I$ by Clifford algebra.]

For N even: $\begin{cases} T^{[ij \dots k]} \text{ with } \underline{\text{even rank}} \text{ couples to } \Psi_{\pm}^T \sigma \Psi_{\pm} \\ T^{[ij \dots k]} \text{ with } \underline{\text{odd rank}} \text{ couples to } \Psi_{\pm}^T \sigma \Psi_{\mp} \end{cases}$

For N odd: $\begin{cases} T^{[ij \dots k]} \text{ with } \underline{\text{even rank}} \text{ couples to } \Psi_{\pm}^T \sigma \Psi_{\mp} \\ T^{[ij \dots k]} \text{ with } \underline{\text{odd rank}} \text{ couples to } \Psi_{\pm}^T \sigma \Psi_{\pm} \end{cases}$

Define $\gamma_{\text{FIVE}} \equiv \underbrace{\gamma^{[i_1 i_2 \dots i_{2N}]}}_{2N \text{ factors}} = \gamma^1 \gamma^2 \gamma^3 \dots \gamma^{2N}$

$$= (\tau_3 \otimes \dots \otimes \tau_3)$$

$$\gamma_{\text{FIVE}} \Psi_{\pm} = \pm \Psi_{\pm}$$

$$\underbrace{\gamma^{[i_1 i_2 \dots i_p]}}_p = \pm \gamma_{\text{FIVE}} \underbrace{\gamma^{[i_{p+1} i_{p+2} \dots i_{2N}]}}_{2N-p}$$

$$\Rightarrow \left[\begin{array}{l} \Psi_{\pm}^T \quad C [T + \underbrace{\gamma^{[ij]} T^{[ij]}}_{-2i \sigma^{ij} \text{ adjoint}} + \dots + \underbrace{\gamma^{(i \dots k)} T^{(i \dots k)}}_{\substack{\text{largest even} \\ \leq N}}] \Psi_{\pm} \\ \pm (N \text{ even}) \\ \mp (N \text{ odd}) \end{array} \right. \left. \begin{array}{l} \Psi_{\pm}^T \quad C [\gamma^i T^i + \gamma^{[ijk]} T^{[ijk]} + \dots + \underbrace{\gamma^{(i \dots k)} T^{(i \dots k)}}_{\substack{\text{largest odd} \\ \leq N}}] \Psi_{\pm} \\ \mp (N \text{ even}) \\ \pm (N \text{ odd}) \end{array} \right]$$

EXAMPLE: SO(10)

$$\left\{ \begin{array}{l} \Psi_+ C (T + \gamma^{[ij]} T^{[ij]} + \gamma^{[ijkl]} T^{[ijkl]}) \Psi_- \\ 16 \quad (1 + 45 + 210) \overline{16} \end{array} \right.$$

$$\left\{ \begin{array}{l} \Psi_{\pm} C (\gamma^i T^i + \gamma^{[ijk]} T^{[ijk]} + \gamma^{[ijklm]} T^{[ijklm]}) \Psi_{\pm} \\ 16 \quad (10 + 120 + \overline{126}) 16 \\ (\overline{16}) \quad (10 + 120 + 126) (\overline{16}) \end{array} \right.$$

Note $T^{[ijklm]} = \pm \epsilon^{ijklm i' j' k' l' m'} T^{[i' j' k' l' m']}$

$$\text{Rank 5 tensor} = 252 = \underbrace{126 + \overline{126}}_{\text{irreducibly tensors}}$$

IN MINIMAL SO(10) MODEL, FERMION MASS TERMS

$$Y_{AB} (16_{LA} 16_{LB}) 10_H$$

$$\begin{aligned} \rightarrow Y_{AB} & \left(\begin{array}{l} 10_A^1 \quad 10_B^1 \quad 5_H^{-2} \\ + 10_A^1 \quad \overline{5}_B^{-3} \quad \overline{5}_H^{+2} \\ + \overline{5}_A^{-3} \quad 10_B^1 \quad \overline{5}_H^{+2} \\ + \overline{5}_A^{-3} \quad \overline{15}_B^5 \quad 5_H^{-2} \\ + 10_A^1 \quad \overline{5}_B^{-3} \quad 5_H^{-2} \end{array} \right) \rightarrow M_u \\ & \hspace{15em} \rightarrow M_d, M_e \\ & \hspace{15em} \rightarrow M_\nu \end{aligned}$$

THE MAJORANA MASS MATRIX OF THE RIGHT-HANDED NEUTRINOS, CAN COME FROM:

$$(Y_R)_{AB} (16_{L_A} / 16_{L_B}) \overline{126}_H \rightarrow M_R$$

$\langle 1^{-10} \rangle \sim M_{\text{GUT}}$

OR A HIGHER-DIMENSION OPERATOR:

$$(Y_R)_{AB} (16_{L_A} / 16_{L_B}) \overline{16}_H \overline{16}_H / M_G$$

$\langle 1^{-5} \rangle \langle 1^{-5} \rangle \sim M_{\text{GUT}}$

THUS, IN MINIMAL $SO(10)$ ONE HAS

$$\underbrace{M_N = M_U \propto M_D = M_L}_{\text{SYMMETRIC}}, \quad \underbrace{M_R}_{\text{SYMMETRIC}}$$

IN MINIMAL PATI-SALAM, ONE HAS SAME RELATIONS, BUT THE DIRAC MATRICES NEED NOT BE SYMMETRIC.

IN MINIMAL $SU(5)$

$$M_N, \quad \underbrace{M_U}_{\text{SYMMETRIC}}, \quad M_D = M_L^T, \quad \underbrace{(M_R)}_{\text{SYMMETRIC}}$$

IN MINIMAL $SO(10)$ (OR PATI SALAM) $M_\nu = M_U M_R^{-1} M_U^T$

$$\Rightarrow \text{(FOR ONE FAMILY MODEL)} \quad m_3 = \frac{m_\mu^2}{M_R}$$

$$M_R = \frac{m_\mu^2}{m_3} = \frac{(174 \text{ GeV})^2}{0.06 \text{ eV}} = \frac{1}{2} \times 10^{15} \text{ GeV}$$


Close to GUT scale!

NOTE ALSO THAT MINIMAL $SO(10) \Rightarrow \theta_{\text{CKM}} = 0$. (Approx. correct)

(E) BREAKING $SO(10)$ TO SM.NEED TWO KINDS OF HIGGS TO DO IT.

$$SO(10) \supset SU(5) \times U(1)_X$$

$$\left\{ \begin{array}{l} 45 \rightarrow 24^0 + 1^0 + 10^{-4} + \overline{10}^4 \\ 16 \rightarrow 10^1 + \overline{5}^{-3} + 1^5 \\ \overline{16} \rightarrow \overline{10}^{-1} + 5^3 + 1^{-5} \\ 10 \rightarrow 5^{-2} + \overline{5}^2 \end{array} \right.$$



We know that an $SU(5)$ adjoint Higgs, 24_H , can be used to break $SU(5) \rightarrow SM$.

In $SO(10)$, an adjoint Higgs is not enough.

The 10^{-4} and $\overline{10}^4$ in a 45_H cannot get VEVs $\sim M_G$, since all components of them are electrically charged. (Even in "flipped $SU(5)$ ".)

And 24^0 and 1^0 leave $U(1)_X$ unbroken, as well as $SU(3) \times SU(2) \times U(1) \subset SU(5)$.

Need another Higgs to break $U(1)_X$

Usual choices $16_H + \overline{16}_H$ or $126_H + \overline{126}_H$

(Need both 126 and $\overline{126}$, or both 16 and $\overline{16}$, so as not to break SUSY at M_G by D terms:

$$\langle D^a \rangle = \left\langle \sum_i \Phi_i^\dagger \lambda^a \Phi_i \right\rangle$$

We saw that $16 \supset 1^5$, $\overline{16} \supset 1^{-5}$
and $126 \supset 1^{10}$, $\overline{126} \supset 1^{-10}$

and that these can give M_R . They also break $U(1)_X$.

So:

45_H (and/or $54_{H, \dots}$) \longrightarrow Breaks $SO(10)$ to rank 5 subgroup
eg. $SU(3)_L \times SU(2)_L \times U(1)_Y \times U(1)_X$ Does 2/3 splitting

$16_H + \overline{16}_H$ \longrightarrow Breaks $U(1)_X$ ("breaks rank") Gives mass to N^c (i.e. ν_R)
or

$126_H + \overline{126}_H$

E.2 "Flipped SU(5)" [DeRujck, Georgi, Glashow 1980 2.24
S.M.B. 1982]

$U(1)_{em}$ can be embedded purely in $SU(5)$ (Georgi-Glashow) or partly in $SU(5)$ and partly in $U(1)_X$ (Flipped $SU(5)$).

$$SO(10) \supset SU(5) \times U(1)_X \\ \supset [SU(3)_C \times SU(2)_L \times U(1)_{Y_5}] \times U(1)_X$$

Let $\boxed{Y/2 = \alpha Y_5/2 + \beta X}$ $U(1)_Y$
($\alpha=1, \beta=0 \Rightarrow$ G.G.)

rep $[SU(3)_C \times SU(2)_L \times U(1)_{Y_5}]^{U(1)_X}$	G-G	Flipped	$Y/2$ $\alpha Y_5/2 + \beta X$
$(1, 1, +1)^1$	e_L^+	N^c	$\alpha + \beta = 0, 1$
$(3, 2, 1/6)^1$	$Q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix}$	$Q_L = \begin{pmatrix} u_L \\ d_L \end{pmatrix}$	$\boxed{\frac{1}{6}\alpha + \beta = \frac{1}{6}}$
$(\bar{3}, 1, -2/3)^1$	u_L^c	d_L^c	$\frac{-2}{3}\alpha + \beta = +1/3, -2/3$
$(1, 2, -1/2)^{-3}$	$L_L = \begin{pmatrix} \nu_L \\ e_L^- \end{pmatrix}$	$L_L = \begin{pmatrix} \nu_L \\ e_L^- \end{pmatrix}$	$\boxed{\frac{-1}{2}\alpha - 3\beta = \frac{-1}{2}}$
$(\bar{3}, 1, 1/3)^{-3}$	d_L^c	u_L^c	$\frac{1}{3}\alpha - 3\beta = -2/3, 1/3$
$(1, 1, 0)^5$	N_L^c	e_L^+	$0\alpha + 5\beta = 1, 0$

$$\left. \begin{aligned} \alpha + \beta &= 0 \\ \frac{1}{6}\alpha + \beta &= \frac{1}{6} \end{aligned} \right\} \Rightarrow \alpha = -1/5, \beta = 1/5$$

$$\boxed{Y/2 = +\frac{1}{5}(-Y_5/2 + X)}$$

Flipped

Check: $\frac{-2}{3}\alpha + \beta = \frac{-2}{3}(-\frac{1}{5}) + \frac{1}{5} = \frac{1}{3} \checkmark$
 $\frac{1}{3}\alpha - 3\beta = \frac{1}{3}(-\frac{1}{5}) - 3\frac{1}{5} = -\frac{2}{3} \checkmark$

HIGGS IN FLIPPED SU(5)

$$(3_c 2_L \frac{1}{2})^X \quad \frac{1}{2} = \frac{-1}{5}(\frac{1}{2}) + \frac{1}{5}X \quad (3_c 2_L \frac{1}{2})$$

$10'$	$\left\{ \begin{array}{l} (1, 1, +1)^1 \\ (3, 2, \frac{1}{6})^1 \\ (\bar{3}, 1, -\frac{2}{3})^1 \end{array} \right.$	$0 = -\frac{1}{5}(1) + \frac{1}{5}(1)$	$\left\{ \begin{array}{l} \langle (1, 1, 0) \rangle \neq 0 \\ (3, 2, \frac{1}{6}) \\ (\bar{3}, 1, \frac{1}{3}) \\ (1, 2, -\frac{1}{2}) \\ (\bar{3}, 1, -\frac{2}{3}) \\ (1, 1, +1) \end{array} \right.$
		$\frac{1}{6} = -\frac{1}{5}(\frac{1}{6}) + \frac{1}{5}(1)$	
$\frac{1}{5}^{-3}$	$\left\{ \begin{array}{l} (1, 2, -\frac{1}{2})^{-3} \\ (\bar{3}, 1, \frac{1}{3})^{-3} \end{array} \right.$	$\frac{1}{3} = -\frac{1}{5}(-\frac{1}{3}) + \frac{1}{5}(1)$	$\left\{ \begin{array}{l} (1, 2, -\frac{1}{2}) \\ (3, 1, -\frac{1}{3}) \\ (1, 2, \frac{1}{2}) \\ (\bar{3}, 1, \frac{1}{3}) \end{array} \right.$
		$-\frac{1}{2} = -\frac{1}{5}(-\frac{1}{2}) + \frac{1}{5}(-3)$	
1^5	$\left\{ \begin{array}{l} (1, 1, 0)^5 \end{array} \right.$	$-\frac{2}{3} = -\frac{1}{5}(\frac{1}{3}) + \frac{1}{5}(-3)$	$\left\{ \begin{array}{l} (1, 1, +1) \end{array} \right.$
		$1 = -\frac{1}{5}(0) + \frac{1}{5}(5)$	
5^{-2}	$\left\{ \begin{array}{l} (1, 2, \frac{1}{2})^{-2} \\ (3, 1, -\frac{1}{3})^{-2} \end{array} \right.$	$-\frac{1}{2} = -\frac{1}{5}(\frac{1}{2}) + \frac{1}{5}(-2)$	$\left\{ \begin{array}{l} (1, 2, -\frac{1}{2}) \\ (3, 1, -\frac{1}{3}) \\ (1, 2, \frac{1}{2}) \\ (\bar{3}, 1, \frac{1}{3}) \end{array} \right.$
		$-\frac{1}{3} = -\frac{1}{5}(-\frac{1}{3}) + \frac{1}{5}(-2)$	
5^2	$\left\{ \begin{array}{l} (1, 2, -\frac{1}{2})^2 \\ (\bar{3}, 1, \frac{1}{3})^2 \end{array} \right.$	$\frac{1}{2} = -\frac{1}{5}(\frac{1}{2}) + \frac{1}{5}(2)$	$\left\{ \begin{array}{l} (1, 2, \frac{1}{2}) \\ (\bar{3}, 1, \frac{1}{3}) \end{array} \right.$
		$\frac{1}{3} = -\frac{1}{5}(\frac{1}{3}) + \frac{1}{5}(2)$	

So: $\left\{ \begin{array}{l} \langle 1^0 (45_H) \rangle : SO(10) \rightarrow SU(5) \times U(1)_X \\ \langle 10' (16_H) \rangle : SU(5) \times U(1)_X \rightarrow SM. \end{array} \right.$

IN "FLIPPED SU(5)" MODELS IT IS ASSUMED THAT STARTING GROUP IS $SU(5) \times U(1)_X$, NOT $SO(10)$. AS WE SHALL SEE, THIS HAS ADVANTAGES. BUT IT LEAVES UNEXPLAINED UNIFICATION OF GAUGE COUPLINGS.

LECTURE III: THE GAUGE HIERARCHY PROBLEM IN SUSY GUTS

A) NON-SUSY GUTS, SU(5)

Adjoint Higgs (24_H): $\langle \Omega^\alpha_\beta \rangle = \Omega \begin{pmatrix} \frac{1}{2} & & & & \\ & \frac{1}{3} & & & \\ & & -\frac{1}{3} & & \\ & & & -\frac{1}{3} & \\ & & & & -\frac{1}{3} \end{pmatrix} = \Omega \frac{1}{2}$
breaks $SU(5) \rightarrow SM$.

CONSIDER $V(5_H)$

$$V_{\text{tree}} = \lambda (5_H^\dagger 5_H)^2 + M^2 (5_H^\dagger 5_H) + \lambda' (\text{tr } 24_H 24_H) 5_H^\dagger 5_H + \lambda'' 5_H^\dagger 24_H 24_H 5_H + M' 5_H^\dagger 24_H 5_H$$

$$= \lambda (H_\alpha^\dagger H^\alpha)^2 + M^2 (H_\alpha^\dagger H^\alpha) + \lambda' (\Omega^\alpha_\beta \Omega^\beta_\alpha) H_\alpha^\dagger H^\alpha + \lambda'' H_\alpha^\dagger \Omega^\alpha_\beta \Omega^\beta_\gamma H^\gamma + M' H_\alpha^\dagger \Omega^\alpha_\beta H^\beta$$

$$\Rightarrow (H, H_c)^\dagger \left[\lambda'' \Omega^2 \begin{pmatrix} \frac{1}{4} I_2 & \\ & \frac{1}{4} I_3 \end{pmatrix} + (M^2 + \frac{5}{8} \lambda' \Omega^2) \begin{pmatrix} I_2 & \\ & I_3 \end{pmatrix} + M' \Omega \begin{pmatrix} \frac{1}{2} I_2 & \\ & \frac{1}{3} I_3 \end{pmatrix} \right] \begin{pmatrix} H \\ H_c \end{pmatrix}$$

$$\Rightarrow m^2(H) = \frac{1}{4} \lambda'' \Omega^2 + (M^2 + \frac{5}{8} \lambda' \Omega^2) + \frac{1}{2} M' \Omega \approx (10^2 \text{ GeV})^2$$

$$m^2(H_c) = \frac{1}{9} \lambda'' \Omega^2 + (M^2 + \frac{5}{8} \lambda' \Omega^2) - \frac{1}{3} M' \Omega \approx (10^{15} \text{ GeV})^2$$

THREE ASPECTS:

(a) Doublet-Triplet ($2/3$) Splitting problem
[WHY CANCELLATION?]

(b) Fine tuning problem:
[Why true to all orders?]

(c) Small number problem: [Where does $(\frac{10^2}{10^{15}})^2 = 10^{-26}$ come from?]

(B) SUSY SU(5)

NOTE: IN SUSY WE CAN LOOK AT MASSES OF THE FERMIONIC PARTNERS OF THE HIGGS DOUBLETS AND TRIPLETS (I.E. THE HIGGSINOS). THESE CAN BE READ OFF DIRECTLY FROM THE SUPERPOTENTIAL

$$W = \lambda \bar{5}_H 24_H 5_H + M \bar{5}_H 5_H + W(24_H) + \dots$$

NOTE: $\bar{5}_H$ AND 5_H ARE DISTINCT LEFT-HANDED CHIRAL SUPERFIELDS.

$$\Rightarrow \bar{5}_H (\lambda \langle 24_H \rangle + M) 5_H$$

↑ Higgsinos

$$\Rightarrow (\bar{2}_H \bar{3}_H) \left[\lambda \Omega \begin{pmatrix} \frac{1}{2} I_2 & \\ & -\frac{1}{3} I_3 \end{pmatrix} + M \begin{pmatrix} I_2 & \\ & I_3 \end{pmatrix} \right] \begin{pmatrix} 2_H \\ 3_H \end{pmatrix}$$


$$2 \equiv (1, 2, \frac{1}{2}), \quad \bar{2} \equiv (1, 2, -\frac{1}{2})$$

$$3 \equiv (3, 1, -\frac{1}{3}), \quad \bar{3} \equiv (\bar{3}, 1, \frac{1}{3})$$

$$\Rightarrow \begin{cases} \mu_2 = \frac{1}{2} \lambda \Omega + M \sim M_{\text{weak}} \\ \mu_3 = -\frac{1}{3} \lambda \Omega + M \sim M_G \end{cases}$$

μ_2 is just " μ parameter" of MSSM

μ_3 is the parameter that comes into p decay amplitude ($d=5$ operator)



(C) SUSY MECHANISMS TO SOLVE THE 2/3 SPLITTING PROBLEM

SUSY AUTOMATICALLY SOLVES THE "FINE-TUNING" PROBLEM, BECAUSE W IS NOT RENORMALIZED — no radiative corrections to it.

FOUR MECHANISMS HAVE BEEN PROPOSED TO SOLVE 2/3 SPLITTING PROBLEM IN SUSY GUTS:

1. "SLIDING SINGLET MECHANISM" [Litten 1981, Manopoulos + Tamvakis 1982]
2. "MISSING V_{EV} MECHANISM" [Dimopoulos + Wilczek 1981] ("DIMOPOULOS WILCZEK MECHANISM")
3. "MISSING PARTNER MECHANISM" [Georgi 1982, Masiero, Manopoulos, Tamvakis + Yanagida 1982]
4. "GIFT MECHANISM" [Berezhiani + Okuli 1989]

SLIDING SINGLET MECHANISM.

$$W = \lambda \bar{5}_H 24_H 5_H + \lambda' \bar{5}_H 1_H 5_H + M \bar{5}_H 5_H + W(24_H) + \dots$$

↑
absorb into 1_H

SINGLET WHICH HAS NO OTHER COUPLINGS THAT GIVE IT POTENTIAL

$$\Rightarrow V(1_H) = \left| (\lambda \langle 24_H \rangle + \lambda' \langle 1_H \rangle) \langle 5_H \rangle \right|^2 + \left| \langle \bar{5}_H \rangle (\lambda \langle 24_H \rangle + \lambda' \langle 1_H \rangle) \right|^2 + \text{SUSY terms.}$$

$$\langle 5_H \rangle = \begin{pmatrix} v_u \\ 0 \end{pmatrix} \Rightarrow V(1_H) = (\lambda \frac{1}{2} \Omega + \lambda' \langle 1_H \rangle) (|v_u|^2 + |v_d|^2) + \text{SUSY terms.}$$



$$\langle \bar{5}_H \rangle = \begin{pmatrix} v_d \\ 0 \end{pmatrix}$$

i.e. IN SUSY LIMIT,

$$V(1_H) = \mu_2 (|v_u|^2 + |v_d|^2) = 0$$

$$\Rightarrow \boxed{\mu_2 = 0}, \mu_3 \neq 0.$$

HOWEVER, IN SU(5) THE SLIDING SINGLET MECH. FAILS WHEN SUSY TAKEN INTO ACCOUNT.



$$\Rightarrow V(1_H) = |\mu_2(1_H) V|^2 + \frac{\lambda' \mu_3}{16\pi^2} m_{\text{SUSY}}^2 \cdot \delta 1_H$$

$$0 = \frac{\partial V}{\partial(\delta 1_H)} = 2\mu_2 V^2 \lambda' + \frac{\lambda' \mu_3}{16\pi^2} m_{\text{SUSY}}^2$$

$$\Rightarrow \mu_2 = -\frac{1}{32\pi^2} \mu_3 \left(\frac{m_{\text{SUSY}}^2}{V^2} \right) \sim M_G$$

THE SLIDING SINGLET MECHANISM CAN BE MADE TO WORK IN SU(6) AND OTHER SU(N) GROUPS.

(Sen 1984, Barr 1998, Maekawa 2003)
+ Yamashita hep-ph/0305116

CONSIDER $SU(6)$

Adjoint VEV:

$$\langle 35_H \rangle = \Omega \begin{pmatrix} 1 & & & & & \\ & 1 & & & & \\ & & -1 & & & \\ & & & -1 & & \\ & & & & -1 & \\ & & & & & -1 \end{pmatrix}$$

Can arise
from minima
of potential

Fundamental VEV:

$$\langle 6_H \rangle = \begin{pmatrix} \Sigma \\ 0 \\ 0 \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$\Sigma \sim M_6$. THIS BREAKS
 $SU(6) \rightarrow SU(5)$.

$\langle 35_H \rangle$ FURTHER BREAKS
IT TO G_{SM} .

$$W \supset \lambda \bar{6}_H 35_H 6_H + \lambda' \bar{6}_H 1_H 6_H$$

$$\Rightarrow (\bar{\Sigma}, \bar{H}, \bar{H}_c) \begin{bmatrix} (\lambda\Omega + \lambda' 1_H) \\ (\lambda\Omega + \lambda' 1_H) I_2 \\ (-\lambda\Omega + \lambda' 1_H) I_3 \end{bmatrix} \begin{pmatrix} \Sigma \\ H \\ H_c \end{pmatrix}$$

$$\Rightarrow V(1_H) = (\mu_2(1_H))^2 (v^2 + \Sigma^2 + \bar{\Sigma}^2) + \text{SUSY terms}$$

$$\mu_2 \sim \frac{1}{32\pi^2} \mu_3 \frac{m_{\text{SUSY}}^2}{v^2 + \Sigma^2 + \bar{\Sigma}^2}$$

There are some technical difficulties, but they can be overcome,

(2) MISSING VEV MECHANISM

(DIMOPOULOS - WILCZEK MECHANISM)

THE PROBLEM IN $SU(5)$ IS THAT THE GENERATORS ARE TRACELESS:

$$\langle 24_H \rangle = \Omega(\frac{1}{2}) = \Omega \begin{pmatrix} \frac{1}{2} & & & & \\ & \frac{1}{2} & & & \\ & & -\frac{1}{3} & & \\ & & & -\frac{1}{3} & \\ & & & & -\frac{1}{3} \end{pmatrix}$$

SO NEED ANOTHER TERM TO CANCEL THE $\frac{1}{2}\Omega$ CONTRIBUTION TO μ_2 .

IN $U(5)$ THE GENERATORS ARE NOT TRACELESS, BUT $U(5)$ IS REALLY $SU(5) \times U(1)$ AND THE ADJOINT "25" IS REALLY TWO IRREDUCIBLE REPS. WHOSE COUPLINGS ARE NOT RELATED BY SYMMETRY: $25 = 24 + 1$. SO CANCELLATION BETWEEN $\langle 24_H \rangle$ AND $\langle 1_H \rangle$ WOULD STILL BE ACCIDENTAL.

BUT $SO(10) \supset U(5)$

generator (in vector rep) of $SO(10)$

$$\begin{pmatrix} a_1 & & & & \\ -a_1 & a_2 & & & \\ & -a_2 & a_3 & & \\ & & -a_3 & a_4 & \\ & & & -a_4 & a_5 \\ & & & & -a_5 \end{pmatrix}$$



generator (in fund. rep.) of $U(5)$

$$\begin{pmatrix} a_1 & & & & \\ & a_2 & & & \\ & & a_3 & & \\ & & & a_4 & \\ & & & & a_5 \end{pmatrix}$$

does not need to be traceless

B-L corresponds to $a_1 = a_2 = 0$
 $a_3 = a_4 = a_5 = -\frac{2}{3}$

$\langle 45_H \rangle \propto B-L$ CAN ARISE FROM MINIMIZING A SIMPLE POTENTIAL.

eg.

$$W_{45} = \frac{\lambda}{M} \text{tr} (45_H)^4 - \lambda' M \text{tr} (45_H)^2$$

$$= \frac{2\lambda}{M} \sum_{i=1}^5 (a_i)^4 - 2\lambda' M \sum_{i=1}^5 (a_i)^2$$

$$\Rightarrow 0 = \frac{\partial W}{\partial a_i} = \frac{8\lambda}{M} a_i^3 - 4\lambda' M a_i$$

$$\Rightarrow a_i = 0 \text{ or } \sqrt{\frac{\lambda'}{2\lambda}} M \equiv \Omega$$

Multiply degenerate minimum, one of them is

$$\langle 45_H \rangle = \Omega \begin{pmatrix} \Omega & & & & \\ & \Omega & & & \\ & & -\Omega & & \\ & & & -\Omega & \\ & & & & -\Omega \end{pmatrix} = \frac{3}{2} \Omega (B-L).$$


This solves 2/3 splitting problem:

$$W \supset 10_{1H} 45_H 10_{2H}$$

$$\rightarrow (\bar{5}_{1H} \binom{24+1}{H} 5_{2H} + 5_{1H} \binom{24+1}{H} \bar{5}_{2H})$$

only give mass to triplets (which have $B-L = \pm 2/3$)

[Need $10_{1H}, 10_{2H}$ different as otherwise term vanishes:
 $T^a T^b T^c$]



NOTE:
 4 LIGHT
 DOUBLET HIGGS,
 NOT 2.

(3) MISSING PARTNER MECHANISM

CONSIDER, IN $SU(5)$:

$$\begin{array}{cccc} \bar{5}_H & R_H & \bar{R}_H & 5_H \\ \left(\begin{array}{c} \bar{2} \\ \bar{3} \end{array} \right) & \left(\begin{array}{c} \text{other} \\ 3 \end{array} \right) & \left(\begin{array}{c} \overline{\text{other}} \\ \bar{3} \end{array} \right) & \left(\begin{array}{c} 2 \\ 3 \end{array} \right) \end{array}$$

SUPPOSE "other" contains no $2 \equiv (1, 2, \frac{1}{2})$.
 .e. doublet "partner" of triplet in R is "missing"
 $\Rightarrow \bar{2}, 2$ can remain light.

THIS CAN BE IMPLEMENTED IN $SU(5)$

$$\begin{array}{ccc} \bar{5}_H & R_H & \langle V_H \rangle \\ & \uparrow & \uparrow \\ & & \text{Needs to leave SM} \\ & & \text{unbroken. i.e. contain} \\ & & (1, 1, 0) \\ & \uparrow & \\ & & \text{Must have } (3, 1, -\frac{1}{3}) \text{ but not } (1, 2, \frac{1}{2}) \end{array}$$

SIMPLEST POSSIBILITY is

$$R = 50 = T^{[\alpha\beta\gamma]}_{[8c]} \quad (\text{traceless})$$

$$V = 75 = T^{[\alpha\beta]}_{[8a]} \quad (\text{traceless})$$

We will now show that 50 contains
 $(3, 1, -\frac{1}{3})$ but not $(1, 2, \frac{1}{2})$.

$$10 \times \bar{5} = 45 + 5$$

$$T^{[\alpha\beta]} T_\gamma = \left[T^{[\alpha\beta]} T_\gamma - \frac{1}{4} T^{[\alpha\sigma} T_\sigma \delta^{\beta\gamma]} \right] \text{ (Traceless)}$$

$$+ \frac{1}{4} \underbrace{\left[T^{[\alpha\sigma} T_\sigma \delta^{\beta\gamma]} \right]}_5$$

$$\bar{10} \times \bar{10} = 50_S + 45_A + 5_S$$

$$T^{[\alpha\beta\gamma]} T_{[\delta\epsilon]} = \left[T^{[\alpha\beta\gamma]}_{[\delta\epsilon]} - (\text{traces}) \right] \text{ (Traceless)}$$

$$+ \left[\underbrace{T^{[\alpha\beta\sigma} T_{[\delta\sigma} \delta^{\gamma\epsilon]}]}_{45} - (\text{traces}) \right]$$

$$+ \left[\underbrace{T^{[\alpha\sigma\tau} T_{\sigma\tau} \delta^{\beta\epsilon]}_{[\delta\delta^{\gamma\epsilon}]}}_5 \right]$$

$$\bar{50} \text{ } (\bar{10} \times \bar{10})_S = 50 + 5$$

$$\begin{bmatrix} (1, 1, -1) \\ (\bar{3}, 2, -\frac{1}{6}) \\ (3, 1, \frac{2}{3}) \end{bmatrix} \times \begin{bmatrix} (1, 1, -1), (\bar{3}, 2, -\frac{1}{6}), (3, 1, \frac{2}{3}) \end{bmatrix}$$

$(1, 1, -2)_S$	$(\bar{3}, 2, -\frac{1}{6})$	$(3, 1, -\frac{2}{3})$
$(\bar{3}, 2, -\frac{1}{6})$	$(3, 1, -\frac{1}{3})_S$ $(6, 3, -\frac{1}{3})_S$ $(3, 3, -\frac{1}{3})_A$ $(6, 1, -\frac{1}{3})_A$	$(8, 2, \frac{1}{2})$ $(1, 2, \frac{1}{2})$
$(3, 1, -\frac{2}{3})$	$(8, 2, \frac{1}{2})$ $(1, 2, \frac{1}{2})$	$(\bar{6}, 1, \frac{1}{3})_S$ $(3, 1, \frac{1}{3})_A$

$$50 = (1, 1, -2) + \cancel{(3, 1, -\frac{1}{3})} + (6, 3, -\frac{1}{3}) + (\bar{6}, 1, \frac{1}{3})$$

$$+ (\bar{3}, 2, -\frac{1}{6}) + (3, 1, -\frac{1}{3}) + (8, 2, \frac{1}{2}) + \cancel{(1, 2, \frac{1}{2})}$$

crossed out reps are = 5

$$\text{So } 50 \supset (3, 1, -\frac{1}{3}) \text{ but } \not\supset (1, 2, \frac{1}{2})$$


IN FLIPPED $SU(5)$ THE MISSING PARTNER MECHANISM WORKS ELEGANTLY (IF NOT UNIFIED IN $SO(10)$!) 2.11

SUPPOSE

$$\begin{aligned}
 & (\bar{5}_H^{-2} \quad 10_H^1 \quad 10_H^0) + (\bar{5}_H^{-2} \quad \bar{10}_H^{-1} \quad \bar{10}_H^{-1}) \\
 & \begin{pmatrix} (1, 2, -\frac{1}{2}) \\ (3, 1, -\frac{2}{3}) \end{pmatrix} \begin{pmatrix} (1, 1, 0) \\ (3, 2, \frac{1}{6}) \\ (\bar{3}, 1, \frac{1}{3}) \end{pmatrix} \begin{pmatrix} \langle (1, 1, 0) \rangle \\ (3, 2, \frac{1}{6}) \\ (\bar{3}, 1, \frac{1}{3}) \end{pmatrix} + \begin{pmatrix} (1, 2, \frac{1}{2}) \\ (\bar{3}, 1, \frac{1}{3}) \end{pmatrix} \begin{pmatrix} (1, 1, 0) \\ (3, 2, -\frac{1}{6}) \\ (3, 1, -\frac{2}{3}) \end{pmatrix} \begin{pmatrix} \langle (1, 1, 0) \rangle \\ (\bar{3}, 2, -\frac{1}{6}) \\ (3, 1, -\frac{2}{3}) \end{pmatrix}
 \end{aligned}$$

THE "OTHER" FIELDS GET EATEN BY HIGGS MECHANISM!

- ⇒ No NEED TO COUPLE $M 10_H^1 \bar{10}_H^{-1}$
- ⇒ $d=5$ PROTON DECAY OPERATORS CAN BE SUPPRESSED.



HOWEVER, IF $SU(5) \times U(1)_X$ UNIFIED IN $SO(10)$, THEN "MISSING PARTNER" IS NOT MISSING.

$$10_H^1 \subset 16_H \quad \text{WHICH CONTAINS } \bar{5}^{-3} = \begin{pmatrix} (1, 2, -\frac{1}{2}) \\ (\bar{3}, 1, -\frac{2}{3}) \end{pmatrix}$$

$$\bar{5}_H^{-2} \bar{5}_H^{-3} \langle 10_H^1 \rangle$$

(D) SO(10): GAUGE HIERARCHY AND PROTON DECAY

As we saw, the 2/3 splitting problem in SO(10) can be solved by assuming $\langle 45_H \rangle \propto B-L$ and having a term

$$\lambda 10_{1H} 45_H 10_{2H} \Rightarrow \left[\bar{5}_{1H} \overbrace{(24_H + 1_H)}^{= (B-L)\Omega} 5_{2H} + 5_{1H} (24_H + 1_H) \bar{5}_{2H} \right]$$

$$\Rightarrow \begin{pmatrix} 2_1 \\ 3_1 \end{pmatrix} \begin{pmatrix} \bar{2}_2 \\ \bar{3}_2 \end{pmatrix} \begin{pmatrix} 2_2 \\ 3_2 \end{pmatrix} \begin{pmatrix} \bar{2}_1 \\ \bar{3}_1 \end{pmatrix}$$

To avoid having 4 light doublets, one needs to give mass to $\bar{2}_2, 2_2$. (Assume MSSM doublets of Higgs H_u, H_d are $2_1, \bar{2}_1$).

Simplest: $M 10_{2H} 10_{2H}$

Look at masses of color-triplet Higgsinos:

$$(\bar{3}_1, \bar{3}_2) \begin{pmatrix} 0 & \frac{2}{3}\lambda\Omega \\ \frac{2}{3}\lambda\Omega & M \end{pmatrix} \begin{pmatrix} 3_1 \\ 3_2 \end{pmatrix}$$

$$\equiv \mathcal{M}_3$$

PROTON DECAY AMPLITUDE $\propto (\mathcal{M}_3^{-1})_{11}$

$$\mathcal{M}_3^{-1} = \begin{pmatrix} \frac{-M}{(\frac{2}{3}\lambda\Omega)^2} & \frac{1}{(\frac{2}{3}\lambda\Omega)} \\ \frac{1}{(\frac{2}{3}\lambda\Omega)} & 0 \end{pmatrix}$$

Can we make p decay amplitude small by setting $M \ll M_G$? YES, BUT

MAKING M SMALL SUPPRESSES HIGGSINO-MEDIATED PROTON DECAY, BUT MAKES UNIFICATION OF GAUGE COUPLINGS WORSE:

$$\delta E_3 \Big|_{\frac{2}{3}} = \frac{\delta \alpha_3(M_G)}{\alpha_3(M_G)} = \frac{3\alpha_G}{5\pi} \cdot [\ln|\det M_3| - \ln|\det M_2|]$$

NOTE: ONLY "SPLIT MULTIPLETS" OF $SU(5)$ CONTRIBUTE AT ONE LOOP TO RGE SPLITTING OF THE THREE SM GAUGE COUPLINGS. THUS, IF $\det M_3 = \det M_2 \Rightarrow \delta E_3 = 0$

TRIPLET MASSES: $M_3 = \begin{bmatrix} 0(\mu) & \frac{2}{3}\lambda\Omega \\ \frac{2}{3}\lambda\Omega & M \end{bmatrix}, M_2 = \begin{bmatrix} \mu & 0 \\ 0 & M \end{bmatrix}$

$$\Rightarrow \delta E_3 \Big|_{\frac{2}{3}} = - \frac{3\alpha_G}{5\pi} \ln \left(\mu \frac{M}{(\frac{2}{3}\lambda\Omega)^2} \right)$$

≈ 0.008 $(M_3^{-1})_{11}$ ← in p decay AMPLITUDE!

In Minimal SUSY $SU(5)$, NEED $(M_3^{-1})_{11}$ at least $1/20 \times$ (and typically $1/100 \times$) the value that gives good unification to sufficiently suppress p decay. $\Rightarrow \delta E_3 \approx +0.04$
 VERY GENERAL RESULT (SM. B. hep-ph/9806217)

- F (1) ONLY MASSES OF THE FORM $(\bar{3}_H 3_H)$ OR $(\bar{2}_H 2_H + \bar{3}_H 3_H) = (\bar{3}_H 5_H)$, BUT NOT $(\bar{2}_H 2_H)$, AND
- (2) NO ARTIFICIAL CANCELLATIONS

\Rightarrow SUPPRESSION OF P DECAY AMPLITUDE INCREASES E_3 .

HOWEVER, SUPPOSE: $10_{1H} 45_{H} 10_{2H} + 10_{2H} (45'_H)^2 10_{2H} / M_G$
 \uparrow B-L \uparrow I_{3R}

$$H_u \begin{pmatrix} 2_1 \\ 3_1 \end{pmatrix} \begin{pmatrix} \bar{2}_2 \\ \bar{3}_2 \end{pmatrix} \begin{pmatrix} 2_2 \\ 3_2 \end{pmatrix} \begin{pmatrix} \bar{2}_1 \\ \bar{3}_1 \end{pmatrix} H_u$$

THIS REQUIRES 2 ADJUNCT HIGGS

(E) STABILITY OF GAUGE HIERARCHY 3.14
IN SO(10) MODELS [K.S. Babu + SMB 1993]

$$M_2 = \begin{bmatrix} \mu & \lambda \langle 45_H \rangle \\ \lambda \langle 45_H \rangle & M \end{bmatrix}$$

\uparrow Need this $\approx \sqrt{M_W M_G}$
 \uparrow Need this $\approx M_W$

- Must forbid $M_G(10_{1H} 10_{1H})$ by some symmetry, Also $M_G(10_{1H} 10_{2H})$.
- Must prevent terms that destabilize $\langle 45_H \rangle \propto B-L$

$$\langle 45_H \rangle = \begin{bmatrix} b \\ -b \\ a \\ -a \\ a \\ -a \\ a \\ -a \end{bmatrix} = \frac{3}{2} a (B-L) + 2b (I_{3R})$$

Must have $b/a \approx \sqrt{M_W/M_G} \sim 10^{-7}$

THERE IS A TECHNICAL DIFFICULTY ASSOCIATED WITH COUPLING ADJOINT HIGGS SECTOR AND SPINOR HIGGS SECTOR

$$W_{\text{Higgs}} = W(45_H) + W(\overline{16}_H, 16_H) + \underbrace{W_{45-16}}_{?} + 10_{1H} 45_H 10_{2H}$$

IF THERE ARE NO COUPLINGS (DIRECT OR INDIRECT) OF 45_H TO $\overline{16}_H + 16_H$, THEN BAD GOLDSTONE MODES RESULT.

WITHOUT 45_H COUPLING TO $\bar{16}_H + 16_H$, THE RELATIVE DIRECTION OF $\langle 45_H \rangle$ AND $\langle 16_H \rangle$ IS NOT DETERMINED BY $W \Rightarrow$ MASSLESS MODE.

WHAT ARE THE GOLDSTONE MODES?

LOOK AT WHICH GENERATORS ARE BROKEN BY BOTH $\langle 45_H \rangle$ AND $\langle 16_H \rangle$. THERE WOULD BE TWO "WOULD-BE GOLDSTONE BOSONS" FOR EACH SUCH GENERATOR, OF WHICH ONLY ONE CAN BE "EATEN".

$\langle 1^5(16_H) \rangle$ BREAKS $SO(10) \rightarrow SU(5)$

$$\begin{aligned} \Rightarrow 45 &= 24 + \overbrace{1 + 10 + 10}^{\text{broken}} \\ &= (1, 1, 0) + \sqrt{2}(1, 1, 0) + \sqrt{2}(1, 1, +1) + \sqrt{2}(1, 1, -1) \\ &\quad + (3, 2, -5/6) \quad + \sqrt{2}(3, 2, 1/6) + \sqrt{2}(\bar{3}, 2, -1/6) \\ &\quad + (3, 2, 5/6) \quad + \sqrt{2}(3, 1, -2/3) + \sqrt{2}(3, 1, 2/3) \\ &\quad + (8, 1, 0) \end{aligned}$$

$\langle 45_H \rangle \propto B-L$ BREAKS $SO(10) \rightarrow SU(3)_c \times U(1)_{B-L} \times SU(2)_L \times SU(2)_R$

$$\begin{aligned} 45 &= (15, 1, 1) + (1, 3, 1) + (1, 1, 3) + (6, 2, 2) \text{ of P-S} \\ &= (1^0, 1, 1) + (1^0, 3, 1) + (1^0, 1, 3) + \sqrt{2}(3^{-2/3}, 2, 2) \\ &\quad (8^0, 1, 1) \quad \sqrt{2}(3^{2/3}, 2, 2) \\ &\quad \sqrt{2}(3^{1/3}, 1, 1) \\ &\quad \sqrt{2}(\bar{3}^{-4/3}, 1, 1) \end{aligned}$$

\Rightarrow GENERATORS IN $(\bar{3}, 2, -1/6) + (3, 2, 1/6)$
 $(\bar{3}, 1, -2/3) + (3, 1, 2/3)$

ARE TWICE BROKEN

THE SIMPLEST WAY TO COUPLE 45_H TO $\overline{16}_H + 16_H$ IS NO GOOD:

$$\overline{16}_H \ 45_H \ 16_H$$

$$\langle 45_H \rangle = \frac{2}{3} a (B-L) + 2 b (I_{3R})$$

$$\langle 1(16_H) \rangle \sim M_G, \quad \langle 1(\overline{16}_H) \rangle \sim M_G.$$

↑ has quantum numbers of N_L^c : $\begin{cases} B-L = +1 \\ I_{3R} = -\frac{1}{2} \end{cases}$

$$\Rightarrow W \supset \underbrace{\# M_G}_{\text{From } (45_H)^2} b^2 + \# [1(\overline{16}_H) b \ 1(16_H)]$$

$$0 = -F_b^* = \frac{\partial W}{\partial b} = \# M_G b + \# \langle 1(\overline{16}_H) \rangle \langle 1(16_H) \rangle$$

$$\Rightarrow \boxed{b \sim M_G} \quad (\text{destroys hierarchy})$$

There are various ways to couple 45_H to $\overline{16}_H, 16_H$ without destroying gauge hierarchy:

$$\left\{ \begin{array}{l} \overline{16}_H \ 45_x \ 16_H + \underbrace{45_x \ 45_{B-L} \ 45_{I_{3R}}}_{\text{THREE DISTINCT ADJUNCT HIGGS}} + \dots \quad (\text{Babu Barr}) \\ \overline{16}_H (45_H + 1_H) \underset{\uparrow \text{VEV}=0}{16'_H} + \overline{16}'_H (45_H + 1'_H) \underset{\uparrow \text{VEV}=0}{16_H} \quad (\text{Barr Raby}) \end{array} \right.$$

LECTURE IV : FERMION MASSES IN SUSY GUTS

WE HAVE ALREADY SEEN THAT IN "MINIMAL SU(5)"

$$(Y_U)_{AB} (10_A 10_B) 5_H \rightarrow M_U = M_U^T = Y_U V_U$$

$$(Y_D)_{AB} (10_A \bar{5}_B) \bar{5}_H \rightarrow M_D = M_D^T = Y_D V_D$$

$$(Y_N)_{AB} (\bar{5}_A 1_B) 5_H \rightarrow M_N = Y_N V_U$$

$$M_D = M_D^T \Rightarrow \begin{cases} m_b^0 = m_\tau^0 & \text{within } 20\% \\ m_s^0 = m_\mu^0 & (\text{expt } \approx \frac{1}{3} m_\mu^0) \\ m_d^0 = m_e^0 & (\text{expt } \approx 3 m_e^0) \end{cases}$$

1) GEORGI - JARLSKOG

$$SU(5): (10_{1L} 10_{2L} 10_{3L}) \begin{bmatrix} 0 & \langle \bar{5}_H \rangle & 0 \\ \langle \bar{5}_H \rangle & \langle \overline{45}_H \rangle & 0 \\ 0 & 0 & \langle \bar{5}_H \rangle \end{bmatrix} \begin{pmatrix} \bar{5}_{1L} \\ \bar{5}_{2L} \\ \bar{5}_{3L} \end{pmatrix} \quad \begin{array}{l} \text{Yukawa} \\ \text{constants} \\ \text{not shown} \end{array}$$

$$(10_L \bar{5}_L) \bar{5}_H: \begin{cases} \psi^{a2} \psi_a \langle H_2 \rangle \rightarrow d_L d_L^c \langle H_d \rangle \\ \psi^{i2} \psi_i \langle H_2 \rangle \rightarrow \ell_L^+ \ell_L^- \langle H_d \rangle \end{cases}$$

$$(10_L \bar{5}_L) \overline{45}_H: \begin{cases} \psi^{a2} \psi_a \langle H_{a2}^a \rangle \rightarrow (-\frac{1}{3}) d_L d_L^c \langle H_d' \rangle \\ \psi^{i2} \psi_i \langle H_{i2}^{i'} \rangle \rightarrow \ell_L^+ \ell_L^- \langle H_d' \rangle \end{cases}$$

NOTE: $\overline{45}$ is traceless $\Rightarrow \sum_{a=3,4,5} H_{a2}^a + H_{i2}^{i'} + \cancel{H_{11}} = 0$

$$\Rightarrow \langle H_{a2}^a \rangle = -\frac{1}{3} \langle H_{i2}^{i'} \rangle$$

NO SUM

$$M_L^0 = \begin{bmatrix} 0 & C & \\ C & B & \\ & & A \end{bmatrix}$$

$$M_D^0 = \begin{bmatrix} C & \frac{1}{3}B & \\ & \frac{1}{3}B & \\ & & A \end{bmatrix}$$

LET $A \gg B \gg C$ ↓ diagonalize

$$\begin{bmatrix} -\frac{C^2}{B} & & \\ & B & \\ & & A \end{bmatrix}$$

$$\begin{bmatrix} +\frac{3C^2}{B} & & \\ & -\frac{1}{3}B & \\ & & A \end{bmatrix}$$

NOTE: $\det M_L^0 = \det M_D^0 = -AC^2$

$$\Rightarrow m_s^0 \cong \frac{1}{3} m_\mu^0 \quad m_d^0 \cong 3 m_e^0$$

These factors of $\frac{1}{3}, 3$ are called "Georgi Jarlskog factors."

SUCH FACTORS OF $\frac{1}{3}, 3$ ARISE VERY NATURALLY IN GUTS. THEY COME FROM FACT THAT THERE ARE 3 COLORS

ANOTHER WAY TO GET SUCH FACTORS

$$B-L = \begin{pmatrix} \frac{1}{3} & & & \\ & \frac{1}{3} & & \\ & & \frac{1}{3} & \\ & & & -1 \end{pmatrix} \quad (\text{generator of } SU(4)_c)$$

= $\frac{1}{3}$ for quarks, -1 for leptons.

NOTE FORM OF 12 BLOCK ABOVE: $\begin{pmatrix} 0 & C \\ C & B \end{pmatrix}$.

Suggested independently by Weinberg, Wilczek + Zee, Fritzsche in 1979 to explain Cabibbo angle.

"TEXTURE"

TEXTURES AND MIXING ANGLES

CONSIDER A TWO-FAMILY MODEL:

$$(\bar{d}_L \bar{s}_L)^{(0)} \begin{pmatrix} 0 & c' \\ c' & B' \end{pmatrix} \begin{pmatrix} d_R^{(0)} \\ s_R^{(0)} \end{pmatrix} + (\bar{u}_L \bar{c}_L)^{(0)} \begin{pmatrix} 0 & c \\ c & B \end{pmatrix} \begin{pmatrix} u_R \\ c_R \end{pmatrix}$$

$$\Rightarrow \begin{pmatrix} d_L \\ s_L \end{pmatrix} = \begin{pmatrix} \cos \theta_{dL} & \sin \theta_{dL} \\ -\sin \theta_{dL} & \cos \theta_{dL} \end{pmatrix} \begin{pmatrix} d_L^{(0)} \\ s_L^{(0)} \end{pmatrix} \quad \text{similar for } d_R, s_R \quad (\theta_{dR})$$

$$\begin{pmatrix} u_L \\ c_L \end{pmatrix} = \begin{pmatrix} \cos \theta_{uL} & \sin \theta_{uL} \\ -\sin \theta_{uL} & \cos \theta_{uL} \end{pmatrix} \begin{pmatrix} u_L^{(0)} \\ c_L^{(0)} \end{pmatrix} \quad \text{similar for } u_R, c_R \quad (\theta_{uR})$$

$$\Rightarrow \tan 2\theta_{dL} = \frac{2c'}{B'} \Rightarrow \theta_{dL} \approx \frac{c'}{B'} \approx \sqrt{\frac{m_d}{m_s}}$$

$$\tan 2\theta_{uL} = \frac{2c}{B} \Rightarrow \theta_{uL} \approx \frac{c}{B} \approx \sqrt{\frac{m_u}{m_c}}$$

bibbo:

$$\theta_C = \theta_{dL} - \theta_{uL} \approx \sqrt{\frac{m_d}{m_s}} - e^{i\phi} \sqrt{\frac{m_u}{m_c}}$$

$\approx 0.2 \qquad \qquad \qquad \approx 0.2 \qquad \qquad \qquad \approx 0.07$

Old numerology: $\sin \theta_C \approx m_\mu / m_K \approx \sqrt{m_d / m_s}$

Note that here matrices are assumed symmetric in SU(5), justified for M_u , not M_D, M_L .

However, this assumption reduces # of parameters and thus allows a prediction.

Also gives $\theta_L = \theta_R$

MILAR TEXTURES FOR 23 BLOCK GIVE RELATIONS LIKE

$$\left\{ \begin{array}{l} V_{cb} = \sin \theta_{23}^{\text{charm}} \approx \sqrt{m_s / m_b} - e^{i\phi'} \sqrt{m_c / m_e} \\ \sim 0.04 \qquad \qquad \sim 0.14 \qquad \qquad \sim 0.04 \\ V_{\mu 3} = \sin \theta_{23}^{\text{lepton}} \approx \sqrt{m_\mu / m_\tau} - e^{i\phi''} \sqrt{m_2 / m_3} \\ \sim 0.7 \qquad \qquad \sim 0.25 \qquad \qquad \sim 0.15 \end{array} \right.$$

3) AN SO(10) MODEL OF QUARK

4.4

AND LEPTON MASSES

(Albright, Babu, Barr 1998)
(Albright, Barr 1999)

(Similar model by
Babu, Pati, Wilczek)

THE VEVs:

$$O(M_G): \begin{cases} \langle 45_H \rangle \propto B-L \\ \langle 16_H \rangle \propto 1(16) \\ \langle \bar{16}_H \rangle \propto 1(\bar{16}) \end{cases}$$

$$O(M_W): \begin{cases} \langle 10_H \rangle \propto \begin{bmatrix} 2(5(10)) \\ \bar{2}(\bar{5}(10)) \end{bmatrix} \\ \langle 16'_H \rangle \propto \bar{2}(\bar{5}(16)) \end{cases}$$

THIRD FAMILY MASSES

$$16_3 \ 16_3 \ \langle 10_H \rangle \rightarrow M_U, M_D, M_L, M_N \propto \begin{pmatrix} 0 & \\ & 0 \\ & & 1 \end{pmatrix} \begin{matrix} m_1 \\ m_2 \\ m_3 \end{matrix}$$


SECOND FAMILY

Need Georgi-Jarlskog factors

This can be achieved if the $\langle 45_H \rangle$ is involved. Simplest effective operator is

$$16_A \ 16_B \ \langle 10_H \rangle \langle 45_H \rangle / M_G$$

This can arise from "integrating out" $\bar{16}' + 16'$



There are two ways to contract the $SO(10)$ indices in the product $16 \ 16 \ 10 \ 45$: 4.5

$$16 \times 16 = 10_S + 120_A + 126_S$$

$$10 \times 45 = 10 + 120 + 320$$

So 16×16 must be contracted in 10 or 120 .

If $\langle 45_H \rangle \propto B-L$ then only 120 channel contributes to quark and lepton masses

Thus this term is antisymmetric in flavor

PROOF $\langle 10_H \rangle \sim (1, 2, 2)$ of Pati-Salam Group.

$\langle 45_H \rangle \sim (15, 1, 1)$ ($\propto B-L =$ a generator of $SU(4)_c$)

$$\Rightarrow \langle 10_H \rangle \langle 45_H \rangle \sim (15, 2, 2)$$

But, this is not in 10 of $SO(10)$.

\Rightarrow it must be in 120 .

CONSIDER, THEN,

$$(16_2 \ 16_3) \langle 10_H \rangle \langle 45_H \rangle / M_G$$

$$\Rightarrow M_U = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & \epsilon/3 \\ 0 & -\epsilon/3 & 1 \end{bmatrix} m_u, \quad M_D = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & \epsilon/3 \\ 0 & -\epsilon/3 & 1 \end{bmatrix} m_D$$

$$M_N = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -\epsilon \\ 0 & \epsilon & 1 \end{bmatrix} m_u, \quad M_L = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -\epsilon \\ 0 & \epsilon & 1 \end{bmatrix} m_D$$

Not realistic. FOUR PROBLEMS:

(1) Georgi-Jarlskog factors are $1/9$ not $1/3$

(2) $M_D \propto M_U \Rightarrow$ No CKM MIXING.

eg. $V_{cb} = 0$

(3) $M_D \propto M_U \Rightarrow m_c^0/m_t^0 = m_s^0/m_b^0$
 $\sim 0.0025 \quad \sim 0.02$


(4) The atmospheric ν angle $\theta_{\mu 3} = \mathcal{O}(\epsilon) \ll 1$

Need another contribution to 23 block that makes M_U not proportional to M_D .


$\langle 45_H \rangle$ does not distinguish U from D.

Simplest possibility: $\langle 16_H \rangle$ contributes.

Consider



THREE WAYS OF CONTRACTING INDICES. ASSUME OPERATOR COMES FROM DIAGRAM:



$$M_U = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & \epsilon/3 \\ 0 & -\epsilon/3 & 1 \end{bmatrix} M_U \quad M_D = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & \epsilon/3 \\ 0 & \sigma^{-\epsilon/3} & 1 \end{bmatrix} M_D \quad \text{v.7}$$

$$M_N = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & -\epsilon \\ 0 & \epsilon & 1 \end{bmatrix} M_U \quad M_L = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & \sigma^{-\epsilon} \\ 0 & \epsilon & 1 \end{bmatrix} M_D$$

ASSUME $\epsilon \ll \sigma \sim 1$

Then: (1) Georgi - Jarlskog factor is $\approx 1/3$

(2) $M_U \text{ not } \propto M_D \Rightarrow V_{cb} \neq 0$

in fact $V_{cb} \approx \frac{\epsilon}{3} \frac{\sigma^2}{1+\sigma^2}$

(3) $\frac{m_c^0}{m_t^0} \approx (\epsilon/3)^2 \ll \frac{m_s^0}{m_b^0} \approx \frac{\epsilon}{3} \frac{\sigma}{1+\sigma^2}$

(4) $V_{cb} \sim \mathcal{O}(\sigma) \sim \mathcal{O}\left(\frac{V_{cb}}{m_s/m_b}\right) \sim 1$

FIRST FAMILY CAN BE INCLUDED SIMPLY

$$M_U = \begin{bmatrix} \eta & 0 & 0 \\ 0 & 0 & \epsilon/3 \\ 0 & -\epsilon/3 & 1 \end{bmatrix} M_U \quad M_D = \begin{bmatrix} 0 & \delta & \delta' \\ \delta & 0 & \epsilon/3 \\ \delta' & \sigma^{-\epsilon/3} & 1 \end{bmatrix} M_D$$

$$M_N = \begin{bmatrix} \eta & 0 & 0 \\ 0 & 0 & -\epsilon \\ 0 & \epsilon & 1 \end{bmatrix} M_U \quad M_L = \begin{bmatrix} 0 & \delta & \delta' \\ \delta & 0 & \sigma^{-\epsilon} \\ \delta' & \epsilon & 1 \end{bmatrix} M_D$$