

SMR 1595 - 7

**Joint DEMOCRITOS - ICTP School on
CONTINUUM QUANTUM MONTE CARLO METHODS
12 - 23 January 2004**

SUMMARY AND PROBLEMS WITH VARIATIONAL METHODS

David M. CEPERLEY

Beckman Institute for Advanced Studies & Technology
University of Illinois at Urbana-Champaign - N.C.S.A.
II-61801 Urbana, U.S.A.

These are preliminary lecture notes, intended only for distribution to participants.


Summary and problems with variational methods

- Powerful method since you can use any trial function
- Scaling (computational effort vs. size) is almost classical
- Learn directly about what works in wavefunctions
- No sign problem
- Optimization is time consuming
- Energy is insensitive to order parameter
- Non-energetic properties are less accurate. $O(1)$ vs. $O(2)$ for energy.
- Difficult to find out how accurate results are.
- Favors simple states over more complicated states, e.g.
 - Solid over liquid
 - Polarized over unpolarized

What goes into the trial wave function comes out! “GIGO”

We need a more automatic method! Projector Monte Carlo

Summary of Variational (VMC)




Dependence of energy on wavefunction

3d Electron fluid at a density $r_s=10$

Kwon, Ceperley, Martin, Phys. Rev. **B58**,6800, 1998

- Wavefunctions
 - Slater-Jastrow (SJ)
 - three-body (3)
 - backflow (BF)
 - fixed-node (FN)
- Energy $\langle \phi | H | \phi \rangle$ converges to ground state
- Variance $\langle \phi | [H-E]^2 | \phi \rangle$ to zero.
- Using 3B-BF gains a factor of 4.
- Using DMC gains a factor of 4.



Projector Monte Carlo

(variants: Green's function MC, Diffusion MC, Reptation MC)

- Project single state using the Hamiltonian

$$\mathbf{f}(t) = e^{-(H-E_T)t} \mathbf{f}(0)$$
- We show that this is a diffusion + branching operator. Maybe we can interpret as a probability. **But is this a probability?**
- **Yes!** for bosons since ground state can be made real and non-negative.
- **But** all excited states must have sign changes. This is the "sign problem."
- For efficiency we do "importance sampling."
- Avoid sign problem with the fixed-node method.

Diffusion Monte Carlo

- How do we analyze this operator?

$$\mathbf{y}(R, t) = e^{-(H-E_T)t} \mathbf{y}(R, 0)$$

- Expand into exact eigenstates of H.

$$H\mathbf{f}_a = E_a\mathbf{f}_a$$

$$\mathbf{y}(R, 0) = \sum_a \mathbf{f}_a(R) \langle \mathbf{f}_a | \mathbf{y}(0) \rangle$$

- Then the evolution is simple in this basis.

$$\mathbf{y}(R, t) = \sum_a \mathbf{f}_a(R) e^{-t(E_a-E_T)} \langle \mathbf{f}_a | \mathbf{y}(0) \rangle$$

- Long time limit is lowest energy state that overlaps with the initial state, usually the ground state.

$$\lim_{t \rightarrow \infty} \mathbf{y}(R, t) = \mathbf{f}_0(R) e^{-t(E_0-E_T)} \langle \mathbf{f}_0 | \mathbf{y}(0) \rangle$$

$$E_0 \approx E_T \Rightarrow \text{normalization fixed}$$

- How to carry out on the computer?

The Green's function

- Operator notation

$$\frac{d\hat{\mathbf{r}}}{dt} = -\hat{H}\hat{\mathbf{r}}$$

$$\hat{\mathbf{r}} = e^{-\hat{H}t}$$

- We define the coordinate green's function (or density matrix) by:

$$G(R \rightarrow R'; t) = \langle R | e^{-t\hat{H}} | R' \rangle$$

Roughly the probability density of going from R_0 to R in "time" t . (but is it a probability?)

$$-\frac{\partial G(R_0 \rightarrow R; t)}{\partial t} = \hat{H}G(R_0 \rightarrow R; t)$$

- Properties:

$$G(R_0 \rightarrow R; 0) = \mathbf{d}(R_0 - R)$$

$$G(R_0 \rightarrow R; t) = \sum_a \mathbf{f}_a^*(R_0) \mathbf{f}_a(R)$$

Froebinius Theorem

When can we consider the wavefunction as a probability?
 First how about the Green's function?

$$G(R_0 \rightarrow R; 0) = \mathbf{d}(R_0 - R) \geq 0$$

Trotter's theorem implies it continues to
 be positive at all times.

$$G(R_0 \rightarrow R; t) \geq 0$$

But if we start with a non-negative function it will stay
 non-negative, and can be interpreted as a p.d.f.

Not true for all Hamiltonians (require off-diagonal matrix
 elements to be non-positive.) (not pseudopotentials, not
 magnetic fields.)

Only true for the bosonic ground state.

Monte Carlo process

- Now consider the variable "t" as a continuous time (*it is really imaginary time*).
- Take derivative with respect to time to get evolution.
- This is a diffusion + branching process.
- Justify in terms of Trotter's theorem.

Requires interpretation of the wavefunction as a probability density.

But is it? Only in the boson ground state. Otherwise there are nodes. Come back to later.

$$-\frac{\partial \mathbf{y}(R, t)}{\partial t} = (H - E_T) \mathbf{y}(R, t)$$

$$H = -\sum_i \frac{\hbar^2}{2m_i} \nabla_i^2 + V(R)$$

$$\left\{ \begin{array}{l} -\frac{\partial \mathbf{y}(R, t)}{\partial t} = -\sum_i \frac{\hbar^2}{2m_i} \nabla_i^2 \mathbf{y}(R, t) \\ -\frac{\partial \mathbf{y}(R, t)}{\partial t} = (V(R) - E_T) \mathbf{y}(R, t) \end{array} \right.$$

Trotter's theorem

- How do we find the solution of:

$$\frac{d\hat{r}}{dt} = (A + B)\hat{r}$$

- The operator solution is:

$$\hat{r} = e^{(A+B)t}$$

- Trotter's theorem (1959):

$$\hat{r} = \lim_{n \rightarrow \infty} \left[e^{\frac{t}{n}A} e^{\frac{t}{n}B} \right]^n$$

- Assumes that A, B and A+B are reasonable operators.

$$\left\langle R_0 \left[e^{\frac{t}{n}A} e^{\frac{t}{n}B} \right]^n \middle| R_n \right\rangle = \left\langle R_0 \middle| e^{\frac{t}{n}A} \middle| R'_1 \right\rangle \left\langle R'_1 \middle| e^{\frac{t}{n}B} \middle| R_1 \right\rangle \dots \left\langle R_{n-1} \middle| e^{\frac{t}{n}A} \middle| R'_n \right\rangle \left\langle R'_n \middle| e^{\frac{t}{n}B} \middle| R_n \right\rangle$$

- This means we just have to figure out what each operator does independently and then alternate their effect. This is rigorous in the limit as $n \rightarrow \infty$.
- In the DMC case A is diffusion operator, B is a branching operator.
- Just like "molecular dynamics" At small time we evaluate each operator separately.

Evaluation of kinetic density matrix

$$\langle r | e^{-t\hat{T}} | r' \rangle = \sum_a f_a^*(r) f_a(r') e^{-tE_a}$$

In PBC eigenfunctions of $\hat{T} = \frac{1}{\sqrt{\Omega}} e^{-i\vec{k}\vec{r}}$

and eigenvalues are $\mathbf{1}k^2$

$$\langle r | e^{-t\hat{T}} | r' \rangle = \sum_k \frac{1}{\Omega} e^{-i\vec{k}\vec{r}} e^{i\vec{k}\vec{r}'} e^{-\mathbf{1}k^2 t}$$

convert to an integral

$$\langle r | e^{-t\hat{T}} | r' \rangle = \frac{1}{(2\mathbf{p})^3} \int d\mathbf{k} e^{i\vec{k}(\vec{r}'-\vec{r})-\mathbf{1}k^2 t} = (4\mathbf{p}\mathbf{1}t)^{-3/2} e^{-(r-r')^2/4\mathbf{1}t}$$

Danger: makes assumption about boundaries and statistics.

This is a diffusion process.

Putting this together

$$\hat{r} = e^{-b(\hat{T} + \hat{V})}$$

- n is number of time slices.
- τ is the "time-step"

$$\hat{r} = \lim_{n \rightarrow \infty} \left[e^{-t\hat{T}} e^{-t\hat{V}} \right]^n$$

$$t = b/n$$

- V is "diagonal"

$$\langle r | e^{-t\hat{T}} | r' \rangle = (4\pi t)^{-3/2} e^{-(r-r')^2/4t}$$

$$\langle r | e^{-t\hat{V}} | r' \rangle = d(r-r') e^{-tV(r)}$$

$$\langle R_0 e^{-nt\hat{H}} R_n \rangle \sim \langle R_0 | e^{-t\hat{T}} | R_1 \rangle e^{-tV(R_1)} \dots \langle R_{n-1} | e^{-t\hat{T}} | R_n \rangle e^{-tV(R_n)}$$

- Error at finite n comes from commutator is roughly: $e^{-\frac{t^2}{2}[\hat{T}, \hat{V}]}$
- Diffusion preserves normalization but potential does not!

Basic DMC algorithm

- Construct an ensemble (population P(0)) sampled from the trial wavefunction. $\{R_1, R_2, \dots, R_P\}$
- Go through ensemble and diffuse each one (timestep τ)

$$R'_k = R_k + \sqrt{2\tau} \mathbf{1} z(t) \leftarrow \begin{matrix} \text{ndrn} \\ \text{uprn} \end{matrix}$$
- number of copies = $\frac{e^{-t(V(R) - E_T)} + u}{\text{floor function}}$
- Trial energy E_T adjusted to keep population fixed.

$$E_0 = \lim_{t \rightarrow \infty} \frac{\int dR H f(R, t)}{\int dR f(R, t)} \approx \langle V(R) \rangle_{f(\infty)}$$

- Problems:
 - Branching is uncontrolled
 - What do we do about fermi statistics?

Population Bias

- Having the right trial energy guarantees that population will on the average be stable, but fluctuations will always cause the population to either grow too large or too small.
- Various ways to control the population
- Suppose P_0 is the desired population and $P(t)$ is the current population. How much do we have to adjust E_T to make $P(t+T)=P_0$?

$$P(t+T) = e^{-T(-dE_T)} P(t) = P_0$$

$$dE_T = \frac{\ln(P(t) / P_0)}{T}$$

$$E_T = E_{T0} + k \ln(P / P_0)$$

- There will be a (small) bias in the energy caused by a limited population.

Importance Sampling

Kalos 1970, Ceperley 1979

- Why should we sample the wavefunction? The physically correct pdf is $|\phi|^2$.
- Importance sample (multiply) by trial wave function.

$$f(R, t) \equiv \mathbf{y}_T(R) \mathbf{f}(R, t) \quad \lim_{t \rightarrow \infty} f(R, t) \equiv \mathbf{y}_T(R) \mathbf{f}_0(R)$$

$$-\frac{\partial f(R, t)}{\partial t} = \mathbf{y}_T(R) H [f(R, t) / \mathbf{y}_T(R)] \quad \text{Commute through H}$$

$$-\frac{\partial f(R, t)}{\partial t} = -I \nabla^2 f - I \nabla (2 f \nabla \ln \mathbf{y}_T(R)) + (\mathbf{y}_T^{-1} H \mathbf{y}_T) f(R, t)$$

- Evolution = diffusion + drift + branching
- Use accept/reject step for more accurate evolution.
make acceptance ratio > 99% . Determines time step.
 - We have three terms in the evolution equation. Trotter's theorem still applies.

Brownian Dynamics

Consider a big molecule in a solvent. In the high viscosity limit the "master equation" (Smoluchowski or Fokker-Planck eq.) is:

$$\frac{\partial \mathbf{r}(R, t)}{\partial t} = D \nabla^2 \mathbf{r}(R, t) - \mathbf{b} D \nabla [F(R) \mathbf{r}(R, t)]$$

$$R(t + \mathbf{t}) = R(t) + \mathbf{t} \mathbf{b} D F(R(t)) + \mathbf{h}(t)$$

$$\langle \mathbf{h}(t) \rangle = 0 \quad \langle \mathbf{h}(t)^2 \rangle = 2\mathbf{t} D$$

$$G(R \rightarrow R') = c \exp\left(-\frac{(R' - R - \mathbf{b} D \mathbf{t} F(R))^2}{2D\mathbf{t}}\right)$$

Also the equation for Diffusion Quantum Monte Carlo without branching. Borrow rejection technique developed for that.

Green's function for a gradient

What is Green's function for the operator?

$$\vec{F} \vec{\nabla}$$

variables separate to 1D problems

Evolution equation for Green's function:

$$\frac{\partial G(x, t)}{\partial t} = -F \frac{\partial G(x, t)}{\partial x} \quad \text{solution } G(x, t) = h(x - Ft)$$

This operator just causes probability distribution to drift in the direction of F.

Smoluchowski equation for Brownian motion it was the effect of gravitational field on the motion of colloids.

In practice, we limit the gradient so the walk is not pushed too far.

- To the pure diffusion algorithm we have added a drift step that pushes the random walk in directions of increasing trial function: $R' = R + 2\mathbf{I}t\nabla \ln \mathbf{y}_T(R)$
- Branching is now controlled by the local energy

$$E_L(R) - E_T = \mathbf{y}^{-1}(R) \widehat{H} \mathbf{y}(R) - E_T$$

- Because of zero variance principle, fluctuations are controlled.
- Cusp condition can limit infinities coming from singular potentials.
- We still determine E_T by keeping asymptotic population stable.

$$E_0 = \lim_{t \rightarrow \infty} \frac{\int dR \mathbf{f}(R, t) H \mathbf{y}_T(R)}{\int dR \mathbf{f}(R, t)} \approx \langle E_y(R) \rangle_{f(\infty)}$$

- Must have accurate "time" evolution. Adding accept/reject step is a major improvement.

How do we deal with fermi statistics?

- Importance sampled Green's function:

$$G(R \rightarrow R') = \frac{\mathbf{y}(R')}{\mathbf{y}(R)} \langle R | e^{-tH} | R' \rangle$$

- Exact property of DMC Green's function

$$|\Psi(R)|^2 G(R \rightarrow R') = |\Psi(R')|^2 G(R' \rightarrow R)$$

- We enforce detailed balance to decrease time step errors.

$$A(s \rightarrow s') = \min \left[1, \frac{G(s' \rightarrow s) |\mathbf{y}(s')|^2}{G(s \rightarrow s') |\mathbf{y}(s)|^2} \right]$$


- VMC satisfies detailed balance.
- Typically we choose time step to have 99% acceptance ratio.
- Method gives exact result if either time step is zero or trial function is exact.

Schematic of DMC

Ensemble evolves according to

- Diffusion
- Drift
- branching

ensemble



Mixed estimators

- Problem is that PMC samples the wrong distribution.

$$\langle A \rangle_M \equiv \frac{\int d\mathbf{R} \mathbf{y}^*(R) A f(R)}{\int d\mathbf{R} \mathbf{y}^*(R) f(R)}$$

- OK for the energy
- Linear extrapolation helps correct this systematic error

$$\langle A \rangle_o \equiv \frac{\int d\mathbf{R} f^*(R) A f(R)}{\int d\mathbf{R} f^*(R) f(R)}$$

$$\langle A \rangle_V \equiv \frac{\int d\mathbf{R} \mathbf{y}^*(R) A \mathbf{y}(R)}{\int d\mathbf{R} \mathbf{y}^*(R) \mathbf{y}(R)}$$

$$\langle A \rangle_o \approx 2\langle A \rangle_M - \langle A \rangle_V + O((f-y)^2)$$


- Other solutions:
 - Maximum overlap
 - Forward walking
 - Reptation/path integrals

$$\langle A \rangle_o \approx \frac{\langle A \rangle_M^2}{\langle A \rangle_V} + O((f-y)^2) \text{ for the density}$$

$$\langle A \rangle_M = \langle A \rangle_V \Rightarrow \int d\mathbf{R} (f-y)^2 \text{ minimized wrt } A$$

Other projector functions can be used

$$G(E) = \begin{cases} e^{-t(E-E_T)} & \text{Diffusion MC} \\ [1+t(E-E_T)]^{-1} & \text{Green's Function MC} \\ [1-t(E-E_T)] & \text{Power MC} \end{cases}$$



$G(E_T) = 1 \Rightarrow$ ground state remains after many iterations

$$t = - \left. \frac{dG}{dE} \right|_0 = \text{time step}$$

for all 3 cases: $\lim_{n \rightarrow \infty} G(E)^n = e^{-nt(E-E_T)}$

- Common effect on long-time (iteration) limit.
- 3rd choice generates a Krylov sequence. Only works for bounded spectra such as a lattice model.

Green's Function Monte Carlo

Kalos, Levesque, Verlet Phys. Rev. A9, 2178 (1974).

- It is possible to make a zero time-step-error method
- Works with the integral formulation of DMC

$$G(R, R') = \left\langle R \left[1 + t(H - E_T) \right]^{-1} \right| R' \rangle = \int_0^\infty \frac{d\mathbf{b}}{t} e^{-\mathbf{b} \left(\frac{1}{t} + H - E_T \right)}$$

- Sample time-step from Poisson distribution
- Express operator in a series expansion and sample the terms stochastically.

$$G(R, R') = H(R, R') + \int dR'' G(R, R'') K(R'', R')$$

- Recent Revival: "Continuous time Monte Carlo" for lattice models.

Exact fermion methods

- How can we do fermion simulations? The initial condition can be made real but not positive (for more than 1 electron in the same spin state)
- In *transient estimate* or *released-node* methods one carries along the sign as a weight and samples the modulus.

$$\mathbf{f}(t) = e^{-(\hat{H}-E_T)t} \text{sign}(\mathbf{f}(R,0)) |\mathbf{f}(R,0)|$$

- Do not forbid crossing of the nodes, but carry along sign when walks cross.
- What's wrong with node release:
 - Because walks don't die at the nodes, the computational effort increases (bosonic noise)
 - The signal is in the cancellation which dominates

Monte Carlo can add but not subtract

Variational-Projector Approach (Transient Estimate)

$$\Psi(\mathbf{b}) = e^{-\frac{\mathbf{b}}{2}H} \Psi$$


$$Z(\mathbf{b}) = \langle \Psi(\mathbf{b}) \Psi(\mathbf{b}) \rangle = \langle \Psi e^{-\mathbf{b}H} \Psi \rangle = \int dR_0 \dots dR_p \Psi(R_0) \langle R_0 e^{-tH} R_1 \rangle \dots \langle R_{p-1} e^{-tH} R_p \rangle \Psi(R_p)$$

$$E(\mathbf{b}) = \frac{\langle \Psi(\mathbf{b}) H \Psi(\mathbf{b}) \rangle}{\langle \Psi(\mathbf{b}) \Psi(\mathbf{b}) \rangle} = \langle E_L(R_0) \rangle_{\mathbf{b}} \quad t = \frac{\mathbf{b}}{p}$$

$$-\frac{dE(\mathbf{b})}{d\mathbf{b}} = \mathbf{s}^2(\mathbf{b}) = \frac{\langle \Psi(\mathbf{b}) (H - E(\mathbf{b}))^2 \Psi(\mathbf{b}) \rangle}{\langle \Psi(\mathbf{b}) \Psi(\mathbf{b}) \rangle} = \langle E_L(R_0) E_L(R_p) \rangle_{\mathbf{b}} - E(\mathbf{b})^2 > 0$$

- $\psi(\beta)$ converges to the exact ground state
- E is an upper bound converging to the exact answer monotonically because σ its derivative is positive.


$$\begin{aligned} Z(\mathbf{b}) &= \int dR_0 \dots dR_p |\Psi(R_0) \langle R_0 e^{-tH} R_1 \rangle \dots \langle R_{p-1} e^{-tH} R_p \rangle \Psi(R_p)| \mathbf{s}(R_0) \mathbf{s}(R_p) \\ &= \langle \mathbf{s}(R_0) \mathbf{s}(R_p) \rangle \end{aligned}$$



Model fermion problem: Particle in a box

Symmetric potential: $V(\mathbf{r}) = V(-\mathbf{r})$

Antisymmetric state: $\phi(\mathbf{r}) = -\phi(-\mathbf{r})$




Sign of walkers fixed by initial position. They are allowed to diffuse freely.
 $f(r)$ = number of positive-negative walkers. Node is dynamically established by diffusion process. (cancellation of positive and negative walkers.)


$$\langle E(t) \rangle = \frac{\sum \mathbf{s}(0) \mathbf{s}(t) E(t)}{\sum \mathbf{s}(0) \mathbf{s}(t)}$$

Scaling in Released-Node

Initial distribution



Later distribution




- At any point, positive and negative walkers will tend to cancel so the signal is drawn out by the fluctuations.
- Signal/noise ratio is : $e^{-t|E_F - E_B|}$ t =projection time
 E_F and E_B are Fermion, Bose energy (proportional to N)
- Converges but at a slower rate. Higher accuracy, larger t .
- For general excited states:
Exponential complexity! $\text{CPUtime} \propto e^{-2(1+\frac{E_F}{E_g})t} \approx e^{-2N\frac{E_F}{E_g}t}$
- Not a fermion problem but an excited state problem.
- Cancellation is difficult in high dimensions.

Exact fermion calculations

- Possible for the electron gas for up to 60 electrons.
- 2DEG at $r_s=1$ $N=26$
- Transient estimate calculation with SJ and BF-3B trial functions.

$$\langle \Psi_T | e^{-tH} | \Psi_T \rangle$$



General statement of the “fermion problem”

- Given a system with **N** fermions and a known Hamiltonian and a property **O**. (usually the energy).
- How much time **T** will it take to estimate **O** to an accuracy **e**? How does **T** scale with **N** and **e**?
- If you can map the quantum system onto an equivalent problem in classical statistical mechanics then:

$$T \propto N^a e^{-2} \quad \text{With } 0 < a < 4$$

This would be a “solved” quantum problem!

- All approximations must be controlled!
- Algebraic scaling in N!

e.g. properties of Boltzmann or Bose systems in equilibrium.

“Solved Problems”

- 1-D problem. (simply forbid exchanges)
- Bosons and Boltzmannons at any temperature
- Some lattice models: Heisenberg model, 1/2 filled Hubbard model on bipartite lattice (Hirsch)
- Spin symmetric systems with purely attractive interactions: $u < 0$ Hubbard model, nuclear Gaussian model.
- Harmonic oscillators or systems with many symmetries.
- Any problem with $\langle i|H|j \rangle \leq 0$
- Fermions in special boxes
- Other lattice models

- Kalos and coworkers have invented a pairing method but it is not clear whether it is approximation free and stable.

The sign problem

- The fermion problem is intellectually and technologically very important.
- Progress is possible but **danger**-the problem may be more subtle than you first might think. New ideas are needed.
- No fermion methods are perfect but QMC is competitive with other methods and more general.
- The fermion problem is one of a group of related problems in quantum mechanics (**e.g** dynamics).
- Feynman argues that general many-body quantum simulation is exponentially slow on a classical computer.
- Maybe we have to “solve” quantum problems using “analog” quantum computers: programmable quantum computers that can emulate any quantum system.

Fixed-node method

- Initial distribution is a pdf.
It comes from a VMC simulation. $f(R,0) = |\mathbf{y}_T(R)|^2$
- Drift term pushes walks away from the nodes.
- Impose the condition: $\mathbf{f}(R) = 0$ when $\mathbf{y}_T(R) = 0$.
- This is the fixed-node BC
- Will give an upper bound to the exact energy, the best upper bound consistent with the FNBC. $E_{FN} \geq E_0$
 $E_{FN} = E_0$ if $\mathbf{f}_0(R)\mathbf{y}(R) \geq 0$ all R
- $f(R,t)$ has a discontinuous gradient at the nodal location.
- Accurate method because Bose correlations are done exactly.
- Scales well, like the VMC method, as N^3 . Classical complexity.
- Can be generalized from the continuum to lattice finite temperature, magnetic fields, ...
- **One needs trial functions with accurate nodes.**

Proof of fixed-node theorem

- Suppose we solve S.E. in a subvolume V determined by the nodes of an antisymmetric trial function.

$$\hat{H} \mathbf{f}_{FN} = E_{FN} \mathbf{f}_{FN} \quad \text{inside } V$$

Extend the solution to all space with the permutation operator.

$$\hat{\mathbf{f}}_{FN}(R) \equiv \frac{1}{N!} \sum_P (-1)^P \mathbf{f}_{FN}(PR)$$

Inside a given sub-volume only permutations of a given sign (\pm) contribute.

Hence the extend solution is non-zero.

Evaluate the variational energy the extended trial function.

$$E_0 \leq \frac{\sum_{PP'} (-1)^{P+P'} \int dR \mathbf{f}_{FN}^*(PR) \hat{H} \mathbf{f}_{FN}(P'R)}{\sum_{PP'} (-1)^{P+P'} \int dR \mathbf{f}_{FN}^*(PR) \mathbf{f}_{FN}(P'R)} = E_{FN} \leq E_{VMC}$$

Edges of volumes do not contribute to the integral

since the extend solution vanishes there.

Nodal Properties

If we know the sign of the exact wavefunction (the nodes), we can solve the fermion problem with the fixed-node method.

- If $\phi(\mathbf{R})$ is real, nodes are $\phi(\mathbf{R})=0$ where \mathbf{R} is the $3N$ dimensional vector.
- Nodes are a $3N-1$ dimensional surface. (Do not confuse with single particle orbital nodes!)
- Coincidence points $\mathbf{r}_i = \mathbf{r}_j$ are $3N-3$ dimensional hyper-planes
- In 1 spatial dimension these "points" exhaust the nodes: *fermion problem is easy to solve in 1D with the "no crossing rule."*
- Coincidence points (and other symmetries) only constrain nodes in higher dimensions, they do not determine them.
- The nodal surfaces define nodal volumes. **How many nodal volumes are there? Conjecture:** there are typically only 2 different volumes (+ and -) except in 1D. (but only demonstrated for free particles.)

Nodal Picture: 2d slice thru 322d space

- Free electron
- Other electrons

- Nodes pass thru their positions
- Divides space into 2 regions
- Wavelength given by interparticle spacing




Fig. 3. A 2D cross section of the ground-state wave function of 161 free (polarized) fermions in a periodic square. All 161 particle positions were sampled using variational Monte Carlo from $\phi(R)^2$. The filled circle indicates the original position of the first particle. The other 160 particles are fixed at positions indicated by the open circles, and nodes of the wave function as a function of the position of the first particle are plotted. The resolution of the contouring program is approximately half of the line scale shown around the border of the plot.

SPIN?

- How do we treat spin in QMC?
- For extended systems we use the S_z representation.
- We have a fixed number of up and down electrons and we antisymmetrize among electrons with the same spin.
- This leads to 2 Slater determinants.
- For a given trial function, its real part is also a trial function (but it may have different symmetries), for example momentum

$$(e^{ikr}, e^{-ikr}) \text{ or } (\cos(kr), \sin(kr))$$

- For the ground state, without magnetic fields, spin-orbit interaction we can always work with real functions.
- However, in some cases it may be better to work with complex functions.

Fixed-Phase method

Ortiz, Martin, DMC 1993

- Generalize the FN method to complex trial functions: $\Psi(R) = e^{-U(R)}$
- Since the Hamiltonian is Hermitian, the variational energy is real:

$$E_V = \frac{\int dR e^{-2\Re U(R)} \left[V(R) + \mathbf{I} \nabla^2 U(R) - \mathbf{I} [\Re \nabla U(R)]^2 + \mathbf{I} [\Im \nabla U(R)]^2 \right]}{\int dR e^{-2\Re U(R)}}$$

- We see only one place where the energy depends on the phase of the wavefunction.
- If we **fix the phase**, then we add this term to the potential energy. In a magnetic field we get also the vector potential.

$$\text{effective potential} = V(R) + \sum_i \mathbf{I}_i [A(r_i) + \Im \nabla_i U(R)]^2$$

- We can now do VMC or DMC and get upper bounds as before.
- The imaginary part of the local energy will not be zero unless the right phase is used.
- Used for twisted boundary conditions, magnetic fields, vortices, phonons, spin states, ...

Problem with core electrons


- Bad scaling in both VMC and DMC
- In VMC, energy fluctuations from core dominate the calculation
- In DMC, time step will be controlled by core dynamics
- Solution is to eliminate core states by a pseudopotential
- Conventional solution: semi-local form

$$\langle r | \hat{v}_{e\text{-core}} | r' \rangle = v_{\text{local}}(r) \mathbf{d}(r - r') + \sum_l v_l(r) P_l(\cos(r \cdot r'))$$

- Ensures that valence electrons go into well defined valence states with the wavefunction and energy for each angular momentum state prescribed.
- PP is non-local: OK for VMC. Leads to an extra MC integral. But DMC uses a locality approximation and good trial functions. **Extra approximation.**

Summary of T=0 methods:

Variational(VMC), Fixed-node(FN), Released-node(RN)



Problems with projector methods

- Fixed-node is a super-variational method
- DMC dynamics is determined by Hamiltonian
- Zero-variance principle allows very accurate calculation of ground state energy if trial function is good.
- Projector methods need a trial wavefunction for accuracy. They are essentially methods that perturb from the trial function to the exact function. (Note: if you don't use a trial function, you are perturbing from the ideal gas)
- Difficulty calculating properties other than energy. We must use "extrapolated estimators" or "forward walking".

$$f(R, \infty) = \mathbf{f}_0(R) \mathbf{y}_T(R) \text{ not } |\mathbf{f}_0(R)|^2$$

- Bad for phase transitions and finite temperature, complex systems.
- Path Integral MC solves some of these problems.