

Advanced School and Conference on Non-commutative Geometry

(9 - 27 August 2004)

Noncommutative deformations of modules

O.A. Laudal

Institute of Mathematics
University of Oslo
P.O. Box 1053 Blindern
N-0316 Oslo
Norway

These are preliminary lecture notes, intended only for distribution to participants

NONCOMMUTATIVE DEFORMATIONS OF MODULES.

O.A.LAUDAL

University of Oslo

Contents:

1. Homological preparations
2. The noncommutative deformation functor
3. Noncommutative modular deformations
4. The generalized Burnside theorem
5. Filtered modules and iterated extensions
6. Noncommutative algebraic geometry
7. The smooth locus of an affine noncommutative scheme

§1. Homological preparations.

Exts and Hochschild cohomology. Let k be a (usually algebraically closed) field, and let A be a k -algebra. Denote by $A\text{-mod}$ the category of right A -modules and consider the exact forgetful functor

$$\pi : A\text{-mod} \longrightarrow k\text{-mod}$$

Given two A -modules M and N , we shall always use the identification

$$\sigma^i : Ext_A^i(M, N) \simeq HH^i(A, Hom_k(M, N)) \text{ for } i = 0, 1, 2,$$

where $Hom_k(M, N)$ is provided with the obvious left and right A -module structures. If L_* and F_* are A -free resolutions of M and N respectively, and if an element

$$\xi \in Ext_A^1(M, N)$$

is represented by the Yoneda cocycle,

$$\hat{\xi} = \{\xi_n\} \in \prod_n Hom_A(L_n, F_{n-1})$$

then $\sigma^1(\xi)$ is gotten as follows. Let σ be a k -linear section of the augmentation morphism

$$\rho : L_0 \longrightarrow M$$

and let for every $a \in A$ and $m \in M$, $\sigma(ma) - \sigma(m)a = d_0(x)$. Put,

$$\sigma^1(\hat{\xi})(a, m) = -\mu(\xi_1(x))$$

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}\text{-}\mathcal{T}\mathcal{E}\mathcal{X}$

where

$$\mu : F_0 \longrightarrow N$$

is the augmentation morphism of F_* . Then,

$$\sigma^1(\hat{\xi}) \in \text{Der}_k(A, \text{Hom}_k(M, N))$$

and its class in $HH^1(A, \text{Hom}_k(M, N))$ equals $\sigma^1(\xi)$.

Recall the spectral sequence associated to a change of rings. If $\pi : A \longrightarrow B$ is a surjective homomorphism of commutative k -algebras, M a B -module and N an A -module, then $\text{Ext}_A^*(M, N)$ is the abutment of the spectral sequence given by,

$$E_2^{p,q} = \text{Ext}_B^p(M, \text{Ext}_A^q(B, N)).$$

There is an exact sequence,

$$0 \longrightarrow E_2^{1,0} \longrightarrow \text{Ext}_A^1(M, N) \longrightarrow E_2^{0,1} \longrightarrow E_2^{2,0},$$

which, for a B -module N , considered as an A -module, implies the exactness of

$$\begin{aligned} 0 &\longrightarrow \text{Ext}_B^1(M, N) \longrightarrow \text{Ext}_A^1(M, N) \\ &\longrightarrow \text{Hom}_B(M, \text{Hom}_B(I/I^2, N)) \longrightarrow \text{Ext}_B^2(M, N) \end{aligned}$$

where $I = \ker \pi$. The corresponding exact sequence,

$$\begin{aligned} 0 &\longrightarrow HH^1(B, \text{Hom}_k(M, N)) \longrightarrow HH^1(A, \text{Hom}_k(M, N)) \\ &\longrightarrow \text{Hom}_{A \otimes A^{op}}(I, \text{Hom}_k(M, N)) \end{aligned}$$

in the noncommutative case is induced by the sequence

$$\begin{aligned} 0 &\longrightarrow \text{Der}_k(B, \text{Hom}_k(M, N)) \longrightarrow \text{Der}_k(A, \text{Hom}_k(M, N)) \\ &\longrightarrow \text{Hom}_{A \otimes A^{op}}(I, \text{Hom}_k(M, N)). \end{aligned}$$

Notice that in general we do not know that the last morphism is surjective. This, however, is true if $B = A/\text{rad}(A)$, where $\text{rad}(A)$ is the radical of A , and A is a finite dimensional, i.e. an artinian, k -algebra. In this case, B is semisimple and the surjectivity above follows from the Wedderburn-Malcev theorem. Notice also that in the commutative case,

$$\text{Hom}_{A \otimes A^{op}}(I, \text{Hom}_k(M, N)) \simeq \text{Hom}_B(I/I^2, \text{Hom}_B(M, N))$$

as it must, since for $\phi \in \text{Hom}_{A \otimes A^{op}}(I, \text{Hom}_k(M, N))$, $a \in A$, and $i \in I$, $ai = ia$, and therefore

$$a\phi(i) = \phi(ai) = \phi(ia) = \phi(i)a, \text{ i.e. } \phi(i) \in \text{Hom}_B(M, N).$$

This implies that for $B = A/\mathfrak{p}$, $M = A/\mathfrak{p}$, $N = A/\mathfrak{q}$, where $\mathfrak{p} \subseteq \mathfrak{q}$ are (prime) ideals of A ,

$$\text{Ext}_A^1(A/\mathfrak{p}, A/\mathfrak{q}) \simeq \text{Hom}_A(\mathfrak{p}/\mathfrak{p}^2, A/\mathfrak{q})$$

and, in particular

$$\text{Ext}_A^1(A/\mathfrak{q}, A/\mathfrak{q}) \simeq \text{Hom}_A(\mathfrak{q}/\mathfrak{q}^2, A/\mathfrak{q}) = N_{\mathfrak{q}},$$

the normal bundle of $V(\mathfrak{q})$ in $\text{Spec}(A)$. If $\mathfrak{q} \subset \mathfrak{p}$ and $\mathfrak{q} \neq \mathfrak{p}$ we find,

$$\text{Ext}_A^1(A/\mathfrak{p}, A/\mathfrak{q}) \simeq \text{Ext}_{A/\mathfrak{q}}^1(A/\mathfrak{p}, A/\mathfrak{q}).$$

In [La 1], chapter 1., we considered the cohomology of a category \underline{c} with values in a bifunctor, i.e. in a functor defined on the category $\text{mor}_{\underline{c}}$ of morphisms of \underline{c} . Recall that a morphism between the objects ψ and ψ' is a commutative diagram,

$$\begin{array}{ccc} c_1 & \xrightarrow{\psi} & c_2 \\ \uparrow & & \downarrow \\ c'_1 & \xrightarrow{\psi'} & c'_2. \end{array}$$

It is easy to see that this cohomology is an immediate generalization of the projective limit functor and its derivatives, or if one likes it better, the obvious generalization of the Hochschild cohomology of a ring. In fact, for every small category \underline{c} and for every bifunctor,

$$G : \underline{c} \times \underline{c} \longrightarrow \text{Ab}$$

contravariant in the first variable, and covariant in the second, one obtains a covariant functor,

$$G : \text{mor}_{\underline{c}} \longrightarrow \text{Ab}.$$

Consider now the complex,

$$D^*(\underline{c}, G)$$

where,

$$D^p(\underline{c}, G) = \prod_{c_0 \rightarrow c_1 \rightarrow \dots \rightarrow c_p} G(c_0, c_p)$$

where the indices are strings of morphisms $\psi_i : c_i \rightarrow c_{i+1}$ in \underline{c} , and the differential,

$$d^p : D^p(\underline{c}, G) \longrightarrow D^{p+1}(\underline{c}, G)$$

is defined as usual,

$$\begin{aligned} (d^p \xi)(\psi_1, \dots, \psi_i, \psi_{i+1}, \dots, \psi_{p+1}) &= \psi_1 \xi(\psi_2, \dots, \psi_{p+1}) \\ &+ \sum_{i=1}^p (-1)^i \xi(\psi_1, \dots, \psi_i \circ \psi_{i+1}, \dots, \psi_{p+1}) + (-1)^{p+1} \xi(\psi_1, \dots, \psi_p) \psi_{p+1}. \end{aligned}$$

As shown in [La 1], the cohomology of this complex is the higher derivatives of the projective limit functor $\lim_{\leftarrow \text{mor}_{\underline{c}}}^{(*)}$ applied to the covariant functor

$$G : \text{mor}_{\underline{c}} \longrightarrow \text{Ab}.$$

This is the "Hochschild" cohomology of the category \underline{c} , denoted

$$H^*(\underline{c}, G) := H^*(D^*(\underline{c}, G)).$$

Example 1.1. Let \underline{c} be a multiplicative subset of a ring R , considered as a category with one object, and let $\tilde{R} : \underline{c} \times \underline{c} \longrightarrow Ab$ be the functor, defined for $\psi, \psi' \in \underline{c}$, by $\tilde{R}(\psi, \psi') = \psi^* \psi'_*$, where ψ^* is left multiplication on R by ψ , and where ψ'_* is right multiplication on R by ψ' , then

$$H^0(\underline{c}, \tilde{R}) = \{\phi \in R \mid \phi\psi = \psi\phi \text{ for all } \psi \in \underline{c}\},$$

i.e. the commutant of \underline{c} in R .

Given a k -algebra A , and consider a subcategory \underline{c} of the category of right A -modules. Let, as above $\pi : \underline{c} \rightarrow k - mod$ be the forgetful-functor, and consider the bifunctor,

$$Hom_\pi : \underline{c} \times \underline{c} \longrightarrow k - mod$$

defined by

$$Hom_\pi(V_i, V_j) = Hom_k(V_i, V_j).$$

Put,

$$O_0(\underline{c}, \pi) := H^0(\underline{c}, Hom_\pi).$$

It is clear that $O_0(\underline{c}, \pi)$ is a k -algebra, and that there is a canonical homomorphism of k -algebras,

$$\eta_0(\underline{c}, \pi) : A \longrightarrow O_0(\underline{c}, \pi),$$

see §5.

Example 1.2. Let A be a commutative k -algebra of finite type, k algebraically closed, and let $Spec(A)$ be the subcategory of A -mod consisting of the modules A/\mathfrak{p} , where \mathfrak{p} runs through $Spec(A)$, the morphisms being only the obvious ones. It is easy to see that the homomorphism

$$\eta_0(Spec(A), \pi) : A \longrightarrow O_0(Spec(A), \pi)$$

identifies $A/rad(A)$ with $O_0(Spec(A), \pi)$. If $rad(A) = 0$ we even find an isomorphism,

$$\eta_0(Simp^*(A), \pi) : A \simeq O_0(Simp^*(A), \pi).$$

Here $Simp^*(A)$ is the subcategory of A -mod where the objects are A and the simple A -modules, A/\mathfrak{m} , i.e. the closed points of $Spec(A)$, and where the morphisms are the obvious quotient morphisms $A \rightarrow A/\mathfrak{m}$. $\eta_0(Simp^*(A), \pi)$ is, however not, in general, an isomorphism. This is easily seen when A is a local k -algebra. To remedy this situation we shall introduce and study a generalization $O(\underline{c}, \pi)$ of $O_0(\underline{c}, \pi)$ defined in terms of the noncommutative deformation theory, see the next §'s.

§2. Noncommutative deformations.

The category \underline{a}_r , test algebras and liftings of modules.

Let \underline{a}_r be the category of “ r -pointed” artinian k -algebras. An object R of \underline{a}_r is a diagram of morphism of artinian k -algebras,

$$\begin{array}{ccc} k^r & \xrightarrow{\iota} & R \\ & \searrow \cong & \downarrow \rho \\ & & k^r \end{array}$$

Put, $Rad(R) := ker\rho$, such that,

$$R/Rad(R) \simeq \prod_{j=1}^r k_j, \quad k_j \simeq k.$$

A morphism $\phi : R \rightarrow S$ of \underline{a}_r is a morphism of such diagrams inducing the identity on k^r , implying that the induced map,

$$k^r \simeq R/Rad(R) \rightarrow S/Rad(S) \simeq k^r$$

is the identity. Pick idempotents $e_i \in k^r \subseteq R$ such that

$$\sum_{i=1}^r e_i = 1, \quad e_i e_j = 0 \text{ if } i \neq j.$$

For every (i, j) , we shall consider the subspace $R_{ij} := e_i R e_j \subseteq R$, and the pairing

$$R_{ij} \otimes_k R_{jk} \rightarrow R_{ik}$$

given in terms of the multiplication in R .

Let

$$R' = (R_{ij})$$

be the matrix algebra, the elements of which are matrices of the form

$$(\alpha_{ij})$$

with $\alpha_{ij} \in R_{ij}$, $i, j = 1, \dots, r$. There is an obvious isomorphism of k -algebras

$$\phi : R \rightarrow R'$$

defined by

$$\phi(\alpha) = (e_i \alpha e_j).$$

identifying the sub k -algebra k^r of R with the algebra of diagonal $r \times r$ -matrices. Now, for any pair (i, j) , $i, j = 1, \dots, r$, consider the symbol ϵ_{ij} , and let's agree to put all products of such symbols equal to zero. Then we define the (i, j) -test algebra $R(i, j)$ as the matrix algebra

$$R(i, j) = k^r \oplus i \begin{pmatrix} & & j \\ 0 & \vdots & 0 \\ \cdots & k \cdot \epsilon_{ij} & \cdots \\ 0 & \vdots & 0 \end{pmatrix} \quad \text{for } i \neq j$$

$$R(i, i) = i \begin{pmatrix} & & i \\ k & \vdots & 0 \\ \cdots & k[\epsilon_{ii}] & \cdots \\ 0 & \vdots & k \end{pmatrix} \quad \text{for } i = j$$

Denote by $HH^*(A, -)$ the Hochschild cohomology of the k -algebra A . If W is an A -bimodule denote by $Der_k(A, W)$ the k -vectorspace of derivations of A in W . Thus $\psi \in Der_k(A, W)$ is a linear map from A to W such that $\psi(a_1 \cdot a_2) = a_1\psi(a_2) + \psi(a_1)a_2$.

In particular, any element $w \in W$ determines a derivation $i(w) \in Der_k(A, W)$ defined by $i(w)(a) = aw - wa$. There is an exact sequence

$$0 \rightarrow HH^0(A, W) \rightarrow W \rightarrow Der_k(A, W) \rightarrow HH^1(A, W) \rightarrow 0$$

If V_i, V_j are right A -modules, then

$$Hom_k(V_i, V_j)$$

is an A -bimodule. In fact if $\phi \in Hom_k(V_i, V_j)$, then $a\phi$ is defined by $(a\phi)(v) = \phi(va)$, and ϕa is defined by $(\phi a)(v) = \phi(v)a$.

Moreover, we know that

$$HH^0(A, Hom_k(V_i, V_j)) = Hom_A(V_i, V_j)$$

$$HH^1(A, Hom_k(V_i, V_j)) = Ext_A^1(V_i, V_j).$$

Fix a finite family $\mathcal{V} = \{V_i\}_{i=1}^r$ of right A -modules, and consider for every

$$\psi \in Der_k(A, Hom_k(V_i, V_j))$$

the left $R(i, j)$ -module and right A -module,

$$V_{ij}(\psi) = \begin{matrix} & & & & j \\ & & & & \vdots \\ i \left(\begin{array}{cccc} V_1 & & & \\ \cdots & V_i & \cdots & \epsilon_{ij}V_j \\ & & & V_j \\ & & & \vdots \\ & & & V_r \end{array} \right) & & & \end{matrix}$$

defined by

$$\begin{pmatrix} v_1 & & & & \\ & v_i & \epsilon_{ij}v'_j & & \\ & & v_j & & \\ & & & & v_r \end{pmatrix} \cdot a = \begin{pmatrix} v_1a & & & & \\ & v_ia & \epsilon_{ij}(\psi(a, v_i) + v'_ja) & & \\ & & v_ja & & \\ & & & & v_ra \end{pmatrix}$$

and the obvious left $R(i, j)$ -action. The $R(i, j)$ - and the A -action commute, therefore we have got a $R(i, j) \otimes A$ -module, such that

$$k^r \otimes_{R(i, j)} V_{ij}(\psi) \simeq \bigoplus_{i=1}^r V_i.$$

$V_{ij}(\psi)$ is called a lifting of \mathcal{V} to $R(i, j)$. It is easy to see that if ψ maps to zero in $HH^1(A, Hom_k(V_i, V_j)) = Ext_A^1(V_i, V_j)$ then the lifting $V_{ij}(\psi)$ is trivial, i.e. isomorphic to the trivial one. Conversely, if $V_{ij}(\psi)$ is trivial, then ψ maps to zero in $Ext_A^1(V_i, V_j)$.

The noncommutative deformation functor.

We are now ready to start the study of noncommutative deformations of the family $\mathcal{V} = \{V_i\}_{i=1}^r$. We shall assume that \mathcal{V} is a swarm, i.e. that for all $i, j = 1, 2, \dots, r$,

$$\dim_k \text{Ext}_A^1(V_i, V_j) < \infty.$$

Given an object $\rho : R = (R_{i,j}) \rightarrow k^r$ of \underline{a}_r , consider the left R -module $(R_{i,j} \otimes_k V_j)$. ρ defines a k -linear and left R -linear map,

$$\rho(R) : (R_{i,j} \otimes_k V_j) \rightarrow \bigoplus_{i=1}^r V_i,$$

inducing a homomorphism of R -endomorphism rings,

$$\tilde{\rho}(R) : (R_{i,j} \otimes_k \text{Hom}_k(V_i, V_j)) \rightarrow \bigoplus_{i=1}^r \text{End}_k(V_i).$$

The right A -module structure on the V_i 's is defined by a homomorphism of k -algebras,

$$\eta_0 : A \rightarrow \bigoplus_{i=1}^r \text{End}_k(V_i).$$

Definition 2.1. *The deformation functor*

$$\text{Def}_{\mathcal{V}} : \underline{a}_r \rightarrow \text{Sets}$$

is defined for every $R \in \underline{a}_r$, as the set,

$$\text{Def}_{\mathcal{V}}(R) \in \underline{\text{Sets}}$$

of isoclasses of homomorphisms of k -algebras,

$$\{\eta' : A \rightarrow (R_{i,j} \otimes_k \text{Hom}_k(V_i, V_j))\} / \sim$$

such that,

$$\tilde{\rho}(R) \circ \eta' = \eta_0,$$

where the equivalence relation \sim is defined by inner automorphisms in the k -algebra

$$\text{End}_R((R_{i,j} \otimes_k V_j)) = (R_{i,j} \otimes_k \text{Hom}_k(V_i, V_j)).$$

Any such isoclass $\tilde{\eta}'$ will be called a deformation or a lifting of V to R , and usually denoted V_R .

One easily proves that $\text{Def}_{\mathcal{V}}$ has the same properties as the ordinary deformation functor.

Let $\pi : R \rightarrow S$ be a morphism of \underline{a}_r , such that $\text{Rad}(R) \cdot \ker \pi = 0$. Morphisms like this will be called *small*. If $V_R \in \text{Def}_{\mathcal{V}}(R)$ it is easy to see that $V_S := S \otimes_R V_R \in \text{Def}_{\mathcal{V}}(S)$ and that $\bar{V} = \ker\{V_R \rightarrow S \otimes_R V_R\}$ is, as a left R -module, an $R/\text{Rad}(R) = k^r$ -module. Put $\ker \pi = (K_{ij})$, then $\bar{V} = (\bar{V}_{ij})$ where $\bar{V}_{ij} = K_{ij} \otimes_k V_j$.

Consider now the k -vector spaces

$$E_{ij}^d = \text{Ext}_A^d(V_i, V_j)^*$$

i.e. the dual k -vectorspaces of $\text{Ext}_A^d(V_i, V_j)$, and consider the k -algebra of matrices,

$$T_2^d = \begin{pmatrix} k & & 0 \\ & \ddots & \\ 0 & & k \end{pmatrix} + (\epsilon_{ij} E_{ij}^d)$$

where as above, we assume all products of the ϵ_{ij} 's are equal to zero. Now let for every $i, j = 1, \dots, r$, and $d = 1, 2$,

$$\left\{ t_{ij}^d(\ell) \right\}_{\ell=1}^{e_{ij}^d}$$

be a basis of E_{ij}^d , and let $\{\psi_{ij}^d(\ell)\}_{\ell=1}^{e_{ij}^d}$ be the dual basis. Thus $e_{ij}^k = \dim_k E_{ij}^k$. Consider the k -algebra

$$T^d = \begin{pmatrix} k & & 0 \\ & \ddots & \\ 0 & & k \end{pmatrix} + (\tilde{E}_{ij}^d)$$

freely generated as matrix algebra by the generators $\left\{ t_{ij}^d(\ell) \right\}_{\ell=1}^{e_{ij}^d}$. An element of \tilde{E}_{ij}^d is then a matrix where the elements are linear combinations of elements of the form:

$$\begin{aligned} \tau_{ij} &= t_{ij_1}^d(l_1) \otimes t_{j_1 j_2}^d(l_2) \otimes \cdots \otimes t_{j_{m-1} j_m}^d(l_m) \\ & \quad j = j_m, \quad 1 \leq l_s \leq e_{j_{s-1} j_s}^d, \quad 1 \leq j_s \leq r, \quad m \geq 1 \\ & \quad \text{of } E_{ij_1}^d \otimes E_{j_1 j_2}^d \otimes \cdots \otimes E_{j_{m-1} j}^d. \end{aligned}$$

Obviously

$$T_2^1 = T^1 / \text{Rad}(T^1)^2.$$

where $\text{Rad}(T^1)$ is the two-sided ideal of T^1 generated by (\tilde{E}_{ij}^1) .

Lemma 2.2. *Let R be an object of \underline{a}_r and suppose that there exists a surjective homomorphism*

$$\phi_2 : T_2^1 \rightarrow R / \text{Rad}(R)^2,$$

then there exists a surjective homomorphism

$$\phi : T^1 \rightarrow R$$

which lifts ϕ_2 .

Definition 2.3. *For every object R of \underline{a}_r , put*

$$T_R = (\text{Rad}(R) / \text{Rad}(R)^2)^*$$

and call it the tangent space of R .

Lemma 2.4. *Let $\phi : R \rightarrow S$ be a morphism of \underline{a}_r . Assume ϕ induces a surjective homomorphism*

$$\phi^1 : T_R^* \rightarrow T_S^*$$

(or an injective homomorphism on the tangent space level). Then ϕ is surjective.

Notice that if we pick any finite dimensional k -vectorspaces F_{ij} , then there is a unique maximal pro-algebra $F = F(F_{ij})$ in \underline{a}_r with tangent space

$$T_F \simeq (F_{ij}^*)$$

F is defined in the same way as T^d , above, with E^d replaced by F .

To prove the existence of a hull for the deformation functor Def_V the basic tool is the obstruction calculus, which in this case is easily established:

Proposition 2.5. *Suppose $R \xrightarrow{\phi} S$ is a surjective small morphism of \underline{a}_r , i.e. suppose $\ker\phi \cdot \text{Rad}(R) = 0$. Put $\ker\phi = (I_{ij})$. Consider any $V_S \in Def_V(S)$. Then there exists an obstruction*

$$o(\phi, V_S) \in (I_{ij} \otimes_k \text{Ext}_A^2(V_i, V_j))$$

which is zero if and only if there exists a lifting $V_R \in Def_V(R)$ of V_S . The set of isomorphism classes of such liftings is a pseudotorsor under

$$(I_{ij} \otimes_k \text{Ext}_A^1(V_i, V_j)).$$

Proof. As a k -vectorspace $V_R = (R_{ij} \otimes V_j)$ maps onto $V_S = (S_{ij} \otimes V_j)$. Since the right action of A commutes with the left S -action the action of an element $a \in A$ on V_S is uniquely given in terms of a family of k -linear maps,

$$a_{ij} : V_i \rightarrow S_{ij} \otimes V_j.$$

We may of course lift these to k -linear maps

$$\sigma(a)_{ij} : V_i \rightarrow R_{ij} \otimes V_j$$

inducing a lift of the action of each element of A on

$$\bigoplus_{j=1}^r S_{ij} \otimes V_j$$

to a k -linear action on

$$\bigoplus_{j=1}^r R_{ij} \otimes V_j.$$

The obstruction for this to be an A -module structure is, as usual, the Hochschild 2-cocycle

$$\psi^2(a, b) = \sigma(ab) - \sigma(a) \cdot \sigma(b) \in (I_{ij} \otimes_k \text{Hom}_k(V_i, V_j)).$$

The fact that this is a 2-cocycle follows from

- (1) $\sigma(c) \cdot \psi^2(a, b) = c \cdot \psi^2(a, b)$
- (2) $\psi^2(a, b) \cdot \sigma(c) = \psi^2(a, b) \cdot c$

and the obvious relation

$$\begin{aligned} d\psi^2(a, b, c) &= a\psi^2(b, c) - \psi^2(ab, c) + \psi^2(a, bc) - \psi^2(a, b) \cdot c \\ &= \sigma(a)(\sigma(bc) - \sigma(a)\sigma(c)) - (\sigma(abc) - \sigma(ab)\sigma(c)) + (\sigma(abc) - \sigma(a)\sigma(bc)) \\ &\quad - (\sigma(ab) - \sigma(a)\sigma(b))\sigma(c) \equiv 0 \end{aligned}$$

Suppose the class of ψ^2 in $(I_{ij} \otimes_k Ext_A^2(V_i, V_j))$ is zero. This means that $\psi^2 = d\phi$, where $\phi \in Hom_k(A, (I_{ij} \otimes_k Hom_k(V_i, V_j)))$, $\psi^2(a, b) = d\phi(a, b) = a\phi(b) - \phi(ab) + \phi(a)b$. Let $\sigma' = \sigma + \phi$ and consider,

$$\sigma'(ab) - \sigma'(a)\sigma'(b) = \sigma(ab) - \sigma(a)\sigma(b) + \phi(ab) - \sigma(a)\phi(b) - \phi(a)\sigma(b) - \phi(a)\phi(b).$$

Since the matrix $\phi(a)\phi(b) = 0$ as $I_{ij} \cdot I_{jk} = 0$, $\forall i, j, k$ and since $\sigma(a)\phi(b) = a\phi(b)$, $\phi(a)\sigma(b) = \phi(a)b$ for the same reason, we find that $\sigma'(ab) - \sigma'(a)\sigma'(b) = 0$, i.e. there is a lifting of the A -module action to $V_R = (R_{ij} \otimes V_j)$.

If we have given one A -module action σ on V_R lifting the action on V_S , then for any other σ' we may consider the difference

$$\sigma' - \sigma : A \rightarrow (I_{ij} \otimes_k Hom_k(V_i, V_j))$$

Consider

$$d(\sigma' - \sigma)(a, b) = a(\sigma'(b) - \sigma(b)) - (\sigma'(ab) - \sigma(ab)) + (\sigma'(a) - \sigma(a))b$$

As above we may substitute $\sigma'(a)$ for a and $\sigma(b)$ for b , and the expression becomes zero. Thus $\sigma' - \sigma = \bar{\xi}$ defines a class

$$\xi \in (I_{ij} \otimes_k Ext_A^1(V_i, V_j)).$$

If $\xi = 0$, then $\bar{\xi} = d\phi$, $\phi \in (I_{ij} \otimes_k Hom_k(V_i, V_j))$ such that $\sigma'(a) - \sigma(a) = a\phi - \phi a$. Let $\phi = (\phi_{ij})$, then ϕ_{ij} defines an isomorphism

$$\bar{\phi} = id + \phi : \bigoplus_j R_{ij} \otimes V_j \rightarrow \bigoplus_j R_{ij} \otimes V_j$$

lifting the identity of $\bigoplus_j S_{ij} \otimes V_j$. Moreover

$$\begin{aligned} \sigma(a)(id + \phi)(v_i) &= \sigma(a)v_i + a\phi(v_i) \\ &= \sigma'(a)(v_i) + \phi(av_i) = (id + \phi)\sigma'(a)(v_i) \end{aligned}$$

since $\phi(\sigma'(a)v_i) = \phi(av_i)$.

Therefore the A -module structures on

$$V_R = (R_{ij} \otimes V_j)$$

defined by σ and σ' are isomorphic. The rest is clear. \square

Theorem 2.6. *The functor $Def_{\mathcal{V}}$ has a prorepresentable hull, or a formal moduli of V , $H \in \underline{\mathcal{A}}_r$, together with a versal family*

$$\tilde{V} = (H_{i,j} \hat{\otimes} V_j) \in \varprojlim_{n \geq 1} Def_{\mathcal{V}}(H/Rad(H)^n)$$

such that the corresponding morphism of functors on $\underline{\mathcal{A}}_r$,

$$\rho : Mor(H, -) \rightarrow Def_{\mathcal{V}}$$

is smooth and an isomorphism on the tangent level. Moreover, H is uniquely determined by a set of matrix Massey products defined on subspaces,

$$D_n \subset \bigoplus_{p=2}^n Ext^1(V_i, V_{j_1}) \otimes \cdots \otimes Ext^1(V_{j_{p-1}}, V_j),$$

with values in $Ext^2(V_i, V_j)$.

Proof. Notice first that ρ being an isomorphism at the tangent level means that ρ is an isomorphism for all objects R of $\underline{\mathcal{A}}_r$ for which $Rad(R)^2 = 0$.

Word for word we may copy the proof (4.2) of [La 1], and the proof of [La 2]. In particular $H/Rad(H)^2 \simeq T_2^1$ and

$$Mor(H, R(i, j)) \simeq Hom_k(E_{ij}^1, k) \simeq Ext_A^1(V_i, V_j) \simeq Def_{\mathcal{V}}(R(i, j)).$$

Notice that the universal lifting of V to T_2^1 is the $T_2^1 \otimes_k A$ -module \tilde{V}_2

$$\begin{pmatrix} V_1 & & 0 \\ & \ddots & \\ 0 & & V_2 \end{pmatrix} + (E_{ij}^1 \otimes_k V_j)$$

with the obvious left T_2^1 -action and the right A -action defined as,

$$((1 \otimes v_i) \cdot a)_{ii} = 1 \otimes v_i \cdot a + \sum_{\ell} t_{i,j}^1(\ell) \otimes (\psi_{i,j}^1(\ell)(a, v_i))$$

where $v_i \in V_i$, and where $\{t_{ij}^1(\ell)\}_{\ell=1}^{e_{ij}}$ is the chosen basis of E_{ij}^1 . Recall that $\{\psi_{ij}^1(\ell)\}_{\ell=1}^{e_{ij}}$, the dual base, consists of elements $\psi_{ij}^1(\ell) \in Ext_A^1(V_i, V_j)$, which may be represented as elements of $Der_k(A, Hom_k(V_i, V_j))$.

To obtain H we kill obstructions for lifting \tilde{V}_2 successively, to $T_3^1 := T^1/Rad(T^1)^3, T_4^1$ etc. just like in the commutative case. The proof of the existence of a prorepresentable hull for $Def_{\mathcal{V}}$ can, of course, also be modeled on the classical proof of M.Schlessinger [Sch]. This has been carried out by Runar Ile, see [Ile]. \square

A general structure theorem for artinian k -algebras.

For every deformation $V_R \in Def_{\mathcal{V}}(R)$ there exists by definition an, up to inner automorphisms, unique homomorphism of k -algebras,

$$\eta_{V_R} : A \rightarrow End_R(V_R) = (R_{ij} \otimes Hom_k(V_i, V_j)).$$

Definition 2.7. Let $\mathcal{V} = \{V_i\}_{i=1}^r$ be any finite swarm of A -modules, and let $H := H(\mathcal{V})$ be the formal moduli for \mathcal{V} , and \tilde{V} the versal family. The k -algebra of observables of the family \mathcal{V} is the k -algebra,

$$O(\mathcal{V}) := \text{End}_H(\tilde{V}) = (H_{ij} \otimes \text{Hom}_k(V_i, V_j))$$

We would like to describe the kernel and the image of the map,

$$\eta : A \rightarrow O(\mathcal{V})$$

To do this we need to consider the matrix Massey products of the form,

$$D_n \rightarrow \text{Ext}_A^2(V_{i_1}, V_{i_n}),$$

the obvious generalizations of the matrix Massey products introduced in [La 2].

Here we shall describe these products using Hochschild cohomology. This is a more convenient way of describing the map η and maybe also an easier way of understanding the nature of the Massey products.

To simplify the notations, put

$$\text{Ext}_A^1(V) = (\text{Ext}_A^1(V_i, V_j))$$

For $l = 2$, the Massey product above is simply the cup product

$$\text{Ext}_A^1(V) \otimes \text{Ext}_A^1(V) \rightarrow \text{Ext}_A^2(V)$$

defined by: Let $(\psi_{ij}^1), (\psi_{ij}^2) \in \text{Ext}_A^1(V)$, and express ψ_{ij}^k as 1-Hochschild cocycles, i.e. $\bar{\psi}_{ij}^1 \in \text{Der}_k(A, \text{Hom}_k(V_i, V_j)), \bar{\psi}_{ij}^2 \in \text{Der}_k(A, \text{Hom}_k(V_i, V_j))$. The cup product $(\psi_{ij}^1) \cup (\psi_{ij}^2) \in \text{Ext}_A^2(V)$, now denoted

$$\langle (\psi_{ij}^1), (\psi_{ij}^2) \rangle \in \text{Ext}_A^2(V)$$

is defined by the 2-cocycle in the Hochschild complex

$$\langle (\psi_{ij}^1), (\psi_{ij}^2) \rangle_{ik}(a, b) = \sum_j \bar{\psi}_{ij}^1(a) \circ \bar{\psi}_{jk}^2(b) \in \text{Hom}_k(V_i, V_k)$$

Suppose $\langle (\psi_{ij}^1), (\psi_{ij}^2) \rangle = 0$, this means that there exists, for each pair (i, k) a 1-cochain ϕ_{ik}^{12} in the Hochschild complex, i.e. a map

$$\phi_{ik}^{12} \in \text{Hom}_k(A, \text{Hom}_k(V_i, V_k))$$

such that $d\phi_{ik}^{12} = \langle (\psi_{ij}^1), (\psi_{ij}^2) \rangle_{ik}$, i.e. such that for all $a, b \in A$,

$$a\phi_{ik}^{12}(b) - \phi_{ik}^{12}(ab) + \phi_{ik}^{12}(a)b = \sum_j \bar{\psi}_{ij}^1(a) \circ \bar{\psi}_{jk}^2(b)$$

Given classes $\psi^1 = (\psi_{ij}^1), \psi^2 = (\psi_{ij}^2), \psi^3 = (\psi_{ij}^3) \in Ext_A^1(V)$ such that $\langle \psi^1, \psi^2 \rangle = \langle \psi^2, \psi^3 \rangle = 0$ there exists $\phi^{12} = (\phi_{ik}^{12}), \phi^{23} = (\phi_{ik}^{23}) \in Hom_k(A, Hom_k(V_i, V_k))$ such that

$$d\phi^{12} = \langle \psi^1, \psi^2 \rangle, \quad d\phi^{23} = \langle \psi^2, \psi^3 \rangle.$$

Then there exists a matrix Massey product

$$\langle \psi^1, \psi^2, \psi^3 \rangle \in Ext_A^2(V)$$

defined by the 2-cocycle

$$\langle \psi^1, \psi^2, \psi^3 \rangle_{ik}(a, b) = \sum_j \phi_{ij}^{12}(a) \psi_{jk}^3(b) - \sum_j \psi_{ij}^1(a) \phi_{jk}^{23}(b)$$

in $Hom_k(A \otimes_k A, Hom_k(V_i, V_j))$.

As in [La 2] we may go on and obtain a sequence of *defining systems* $\{D_n\}_{n=2}^\infty$ and Massey products, computing the relations of $H(\mathcal{V})$.

Now if $a \in A$, denote by $\tilde{a}_i \in Hom_k(V_i, V_i)$ its action on $V_i, i = 1, \dots, d$. Let $End_0(V)$ be the diagonal matrix $(End_k(V_i, V_i))$, contained in the matrix $End_k(V) := (End_k(V_i, V_j))$. Put,

$$End(V)a = (\tilde{a}_1, \dots, \tilde{a}_d) \in End_0(V) \subseteq End(V)$$

If $a \in A$ is such that $End(V)a = 0$, this means that a acts trivially on each V_i . Let $\psi \in Ext_A^1(V)$ be represented by 1-cocycles $\psi_{ij} \in Der_k(A, End_k(V_i, V_j))$. If $End(V)a = End(V)b = 0$, we find,

$$\psi_{ij}(ab) = a\psi_{ij}(b) + \psi_{ij}(a)b = 0$$

This shows that $\psi \in Ext_A^1(V)$ defines a unique k -linear map,

$$\psi : \{a \in A \mid End(V)a = 0\} \rightarrow End_k(V)$$

vanishing on all squares.

Let $a \in A, End(V)a = 0$, and put

$$Ext_A^1(V)a = 0$$

when $\psi(a) = 0, \forall \psi \in Ext_A^1(V)$. Consider the sub k -vector space of A

$$K_1 = \{a \in A \mid End(V)a = Ext_A^1(V)a = 0\}$$

Let $\sum \alpha_{ij} \psi^i \otimes \psi^j \in Ext_A^1(V) \otimes Ext_A^1(V)$ such that its Massey (cup-)product is zero, i.e. such that:

$$\sum \alpha_{ij} \langle \psi^i, \psi^j \rangle = 0$$

Then there exists a 1-cochain $\phi \in Hom_k(A, (Hom_k(V_i, V_j)))$ such that

$$d\phi = \sum_{ij} \alpha_{ij} \langle \psi^i, \psi^j \rangle$$

Since $d\phi = 0$ implies that ϕ represents an element of $Ext_A^1(V)$ it is clear that ϕ defines a unique k -linear map

$$\phi : K_1 \rightarrow End_k(V).$$

Let us denote by

$$ker\langle Ext_A^1(V), Ext_A^1(V) \rangle$$

the subset of $Ext_A^1(V) \otimes Ext_A^1(V)$ for which the Massey product (i.e. the cup product) is zero. Then we may put

$$ker\langle Ext_A^1(V), Ext_A^1(V) \rangle a = 0$$

if for every $d\phi \in ker\langle Ext_A^1(V), Ext_A^1(V) \rangle$, $\phi(a) = 0$.

Let

$$K_2 = \{a \in A \mid End(V)a = Ext_A^1(V)a = ker\langle Ext_A^1(V), Ext_A^1(V) \rangle a = 0\}$$

Continuing in this way we find a sequence of ideals $\{K_n\}_{n \geq 0}$, where $K_0 = ker\{A \rightarrow End(V)\}$ and, in general, $K_n = \{a \in A \mid D_n a = 0\}$.

Theorem 2.8. *Let A be any k -algebra and let $\mathcal{V} = \{V_i\}_{i=1}^r$ be a swarm of A -modules. Then the kernel of the canonical map*

$$\eta : A \rightarrow O(\mathcal{V})$$

is determined by the matrix Massey product structure of $Ext_A^i(V)$, $i = 1, 2$. In fact

$$ker \eta = \bigcap_{n \geq 0} K_n$$

Proof. By definition, the homomorphism of k -algebras

$$\eta : A \rightarrow O(\mathcal{V})$$

lifts the k -algebra homomorphism,

$$\eta_0 : A \rightarrow \prod_{i=1}^d End_A(V_i).$$

Modulo $Rad(H)^2$ η induces the homomorphism,

$$\eta_1 : A \rightarrow \prod_{i=1}^d End_k(V_i) \oplus (E_{ij}^1 \otimes Hom_k(V_i, V_j))$$

with,

$$\eta_1(a)_{ij} = \delta_{ij} \otimes \eta_0(a)_i + \sum_l t_{ij}(l) \otimes \psi_{ij}^1(l)(a_i, -), \quad \delta_{ij} = \begin{cases} 0 & i \neq j \\ 1 & i = j \end{cases}$$

Now, by construction H is the quotient of the formally free k -algebra T^1 generated by the independent variables $\{t_{ij}(l), l = 1, \dots, l_{ij}\}$ as explained above. The relations of T^1 are generated by linear combinations of monomials in these variables of the form,

$$y_{ik} = \sum_{r=1}^{\infty} \sum_{\underline{j}, \underline{l}} \alpha_{i, j_1, \dots, j_{r-1}, k}^{l_1, \dots, l_r} t_{ij_1}(l_1) t_{j_1 j_2}(l_2) \cdots t_{j_{r-1}, k}(l_r),$$

corresponding to elements,

$$y_{ik} \in Ext_A^2(V_i, V_k)^*.$$

The coefficients α are expressed in terms of partially, but inductively well defined, matrix Massey products,

$$\langle \rangle_r : D_r \longrightarrow Ext_A^2(V)$$

such that, if the Massey product $\langle \psi_{ij_1}^1(l_1), \dots, \psi_{j_{r-1}, k}^1(l_r) \rangle$ is defined, then

$$y_{ik}(\langle \psi_{ij_1}^1(l_1), \dots, \psi_{j_{r-1}, k}^1(l_r) \rangle) = \alpha_{ij_1, \dots, j_{r-1}, k}^{l_1, \dots, l_r}.$$

We therefore obtain a basis for H , as k -vector space, by picking, in a coherent way, a k -basis for

$$coker\{Ext_A^2(V)^* \rightarrow D_r^*\} = (\ker \langle \rangle_r)^*$$

Since $K_r = \ker \langle \rangle_r$, the conclusion of the Theorem follows. \square

Remark 2.9. Let E_{ij} be an extension of V_i by V_j , then as a k -vector space $E_{ij} = V_j \oplus V_i$ and the right action by A is defined for $(v_j, v_i) \in E_{ij}$, $a \in A$ by,

$$(v_j, v_i)a = (v_j a + \psi_{ij}^1(a, v_i), v_i a),$$

where,

$$\psi_{ij}^1 \in Der_k(A, Hom_k(V_i, V_j))$$

defines an element,

$$\bar{\psi}_{ij}^1 \in Ext_A^1(V_i, V_j)$$

corresponding to E_{ij} . Suppose we consider an extension E_{ijk} of E_{ij} by V_k . Then as a k -vector space $E_{ijk} \simeq V_k \oplus E_{ij} = V_k \oplus V_j \oplus V_i$ and the action by A is defined by

$$(v_k, v_j, v_i)a = (v_k a + \phi(a, (v_j, v_i)), v_j a + \psi_{ij}^1(a, v_i), v_i a).$$

By additivity

$$\phi(a, (v_j, v_i)) = \phi(a, (v_j, 0)) + \phi(a, (0, v_i)).$$

Put

$$\psi_{ij}^{1,0}(a, v_i) = \psi_{ij}^1(a, v_i), \quad \psi_{jk}^{0,1}(a, v_j) = \phi(a, (v_j, 0)), \quad \psi_{ik}^{1,1}(a, v_i) = \phi(a, (0, v_i)),$$

then the conditions on the action imply

$$\begin{aligned}\psi_{jk}^{0,1} &\in \text{Der}_k(A, \text{Hom}_k(V_j, V_k)) \\ \psi_{jk}^{1,1} &\in \text{Hom}_k(A, \text{Hom}_k(V_i, V_k))\end{aligned}$$

and

$$d\psi_{ik}^{1,1} = \psi_{jk}^{0,1} \circ \psi_{ij}^{1,0}.$$

This means that $\bar{\psi}_{jk}^{0,1} \in \text{Ext}_A^1(V_j, V_k)$ and that the cup product,

$$\bar{\psi}^{0,1} \cup \bar{\psi}^{1,0} \in \text{Ext}_A^2(V_i, V_k)$$

is zero.

Now, consider an extension E_{ijkl} of E_{ijk} by V_l . As before the action of A on E_{ijkl} is given by

$$\begin{aligned}(v_l, v_k, v_j, v_i) \cdot a \\ = (v_l \cdot a + \phi(a, v_k, v_j, v_i), v_k \cdot a + \psi_{ik}^2(a, v_i) + \psi_{jk}^{0,1}(a, v_j), v_j \cdot a + \psi_{ij}^1(a, v_i), v_i \cdot a).\end{aligned}$$

Put, as above,

$$\begin{aligned}\psi^{1,0,0} &= \psi^{1,0} \\ \psi^{0,1,0} &= \psi^{0,1} \\ \psi_{kl}^{0,0,1}(a, v_k) &= \phi(a, v_k, 0, 0) \\ \psi_{jl}^{0,1,1}(a, v_j) &= \phi(a, 0, v_j, 0) \\ \psi_{ik}^{1,1,1}(a, v_i) &= \phi(a, 0, 0, v_i)\end{aligned}$$

The conditions on ϕ are expressed by:

$$\begin{aligned}d\psi_{kl}^{0,0,1} &= 0 \\ d\psi_{jl}^{0,1,1} &= \psi_{kl}^{0,0,1} \circ \psi_{jk}^{0,1,0} \\ d\psi_{il}^{1,1,1} &= \psi_{jl}^{0,1,1} \circ \psi_{ij}^{1,0,0} + \psi_{kl}^{0,0,1} \circ \psi_{ik}^{1,1,0}\end{aligned}$$

This means that $\bar{\psi}_{kl}^{0,0,1} \in \text{Ext}_A^1(V_k, V_l)$, that the cup product $\bar{\psi}_{kl}^{0,0,1} \cup \bar{\psi}_{jk}^{0,1,0} \in \text{Ext}_A^2(V_j, V_l)$ is zero, and that the Massey product

$$\langle \bar{\psi}_{kl}^{0,0,1}, \bar{\psi}_{jk}^{0,1,0}, \bar{\psi}_{ij}^{1,0,0} \rangle \in \text{Ext}_A^2(V_i, V_l)$$

is zero.

It is clear how to continue.

Corollary 2.10. *Suppose the k -algebra A is of finite dimension, and let the finite swarm $\mathcal{V} = \{V_i\}_{i=1}^r$ contain all simple representations, then*

$$\eta : A \rightarrow O(\mathcal{V})$$

is injective.

Proof. Let $a \in A$, and suppose $\eta(a) = 0$. Since A as a right A -module is an extension of the V'_i 's we may assume there are exact sequences of right A -modules

$$0 \longrightarrow Q_1 \longrightarrow A \longrightarrow \bigoplus_{i \in I_1} V_i \longrightarrow 0$$

$$0 \longrightarrow Q_2 \longrightarrow Q_1 \longrightarrow \bigoplus_{i \in I_2} V_i \longrightarrow 0$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

$$0 \longrightarrow Q_N \longrightarrow Q_{N-1} \longrightarrow \bigoplus_{i \in I_N} V_i \longrightarrow 0$$

with $Q_N = \bigoplus_{i \in I_{N+1}} V_i$, $Q_{N+1} = 0$. Since $\text{End}(V)a = 0$ it follows from the first exact sequence above that $1 \cdot a = a \in Q_1$. Consider the exact sequence

$$0 \longrightarrow \bigoplus_{i \in I_2} V_i \longrightarrow A/Q_2 \longrightarrow \bigoplus_{i \in I_1} V_i \longrightarrow 0$$

Since $\text{Ext}_A^1(V)a = 0$ it follows that $1 \cdot a = a \in Q_2$. In fact, multiplication by a is zero on V_i , $i = 1, \dots, r$ and on A/Q_2 it is therefore given by the elements in $\text{Ext}_A^1(V)$. Continuing in this way, we consider the extensions of extensions,

$$0 \longrightarrow \bigoplus_{i \in I_3} V_i \longrightarrow A/Q_3 \longrightarrow A/Q_2 \longrightarrow 0$$

$$\vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots \quad \quad \quad \vdots$$

$$0 \longrightarrow \bigoplus_{i \in I_{N+1}} V_i \longrightarrow A \longrightarrow A/Q_N \longrightarrow 0.$$

Referring to (2.9), we know that the multiplication by $a \in A$ on the right in the middle term is given inductively, by a family of cochains $\psi_{ij}^{\underline{\epsilon}} \in \text{Hom}_k(A, \text{Hom}_k(V_i, V_j))$, with $\underline{\epsilon} \in \{0, 1\}^n$, for $2 \leq n$, such that

$$d\psi_{ik}^{\underline{\epsilon}} - \sum_{\substack{\underline{\epsilon}_1 + \underline{\epsilon}_2 = \underline{\epsilon} \\ j}} \psi_{ij}^{\underline{\epsilon}_1} \circ \psi_{jk}^{\underline{\epsilon}_2}.$$

Now, this means that all these extensions are defined in terms of a series of well defined Massey products each one containing 0. By the proof of Theorem (2.8), we find that for all i, j and all $\underline{\epsilon}$, $\psi_{ij}^{\underline{\epsilon}}(a, -) = 0$.

This means that the action of $a \in A$ must be 0, so $a = 0$. \square

The same proof works for the following,

Corollary 2.10 bis. *Suppose the k -algebra A is an iterated extension of the objects in the finite swarm $\mathcal{V} = \{V_i\}_{i=1}^r$. Then*

$$\eta : A \rightarrow O(\mathcal{V})$$

is injective.

Corollary 2.11. *Suppose A is an object of \underline{a}_r , and let $\mathcal{V} = \{V_i\}_{i=1}^r$ be the family of simple representations, with $V_i \simeq k_i$. Then*

$$A \simeq H$$

Proof. Obviously A is a left A - and a right A -module, flat over A , therefore $A \in \text{Def}_{\mathcal{V}}(A)$. Let $R \in \underline{a}_r$, and pick an element $V_R \in \text{Def}_{\mathcal{V}}(R)$. Since

$$\text{End}(V) = \begin{pmatrix} k & \cdots & k \\ \vdots & & \vdots \\ k & \cdots & k \end{pmatrix},$$

this amounts to a homomorphism of k -algebras $A \rightarrow \text{End}_R(V_R) = R$, implying that A is versal. But then the unicity of the hull of $\text{Def}_{\mathcal{V}}$ gives us an isomorphism:

$$\phi : H \rightarrow A$$

□

Example 2.12. Reconstructing an ordered set Λ and $k[\Lambda]$, from the swarm of simple modules.

Let Λ be an ordered set, see §1, and let $A = k[\Lambda]$, $V = \{k_\lambda\}_{\lambda \in \Lambda}$. Then the Corollary above implies that $H \simeq k[\Lambda]$.

1. By the general theory we know that $A = k[\Lambda]$ is the matrix algebra generated freely by the immediate relations $\lambda_1 \gg \lambda_2$, i.e. those for which $\{\lambda' \in \Lambda \mid \lambda_1 > \lambda' > \lambda_2\} = \emptyset$, modulo relations of the form


$$\begin{aligned} & (\lambda' > \lambda_2^1)(\lambda_2^1 > \lambda_3^1) \cdots (\lambda_{n_1}^1 > \lambda) \\ & = (\lambda' > \lambda_2^2)(\lambda_2^2 > \lambda_3^2) \cdots (\lambda_{n_2}^2 > \lambda) \end{aligned}$$

They correspond to the first obstructions, given by the n_i term well defined Massey products

$$\begin{aligned} & \text{Ext}_A^1(k_{\lambda'}, k_{\lambda_2^1}) \otimes \cdots \otimes \text{Ext}_A^1(k_{\lambda_{n_1}^1}, k_\lambda) \rightarrow \text{Ext}_A^2(k_{\lambda'}, k_\lambda) \\ & \text{Ext}_A^1(k_{\lambda'}, k_{\lambda_2^2}) \otimes \cdots \otimes \text{Ext}_A^1(k_{\lambda_{n_2}^2}, k_\lambda) \rightarrow \text{Ext}_A^2(k_{\lambda'}, k_\lambda) \end{aligned}$$

There are as many relations as there are base elements of $\text{Ext}_A^2(k_{\lambda'}, k_\lambda)$.

2. Let us check this for the diamond, i.e. for Λ :



One easily computes the Ext 's,

$$Ext_A^1(k_{\lambda_i}, k_{\lambda_j}) = \begin{cases} 0 & i = j \\ k & \text{for } i = 1, j = 2, 3 \\ k & \text{for } i = 2, 3, j = 4 \end{cases}$$

$$Ext_A^2(k_{\lambda_i}, k_{\lambda_j}) = \begin{cases} 0 & \text{for } (i, j) \neq (1, 4) \\ k & \text{for } i = 1, j = 4 \end{cases}$$

The two cup-products

$$Ext_A^1(k_{\lambda_1}, k_{\lambda_j}) \otimes Ext_A^1(k_{\lambda_j}, k_{\lambda_4}) \rightarrow Ext_A^2(k_{\lambda_1}, k_{\lambda_4}) \quad \text{for } j = 2, 3,$$

are non-trivial. At the tangent level we have:

$$H_2 = \begin{pmatrix} k & k & k & 0 \\ 0 & k & 0 & k \\ 0 & 0 & k & k \\ 0 & 0 & 0 & k \end{pmatrix}$$

Therefore H must be a quotient of the matrix ring,

$$T^1 = \begin{pmatrix} k & t_{12} \cdot k & t_{13} \cdot k & (t_{12}t_{24} \cdot k + t_{13}t_{34} \cdot k) \\ 0 & k & 0 & t_{24} \cdot k \\ 0 & 0 & k & t_{34} \cdot k \\ 0 & 0 & 0 & k \end{pmatrix}$$

The kernel of $T^1 \rightarrow H$ is given in terms of the cup products above. In fact, since we have $t_{13}^* \cup t_{34}^* = t_{12}^* \cup t_{24}^* = y^*$ where y^* is the generator of $Ext_A^2(k_{\lambda_1}, k_{\lambda_4})$, the kernel of $T^1 \rightarrow H$ is simply $t_{13} \otimes t_{34} + t_{12} \otimes t_{24}$ such that

$$H = \begin{pmatrix} k & k & k & k \\ 0 & k & 0 & k \\ 0 & 0 & k & k \\ 0 & 0 & 0 & k \end{pmatrix} \simeq k[\Lambda]$$

as it should.

In general, we may reconstruct Λ from the tangent space T_H and the Massey-products above.

The corresponding problem for finite groups, i.e. reconstructing G from $k[G]$ is called the isomorphism problem. Due to some nice examples of Dade, we know that this is hopeless. In fact there are two non isomorphic finite groups such that their group-algebras are isomorphic for all fields.

Example 2.13. Given any scheme $\underline{H} = Spec(H)$, say the 2-dimensional affine space given by $H = k[x_1, x_2]$. We shall be interested in the noncommutative moduli space parametrizing subschemes of length 2 of \underline{H} . We may do this by simply considering a point in the space $Spec(H)$ together with a tangent direction, i.e. the right H -module of the form,

$$V = k[x_1, x_2]/(x_1^2, x_2),$$

and compute the formal moduli of V .

Lemma 2.14. *The formal moduli, $H(V)$ of the H -module $V = H/(x_1^2, x_2)$, is given as the completion of the k -algebra,*

$$\Omega = k\{t_1, t_2, \omega_1, \omega_2\}/(y_1, y_2)$$

where

$$y_1 = [t_1, t_2] - t_1[\omega_1, \omega_2] \quad y_2 = [t_1, \omega_2] - [t_2, \omega_1] - \omega_1[\omega_1, \omega_2],$$

and where the family of left Ω -and right H -modules,

$$\Omega \otimes_k k^2$$

is defined by the actions of x_1 and x_2 , given by,

$$x_1 = \begin{pmatrix} 0 & t_1 \\ 1 & \omega_1 \end{pmatrix}, \quad x_2 = \begin{pmatrix} t_2 & t_1\omega_2 \\ \omega_2 & t_2 + \omega_1\omega_2 \end{pmatrix}$$

Proof. Consider the obvious free resolution of $V := H/(x_1^2, x_2)$ as an H -module,

$$V \xleftarrow{\rho} H \xleftarrow{d_0} H^2 \xleftarrow{d_1} H \xleftarrow{d_2} 0$$

where we have,

$$d_0 = (x_1^2, x_2), \quad d_1 = \begin{pmatrix} x_2 \\ -x_1^2 \end{pmatrix}.$$

Consider the Yoneda complex, and pick a basis

$$\{\hat{t}_1, \hat{t}_2; \hat{\omega}_1, \hat{\omega}_2, \}$$

of $Ext_H^1(V, V)$ represented by the morphisms of the diagram,

$$\begin{array}{ccccccc} V & \xleftarrow{\rho} & H & \xleftarrow{d_0} & H^2 & \xleftarrow{d_1} & H \xleftarrow{\quad} 0 \\ & & & \searrow^{\hat{\omega}_j} & & \searrow^{\hat{\omega}_j^2} & \\ V & \xleftarrow{\rho} & H & \xleftarrow{d_0} & H^2 & \xleftarrow{d_1} & H \xleftarrow{\quad} 0 \\ & & & \searrow^{\hat{\omega}_j} & & \searrow^{\hat{\omega}_j^2} & \\ V & \xleftarrow{\rho} & H & \xleftarrow{d_0} & H^2 & \xleftarrow{d_1} & H \xleftarrow{\quad} 0 \\ & & & \searrow^{\hat{t}_i} & & \searrow^{\hat{t}_i^2} & \end{array}$$

Here,

$$\begin{aligned} \hat{t}_1^1 &= (1, 0), \quad \hat{t}_2^1 = (0, 1,); \\ \hat{\omega}_1^1 &= (x_1, 0), \quad \hat{\omega}_2^1 = (0, x_1) \end{aligned}$$

and,

$$\hat{t}_1^2 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}, \quad \hat{t}_2^2 = \begin{pmatrix} -1 \\ 0 \end{pmatrix},$$

and finally,

$$\hat{\omega}_1^2 = \begin{pmatrix} 0 \\ x_1 \end{pmatrix}, \quad \hat{\omega}_2^2 = \begin{pmatrix} -x_1 \\ 0 \end{pmatrix}.$$

Using this it is easy to see that ,

$$\hat{t}_i \cup \hat{t}_i = 0, \quad \hat{t}_1 \cup \hat{t}_2 = -\hat{t}_2 \cup \hat{t}_1 = \hat{y}_1,$$

and that

$$\hat{t}_1^1 \hat{\omega}_2^2 = \hat{\omega}_1^1 \hat{t}_2^2 = -\hat{y}_2, \quad \hat{\omega}_i^1 \hat{t}_i^2 = 0, \quad \hat{\omega}_i^1 \hat{\omega}_j^2 = 0, \quad \hat{t}_2^1 \hat{\omega}_1^2 = \hat{\omega}_2^1 \hat{t}_1^2 = \hat{y}_2,$$

where,

$$\{\hat{y}_1, \hat{y}_2\}$$

is a basis for of $Ext_H^2(V, V)$ represented by the morphisms of the diagram,

$$\begin{array}{ccccccc} V & \xleftarrow{\rho} & H & \xleftarrow{d_0} & H^2 & \xleftarrow{d_1} & H \xleftarrow{\quad} 0 \\ & & & & & \nearrow \hat{y}_1 & \\ & & & & & \hat{y}_2 & \\ V & \xleftarrow{\rho} & H & \xleftarrow{d_0} & H^2 & \xleftarrow{d_1} & H \xleftarrow{\quad} 0 \end{array}$$

where $\hat{y}_1 = (1)$ and $\hat{y}_2 = (x_1)$. Therefore

$$-\hat{y}_2 = \hat{t}_1 \cup \hat{\omega}_2 = \hat{\omega}_1 \cup \hat{t}_2 = -\hat{t}_2 \cup \hat{\omega}_1 = -\hat{\omega}_2 \cup \hat{t}_1, \quad \hat{\omega}_i \cup \hat{\omega}_j = \hat{t}_i \cup \hat{t}_j = 0.$$

Now, consider the dual basis $\{t_1, t_2; \omega_1, \omega_2\}$ generating the hull of the deformation functor $Def_{k[\epsilon]}$, we find after a simple computation of the 3. order Massey products the formulas we want.

Notice that we just have to compute the tangent situation and check that our formulas give us a lifting of the quadratic relations and of the corresponding H -action, to know that our result holds.

□

By a simple computation one checks that the k -points of Ω form an open dense part of $Hilb^2 \mathbf{A}^2$ containing V . $Hilb^2 \mathbf{A}^2$ is the blow-up of $(\mathbf{A}^2 \times \mathbf{A}^2)/\mathbf{Z}_2$ along the diagonal. However, there are other simple representations of Ω . The homomorphism,

$$\Omega \rightarrow k[t_1, t_2, \frac{\partial}{\partial t_1}, \frac{\partial}{\partial t_2}]$$

mapping ω_i to $\frac{\partial}{\partial t_i}$, shows that $k[t_1, t_2]$ is a simple representation of Ω .

§3. Noncommutative modular deformations. Let V be any right A -module such that $\dim_k Ext_A^1(V, V) < \infty$. Consider the formal moduli $H^A =: H$, the formal versal family $\tilde{V} = H \otimes V$, and the corresponding morphism of functors,

$$\rho : Mor_{\underline{a}_r}(H, -) \rightarrow Def_V.$$

We know that ρ is not, in general, injective. However, V is also a right $A \otimes \text{End}_A(V)$ -module. As such it has a formal moduli $H^{A, \text{End}}$, and there is a natural k -algebra homomorphism, $H^A \rightarrow H^{A, \text{End}}$. Let H_0^A be the unique maximal common quotient of H^A and $H^{A, \text{End}}$. Using the same construction as in [La, Pf], §2, we prove that the composition,

$$\rho_0 : \text{Mor}_{\underline{a}_r}(H_0, -) \rightarrow \text{Mor}_{\underline{a}_r}(H, -) \rightarrow \text{Def}_V$$

is injective.

At the tangent level, the homomorphisms,

$$H^A \rightarrow H^{A, \text{End}} \leftarrow H^{\text{End}},$$

looks like the canonical homomorphisms,

$$\text{Ext}_A^1(V, V) \leftarrow \text{Ext}_{A \otimes_k \text{End}}^1(V, V) \rightarrow \text{Ext}_{\text{End}}^1(V, V).$$

Representing elements of the Ext-groups as derivations, it is easy to see that the two images are contained in the subspace $\text{Ext}_A^1(V, V)^{\text{End}}$, respectively $\text{Ext}_{\text{End}}^1(V, V)^A$. Therefore the tangent space of H_0 must be contained in the subspace of invariants under $\text{End}_A(V)$ of the tangent space of H , $\text{Ext}_A^1(V, V)^{\text{End}}$.

The tangent space of the modular (prorepresenting) substratum, and almost split sequences.

Consider as above a swarm $\mathcal{V} = \{V_i\}_{i=1}^r$ of A -modules, and consider the k^r -algebra

$$\text{End}_A(V) = (\text{Hom}_A(V_i, V_j)).$$

Suppose from now on that the modules V_i are non-isomorphic, indecomposables, and that for each $i = 1, \dots, r$, $\text{End}_A(V_i)$ is a commutative local ring with maximal ideal \underline{m}_i .

Lemma 3.1. *Under the above assumptions, the radical of $\text{End}_A(V)$ has the form*

$$\text{rad}(V) = \begin{pmatrix} \underline{m}_1 \text{End}_A(V_1) & & & \vdots & & \\ & \cdots & \cdots & \underline{m}_i \text{End}_A(V_i) & \text{Hom}_A(V_i, V_j) & \cdots \\ & & & & \underline{m}_j \text{End}_A(V_j) & \\ & & & \vdots & & \underline{m}_r \text{End}_A(V_r) \end{pmatrix}$$

Proof. We need only check that $\text{rad}(V)$ is an ideal, and this amounts to proving that if $\phi_{ij} \in \text{Hom}(V_i, V_j)$ $i \neq j$ and $\phi_{ji} \in \text{Hom}(V_j, V_i)$ then

$$\phi_{ji}\phi_{ij} \in \underline{m}_i \subseteq \text{End}_A(V_i).$$

Suppose $\phi_{ji}\phi_{ij}$ is not in \underline{m}_i , then $\phi_{ji}\phi_{ij}$ is an isomorphism, and we may as well assume that $\phi_{ji}\phi_{ij} = \text{id}_{V_i}$. But then $V_j \simeq V_i \oplus \ker \phi_{ji}$ which contradicts the indecomposability of V_j . \square

In particular this lemma proves that if A is artinian and all V_i are of finite type, then for some N ,

$$\text{rad}(V)^N = 0$$

Obviously there is a left and a right action of $End_A(V)$ on

$$T_H = (Ext_A^1(V_i, V_j)).$$

The difference between these actions defines the action of the Lie algebra $End_A(V)$ on T_H . The invariants of T_H under the Lie algebra $rad(V)$, is equal to the invariants under $End_A(V)$, therefore equal to,

$$T_{H_0} := \{\xi \in T_H | \forall \phi \in End_A(V), \quad \phi\xi - \xi\phi = 0\},$$

containing the tangent space of the *modular*, or the *prorepresentable* substratum H_0 of H .

Lemma 3.2. *Let $\xi \in T_{H_0}$, with $\xi = (\xi_{i,j})$, then for all $\phi = (\phi_{k,l}) \in rad(V)$ we have for $i \neq j$, and all l ,*

$$\begin{aligned} \phi_{l,i}\xi_{i,j} &= 0 \\ \xi_{i,j}\phi_{j,l} &= 0. \end{aligned}$$

Moreover, for all i, j

$$\phi_{i,j}\xi_{j,j} = \xi_{i,i}\phi_{i,j}$$

Proof. Just computation. \square

Definition 3.3. *In the above situation, an extension $\xi \in Ext_A^1(V_i, V_j)$ is called a left almost split extension (resp. a right almost split extension), lase (resp. rase) for short, if for all $\phi_{ki} \in r(V)_{ki}$ (resp. $\phi_{jk} \in r(V)_{jk}$)*

$$\phi_{ki}\xi = 0 \quad (\text{resp.} \quad \xi\phi_{jk} = 0).$$

An extension ξ which is both a lase and a rase is called an *ase*, an *almost split extension*.

This, of course, is nothing but a trivial generalization of the notion of almost split sequence, due to Auslander, see [R].

Denote by $Ext_l^1(V_i, V_j)$ (resp. $Ext_r^1(V_i, V_j)$) the subspace of $Ext_A^1(V_i, V_j)$ formed by the lase's (resp. rase's), and put

$$\begin{aligned} T_H^l &= (Ext_l^1(V_i, V_j)) \subseteq T_H \\ T_H^r &= (Ext_r^1(V_i, V_j)) \subseteq T_H \\ T_H^a &= T_H^l \cap T_H^r =: (Ext_a^1(V_i, V_j)) \subseteq T_H. \end{aligned}$$

Observe that since the left and the right action of $End(V)$ on T_H commute, $End(V)$ acts at right on T_H^l and at left on T_H^r . Moreover, by the lemma above

$$T_H^a = T_H^l \cap T_H^r \subseteq T_{H_0}.$$

Observe also that if $End_A(V_i) = k \oplus \underline{m}_i$ the diagonal part of T_{H_0} is exactly the tangent space of the deformation functor of the full subcategory of mod_A generated by V , see [La 1].

The structure of the modular substratum, and the existence of almost split sequences for artinian k -algebras.

Assume that A is artinian, and that the V_i 's are of finite type. Then T_H is a k -vectorspace of finite dimension, and the radical $rad(V)$ of $End(V)$ acts nilpotently on T_H .

Corollary 3.4. *Given $i \in \{1, \dots, r\}$, assume there exists one $j \in \{1, \dots, r\}$ such that $Ext_A^1(V_i, V_j) \neq 0$. Then there exists a $\tau(i) \in \{1, \dots, r\}$ such that,*

$$Ext_r^1(V_i, V_{\tau(i)}) \neq 0.$$

Proof. This is simply Engels theorem for the right action of $rad(V)$ on T_H . \square

Theorem 3.5. *Suppose \mathcal{V} is such that every extension $\xi \in Ext_A^1(V_i, V_j)$ is of the form $0 \rightarrow V_j \rightarrow E \rightarrow V_i \rightarrow 0$ with E a direct sum of V_k 's. Then, for every $i = 1, \dots, r$, such that there exists a $j = 1, \dots, r$ for which $Ext_A^1(V_i, V_j) \neq 0$, there is a unique ase of the form*

$$0 \rightarrow V_{\tau(i)} \rightarrow E_i \rightarrow V_i \rightarrow 0$$

Moreover, if we agree to put $\tau(i) = i$ for those i 's for which $Ext_A^1(V_i, V_k) = 0$ for all k , then

$$\tau : \{1, \dots, r\} \rightarrow \{1, \dots, r\}$$

is a permutation.

Proof. We already know that there exists a rase of the form

$$\xi_i : 0 \rightarrow V_{\tau(i)} \rightarrow E_i \rightarrow V_i = 0.$$


We shall prove that ξ_i is also a lase. Let $\phi_{ki} \in Hom_A(V_k, V_i)$ for $k \neq i$, or pick an element $\phi_{ii} \in \underline{m}_i \subseteq End(V_i)$, and consider the commutative diagram,

$$\begin{array}{ccccccc}
 & & 0 & & & & \\
 & & \downarrow & & & & \\
 0 & \longrightarrow & V_{\tau(i)} & \longrightarrow & E_i & \longrightarrow & V_i \longrightarrow 0, & \xi_i \\
 & & \downarrow & \nearrow \psi_{ki} & & & \nearrow & \\
 & & E_k = V_k \times_{V_i} E_i & & & & & \\
 & & \downarrow & \nearrow \phi_{ki} & & & & \\
 & & V_k & & & & & \\
 & & \downarrow & & & & & \\
 & & 0 & & & & &
 \end{array}$$

Suppose $V_{\tau(i)} \rightarrow E_k$ is not split, then

$$0 \rightarrow E_k \rightarrow E_k \oplus^{V_{\tau(i)}} E_i \rightarrow V_i \rightarrow 0$$


is split, since ξ_i is a rase. Let $\text{pr.}: E_k \oplus^{V_{\tau(i)}} E_i \rightarrow E_k$ be the splitting. But then the two following diagrams commute:



Here ψ_{ik} is the composition of $E_i \rightarrow E_k \oplus^{V_{\tau(i)}} E_i$ and the projection $E_k \oplus^{V_{\tau(i)}} E_i \rightarrow E_k$ and ϕ_{ik} the induced map.

This means that $(\phi_{ki}\phi_{ik})\xi_i = \xi_i$ which is impossible since $(\phi_{ki}\phi_{ik})$ acts nilpotently on $\text{Ext}_A^1(V_i, V_{\tau(i)})$, and ξ is nonzero. Therefore $V_{\tau(i)} \rightarrow E_k$ splits and ξ_i is also a lase, therefore an ase.

The unicity and the permutation property follows immediately from the following: Assume there exist two ase's ξ_i and ξ'_i of the form:



Then, since ρ'_i is not split, there exist liftings ψ_i, ψ'_i inducing morphisms ϕ_i, ϕ'_i . But then $(\phi_i\phi'_i)\xi_i = \xi_i$ which means that ξ_i is zero. Therefore an ase is unique and in particular, $\tau(i) = \tau(i')$. Dually we prove that $\tau(i) = \tau(i')$ implies $i = i'$, so that τ is a permutation.

We see that T_V^a looks like:

$$\left(\text{Ext}_a^1(V_i, V_j) \right)$$

where

$$\text{Ext}_a^1(V_i, V_j) = \begin{cases} 0 & \text{if } j \neq \tau(i) \\ k & \text{if } j = \tau(i) \text{ and some } \text{Ext}_a^1(V_i, V_j) \neq 0 \end{cases}$$

□

Corollary 3.6. *With the assumptions of the theorem above, we find that*

$$T_{H_0} = \{(\alpha_{ij}) \mid \left. \begin{cases} \alpha_{ij} \in k, \alpha_{ij} = 0 \text{ if } j \neq \tau(i) \\ \alpha_{ii} \in \text{Ext}_A^1(V_i, V_i) \forall \phi_{ji} \in \text{End}_A(V_i), \phi_{ij}\alpha_{ii} = \alpha_{jj}\phi_{ji}, \text{ if } i=j \end{cases} \right\}$$

Remark 3.7. Consider again a not necessarily finite swarm $\mathcal{V} = \{V_i\}_{i=1}^{\aleph}$ of noetherian A -modules. The k^r -algebra

$$\text{End}_A(V)_r := (\text{Hom}_A(V_i, V_j)), \quad i, j \leq r$$

acts on

$$(\text{Ext}_A^1(V_i, V_j)), \quad i, j \leq r,$$

in the way described above. Suppose that the modules V_i are non-isomorphic, indecomposables, and that for each i , $\text{End}_A(V_i)$ is a local ring with maximal ideal \underline{m}_i . Suppose moreover that any iterated extension is a direct sum of such V_i 's. This is obviously the case when $\mathcal{V} = \{V_i\}_{i=1}^{\aleph}$ is the family of all indecomposable A -modules, but holds in many other interesting cases, see [R].

Let $H := H(\mathcal{V})$ and \tilde{V} be the prorepresentable hull and the formal versal family, as defined in §2. For every quotient R of H in \underline{a}_{\aleph} , such that

$$\dim_k \text{Rad}(R)/\text{Rad}(R)^2 < \infty$$

we consider the image $\tilde{V}(R) \in \text{Def}_{\mathcal{V}}(R)$ of \tilde{V} . Denote by $L_i(R)$ the i^{th} -line of $\tilde{V}(R)$. $L_i(R)$ is an A -module and a finite iterated extension of the V_j , therefore a finite sum of indecomposables $\{L_i(R, p)\}$, from our family. Obviously there is a canonical surjection,

$$L_i(R) \rightarrow V_i,$$

and a homomorphism of k -algebras,

$$\iota : H \rightarrow \text{End}_A(\tilde{V}(R)),$$

defined by left multiplication. Any element,

$$r_{i,j} \in R_{i,j} \subset R$$

defines a homomorphism of right A -modules,

$$r_{i,j_*} : L_j(R) \rightarrow L_i(R).$$

In particular, if $r_{i,j}$ is in the socle of R , this morphism induces a homomorphism of right A -modules,

$$r_{i,j_*} : V_j \rightarrow L_i(R).$$

Using this, we may consider different quiver-structures on the set of indecomposable modules, $\{V_i\}_{i=1}^n$. The Auslander-Reiten quiver, see [R], is obtained by picking $R = H_0/Rad(H_0)^2$, a basis $\{h_{i,j}\}$ of $Rad(H_0)/Rad(H_0)^2$, the dual tangent space of H_0 and letting the arrows arriving at an indecomposable V_i be the compositions,

$$L_i(R, p) \rightarrow L_i(R) \rightarrow V_i,$$

and the arrows leaving an indecomposable V_j be the compositions,

$$V_j \xrightarrow{h_{i,j}} L_i(R) \rightarrow L_i(R, p).$$

For an arbitrary quotient R of H , we may construct another quiver containing more information than the Auslander-Reiten quiver. Consider representatives

$$\{r_{i,j}\} \in Rad(R)$$

of a basis of the dual tangent space $Rad(R)/Rad(R)^2$ of R , and let the arrows of the quiver be the compositions

$$L_j(R, q) \rightarrow L_j(R) \xrightarrow{r_{i,j}^*} L_i(R) \rightarrow L_j(R, p).$$

There is a ring homomorphism,

$$R \rightarrow End_A(\tilde{V}) = (Hom_A(L_j, L_i)) = (Hom_A(V_q, V_p)^{n_{p,q}}).$$

If this homomorphism is surjective, or an isomorphism, we find that the arrows of the quiver generate, in an obvious way, all morphisms of the full sub-category of A -modules defined by the family of indecomposables $\mathcal{V} = \{V_i\}$. In both case, it is easy to see that the relations in the quiver correspond to non-trivial cup and Massey products of $Ext_a^*(V_i, V_j)$. When A is artinian, and the family $\{V_i\}$ generates the category of A -modules, it turns out that \tilde{V} is a projective generator. H defines a quiver, with vertices corresponding to the indecomposable projectives, and

$$H = End_A(\tilde{V})$$

is Morita equivalent to A . Moreover, H is determined by the quiver (with relations). In particular, if $\{V_i\}$ is the family of simple A -modules, we shall see in the next paragraph that,

$$A \rightarrow (H_{i,j} \otimes Hom_k(V_i, V_j))$$

is an isomorphism, and that

$$L_i = \bigoplus_{j=1,2,\dots} H_{i,j} \otimes V_j$$

is a projective A -module, for $i = 1, 2, \dots, r$. Since therefore

$$H \simeq End_A(\tilde{V}) = (Hom_A(L_j, L_i))$$

is Morita-equivalent to A , the quiver of (projective) summands of \tilde{V} determines the Morita-equivalence class of A .

We shall end this paragraph by proving the following easy result, see [K] for the notions of Frobenius extension and Frobenius bi-module.

Proposition 3.8. *Suppose the following conditions hold:*

(i) *The family $\mathcal{V} = \{V_i\}$ of right A -modules are either finite dimensional as k -vector spaces, or such that,*

$$\text{Ext}_A^p(V_i, V_j) = \text{Ext}_A^p(V_j^*, V_i^*).$$

(ii) *The hull of the noncommutative deformation functor, $H(\mathcal{V}) = (H_{i,j})$ is a finite dimensional k -vectorspace.*

(iii) *For each i , the projective cover of V_i has a (finite) filtration with graded components contained in the family \mathcal{V} .*

Then

$$\eta : A \rightarrow O(\mathcal{V})$$

is a Frobenius extension.

Proof. The assumption 1. implies that the versal family $P = \tilde{V}$ as a left H and right A -module has the duality property, ${}^*P = P^*$. The assumption 2. implies that P as left H -module is finite projective, and the assumption 3. guarantees that P , as right A -module, is finite projective, therefore a Frobenius bi-module. \square

§4. The generalized Burnside theorem.

In §2 we proved the following result,

Corollary 2.10. *Suppose the k -algebra A is of finite dimension and assume the swarm $\mathcal{V} = \{V_i\}_{i=1}^r$ contains all simple A -modules, then the natural k -algebra homomorphism*

$$\eta : A \rightarrow O(\mathcal{V}) = (H_{ij} \otimes_k \text{Hom}_k(V_i, V_j))$$

is injective.

Recall also the classical Burnside-Wedderburn-Malcev theorems, see [Lang], and [Curtis and Reiner].

Theorem (Burnside). *Let V be a finite dimensional k -vectorspace. Assume k is algebraically closed and let A be a subalgebra of $\text{End}_k(V)$. If V is a simple A -module, then $A = \text{End}_k(V)$.*

Theorem (Wedderburn). *Let A be a ring, and let V be a simple faithful A -module. Put $D = \text{End}_A(V)$ and assume V is a finite dimensional D -vector space. Then $A \simeq \text{End}_D(V)$.*

Theorem (Wedderburn-Malcev). *Let A be a finite dimensional k -algebra, k -any field. Let \mathfrak{r} be the radical of A , and suppose the residue class algebra A/\mathfrak{r} is separable. Then there exists a semi-simple subalgebra S of A such that A is the semidirect sum of S and \mathfrak{r} . If S_1 and S_2 are subalgebras such that $A = S_i \oplus \mathfrak{r}$, $i = 1, 2$, then there exists an element $n \in \mathfrak{r}$, such that $S_1 = (1 - n) \cdot S_2 \cdot (1 - n)^{-1}$.*

In this § we shall prove a generalization of the theorem of Burnside. In fact, assuming the field k is algebraically closed and that $\mathcal{V} = \{V_i\}_{i=1}^r$ is the family of all simple A -modules we shall prove that the homomorphism η of the above Corollary (2.10), is an isomorphism.

When A is semi-simple we know that $\text{Ext}_A^1(V_i, V_j) = 0$ for all $i, j = 1, \dots, r$, therefore the formal moduli H of V is isomorphic to k^r . This implies that

$$\text{End}_H(\tilde{V}) = \bigoplus_{i=1}^r \text{End}_k(V_i),$$

which is the classical extension of Burnside's theorem.

We shall need the following elementary lemma

Lemma 4.1. *Let the k -algebra A be a direct sum of the right- A -modules V_i , $i = 1, \dots, d$ of the family $\mathcal{V} = \{V_i\}_{i=1}^r$. Then left multiplication with an element $a \in A$ induces A -module homomorphisms*

$$a_{ij} \in \text{Hom}_A(V_i, V_j), \quad i, j = 1, \dots, d.$$

Moreover, any k -linear map $x : A \rightarrow A$ expressed as $x = (x_{ij}) \in \text{End}_k(V) := (\text{Hom}_k(V_i, V_j))$, commuting with all $\varphi = (\varphi_{ij}) \in \text{End}_A(V) := (\text{Hom}_A(V_i, V_j))$ is necessarily a right multiplication by some element $\tilde{x} \in A$.

Proof. Trivial, since x commuting with all $\varphi \in (\text{Hom}_A(V_i, V_j))$ commutes with all left-multiplications by $a \in A$, and therefore $x(a) = a \cdot x(1)$, and we may put $\tilde{x} = x(1)$. \square

Corollary 4.2. *Assume that the family of right A -modules $\mathcal{V} = \{V_i\}_{i=1}^r$ is such that*

$$(i) \quad A \simeq \bigoplus_{i=1}^m V_i^{n_i}$$

$$(ii) \quad \text{Hom}_A(V_i, V_j) = 0 \text{ for } i \neq j$$

Then the canonical morphism of k -algebras

$$\eta : A \rightarrow \bigoplus_{i=1}^{n_i} \text{End}_k(V_i)$$

is injective. Moreover, η induces an isomorphism

$$A \simeq \bigoplus_{i=1}^{n_i} \text{End}_{D_i}(V_i)$$

where $D_i = \text{End}_A(V_i)$.

This, in particular, implies the Wedderburn theorem for semisimple k -algebras A .

Theorem 4.3 (A generalized Burnside theorem). *Let A be a finite dimensional k -algebra, k an algebraically closed field. Consider the family $\mathcal{V} = \{V_i\}_{i=1}^r$ of simple A -modules, then*

$$A \simeq O(\mathcal{V}) = (H_{i,j} \otimes \text{Hom}_k(V_i, V_j))$$

Proof. We know that the canonical map

$$\eta : A \rightarrow O(\mathcal{V})$$

is injective. Since $\text{Rad}(A)^n = 0$ for some n , we know that $\hat{A} = A$. The theorem therefore follows from the following lemmas.

Lemma 4.4. *Let A and B be finite type k -algebras and let $\varphi : A \rightarrow B$ be a homomorphism of k -algebras such that the induced morphism*

$$\varphi_2 : A \rightarrow B/\text{Rad}(B)^2$$

is surjective, then

$$\hat{\varphi} : \hat{A} \rightarrow \hat{B}$$

is surjective.

Proof. Well-known. \square

Lemma 4.5. *Let A be a finite dimensional k -algebra, k an algebraically closed field. Let $\mathcal{V} = \{V_i\}_{i=1}^r$ be the family of simple A -modules. Then the homomorphism*

$$\eta : A \rightarrow O(\mathcal{V})$$

induces an isomorphism

$$\text{Rad}(A)/\text{Rad}(A)^2 \simeq (\text{Ext}_A^1(V_i, V_j)^* \otimes_k \text{Hom}_k(V_i, V_j)).$$

Proof. The classical Burnside theorem implies that the canonical homomorphism of k -algebras

$$A \twoheadrightarrow \bigoplus_{i=1}^r \text{End}_k(V_i)$$

induces an isomorphism,

$$A/\text{Rad}(A) \simeq \bigoplus_{i=1}^r \text{End}_k(V_i).$$

According to the Wedderburn-Malcev theorem we may assume that $A/\text{Rad}(A)^2$ is a semidirect sum,

$$A/\text{Rad}(A) \oplus \text{Rad}(A)/\text{Rad}(A)^2.$$

Since $Rad(A)/Rad(A)^2$ is both a left and a right $\bigoplus_{i=1}^r End_k(V_i)$ -module

$$Rad(A)/Rad(A)^2 = (E_{ij})$$

each E_{ij} being an $End_k(V_i)^{op} \otimes_k End_k(V_j)$ -module. This, however, means that

$$E_{ij} \simeq Hom_k(V_i, V_j) \otimes k^{r_{ij}}$$

as a right $End_k(V_i)^{op} \otimes_k End_k(V_j)$ -module. Since we already know that η is an injection, we must have,

$$E_{ij} \simeq Hom_k(V_i, V_j) \otimes k^{r_{ij}} \subset Ext_A^1(V_i, V_j) \otimes Hom_k(V_i, V_j).$$

We must show that this inclusion is an equality. Applying Hochschild cohomology as in §1, we find:

$$Ext_A^1(V_i, V_j) = HH^1(A, Hom_k(V_i, V_j)) = Der_k(A, Hom_k(V_i, V_j))/im d^\circ$$

where d° is the differential

$$Hom_k(V_i, V_j) \rightarrow Der_k(A, Hom_k(V_i, V_j)).$$

Clearly any derivation

$$\xi \in Der_k(A, Hom_k(V_i, V_j))$$

which is zero on $Rad(A)$ induces a derivation

$$\xi_0 \in Der_k(A/Rad(A), Hom_k(V_i, V_j))$$

which, since $A/Rad(A)$ is semisimple, obviously is a coboundary, i.e. an element of $im d^\circ$.

Moreover, any derivation $\xi \in Der_k(A, Hom_k(V_i, V_j))$ induces the zero map on $Rad(A)^2$ since $\xi(r_1 \cdot r_2) = r_1 \xi(r_2) + \xi(r_1) r_2 = 0$ for $r_1, r_2 \in Rad(A)$, and any coboundary $\nu \in im d^\circ$ must vanish on $Rad(A)$ since $\nu(r) = \varphi r - r \varphi$, for some $\varphi \in Hom_k(V_i, V_j)$. Now, every $A^{op} \otimes_k A$ -linear map $Rad(A)/Rad(A)^2 \rightarrow End_k(V_i, V_j)$ extends to a derivation of $Der_k(A/Rad(A)^2, Hom_k(V_i, V_j))$. In fact, let φ be an $A^{op} \otimes_k A$ -linear map

$$Rad(A)/Rad(A)^2 \rightarrow End_k(V_i, V_j)$$

and define the map

$$\psi : A/Rad(A)^2 = A/Rad(A) \bigoplus Rad(A)/Rad(A)^2 \rightarrow End_k(V_i, V_j)$$

by

$$\psi(s, r) = \varphi(r) + \varphi(\rho(s))$$

where ρ is the 1-Hochschild cochain on $A/\text{Rad}(A)$ with values in $\text{Rad}(A)/\text{Rad}(A)^2$ that, according to the Wedderburn-Malcev theorem, defines the semidirect sum referred to above. Then,

$$\begin{aligned}\psi((s_1, r_1) \cdot (s_2, r_2)) &= \psi((s_1 \cdot s_2, s_1\rho(s_2) - \rho(s_1 \cdot s_2) + \rho(s_1)s_2 + s_1r_2 + r_1s_2)) \\ &= \varphi(s_1r_2 + r_1s_2 + s_1\rho(s_2) - \rho(s_1 \cdot s_2) + \rho(s_1) \cdot s_2) + \varphi(\rho(s_1 \cdot s_2)) \\ &= (s_1, r_1)\psi((s_2, r_2)) + \psi((s_1, r_1))(s_2, r_2)\end{aligned}$$

Therefore,

$$\begin{aligned}\text{Ext}_A^1(V_i, V_j) &= \text{Hom}_{A^{\text{op}} \otimes A}(\text{Rad}(A)/\text{Rad}(A)^2, \text{Hom}_k(V_i, V_j)) \\ &= \{\varphi : \text{Rad}(A)/\text{Rad}(A)^2 \rightarrow \text{Hom}_k(V_i, V_j) \mid \forall a \in A, r \in \text{Rad}(A), \text{ s.t.} \\ &\quad \varphi(a \cdot r) = a \cdot \varphi(r) \text{ and } \varphi(ra) = \varphi(r) \cdot a\}\end{aligned}$$

Since $\text{Rad}(A)/\text{Rad}(A)^2 \simeq (E_{ij})$ with

$$E_{ij} \simeq (V_i^* \otimes V_j)^{r_{ij}}$$

it is clear that

$$\begin{aligned}\text{Hom}_{A^{\text{op}} \otimes A}(\text{Rad}(A)/\text{Rad}(A)^2, \text{Hom}_k(V_i, V_j)) \\ \simeq \text{Hom}_{\text{End}_k(V_i)^{\text{op}} \otimes \text{End}_k(V_j)}((V_i^* \otimes V_j)^{r_{ij}}, (V_i^* \otimes V_j)) \\ \simeq k^{r_{ij}}\end{aligned}$$

which means that

$$E_{ij} \simeq \text{Ext}_A^1(V_i, V_j)^* \otimes_k \text{Hom}_k(V_i, V_j).$$

□

Now suppose, as above, that A is a finite dimensional k -algebra, and let $\mathcal{V}_A = \{V_i\}_{i=1}^r$ be any family of finite dimensional A -modules. Obviously

$$\dim_k \text{Ext}_A^p(V_i, V_j) < \infty$$

for all $p = 0, 1, 2, \dots$ and therefore the endomorphism ring

$$\mathcal{O}(\mathcal{V}_A) := \text{End}_H(\tilde{V})$$

is a k -algebra such that

$$\mathcal{O}(\mathcal{V})/\text{Rad}(\mathcal{O}) = \bigoplus_{i=1}^r \text{End}_k(V_i).$$

This implies that $\mathcal{V} = \{V_i\}_{i=1}^r$ is the family of all simple $\mathcal{O}(\mathcal{V})$ -modules. The generalized Burnside theorem applies also in this case, showing that the operation

$$(A, \mathcal{V}) \mapsto (\mathcal{O}(\mathcal{V}), \mathcal{V})$$

is a closure operation. Moreover, we have the following,

Proposition 4.6. *Let $\tau : A \rightarrow B$ be any homomorphism of finite dimensional k -algebras. Consider a family $\mathcal{V}_B = \{V_i\}_{i=1}^r$ of finite dimensional B -modules and let \mathcal{V}_A be the corresponding family of A -modules. Suppose moreover that \mathcal{V}_B is the family of all simple B -modules. Then there exists an, up to isomorphisms, unique homomorphism of k -algebras*

$$O(\tau) : O(\mathcal{V}_A) \rightarrow O(\mathcal{V}_B) \simeq B$$

extending τ .

Proof. There is an obvious forgetful functor defining a morphism of functors on \underline{a}_r ,

$$\tau^* : Def_{\mathcal{V}_B} \rightarrow Def_{\mathcal{V}_A}$$

which in its turn induces a k -algebra homomorphism

$$\eta : H(\mathcal{V}_B) \rightarrow H(\mathcal{V}_A)$$

unique up to isomorphisms, and therefore a k -algebra homomorphism

$$O(\mathcal{V}_A) := (H(\mathcal{V}_A)_{i,j} \otimes Hom_k(V_i, V_j)) \rightarrow (H(\mathcal{V}_B)_{i,j} \otimes Hom_k(V_i, V_j)) =: O(\mathcal{V}_B)$$

obviously extending τ . By the generalized Burnside theorem, $O(\mathcal{V}_B) \simeq B$, and the Proposition follows.

□

Remark 4.7. Up to now we have only considered finite families of A -modules such that

$$\dim_k Ext_A^p(V_i, V_j) < \infty, \quad p = 1, 2.$$

Neither of these conditions are essential. Introducing natural topologies we may, as in [La 1], treat general families of finite type A -modules. Notice also that if $r_1 \leq r_2$, there is an obvious canonical morphism

$$\underline{a}_{r_1} \rightarrow \underline{a}_{r_2}$$

inducing a restriction morphism of functors

$$Def_{\mathcal{V}(2)} \rightarrow Def_{\mathcal{V}(1)}$$

where $\mathcal{V}(1) = \{V_i\}_{i=1}^{r_1}$, $\mathcal{V}(2) = \{V_i\}_{i=1}^{r_2}$. Therefore we obtain an up to isomorphisms unique k -algebra homomorphism

$$r_{2,1} : H_{A, \mathcal{V}(2)} \rightarrow H_{A, \mathcal{V}(1)}.$$

However, this *restriction* morphism is not, in general, unique. The resulting problems will be dealt with later.

§5. Filtered modules and iterated extensions. Let as above $\mathcal{V} = \{V_i\}_{i=1}^r$ be a family of right A -modules, and let $E_{i_1, \dots, i_s} : E_s \subset E_{s-1} \subset \dots \subset E_1 = E$ be a filtered module such that $E_k/E_{k+1} \simeq V_{i_k}$. We shall, as before, refer to any such filtered module as an iterated extension of \mathcal{V} . Notice that for every p there is an extension,


$$\xi_{p,p+1} \in \text{Ext}_A^1(V_{i_p}, V_{i_{p+1}})$$

given by the exact sequence.

$$0 \rightarrow E_{p+1}/E_{p+2} \rightarrow E_p/E_{p+2} \rightarrow E_p/E_{p+1} \rightarrow 0.$$

Corresponding to the iterated extension E_{i_1, \dots, i_s} we shall associate two ordered directed graphs, $\Gamma(\underline{i})$ and $\Gamma(E_{\underline{i}})$. The first is gotten as the graph with nodes in bijection with the modules of the family \mathcal{V} , and with arrows $\gamma_{i_p, i_{p+1}} := \epsilon(i_p, i_{p+1})$ connecting the node i_p with the node i_{p+1} . The second, *the extension type* of the iterated extension, is obtained from the first identifying two arrows $\epsilon(i_p, i_{p+1})$ and $\epsilon(i_q, i_{q+1})$ if the corresponding extensions $\xi_{p,p+1}$ and $\xi_{q,q+1}$ coincide. The "ordered" k -algebra $k[\Gamma]$ of the ordered graph Γ is the quotient algebra of the usual algebra of the graph Γ by the ideal generated by all admissible words which are not "intervals" of the ordered graph. Say $\dots \gamma_{i,j}(n-1) \gamma_{j,j}(n) \gamma_{j,k}(n+1) \dots$ is an interval of the ordered graph, then $\gamma_{i,j}(n-1) \cdot \gamma_{j,k}(n+1) = 0$ in $k[\Gamma]$.

Example 5.1. Let us draw up all extension types for $r, s \leq 3$.



The last example is a $\Gamma(E_{\underline{i}})$ corresponding to the situation, $i_1 = i_2 = i_3 = 1$, and $\xi_{i_1, i_2} = \xi_{i_2, i_3}$. The associated k -algebras are, respectively, the matrix algebras,

$$\begin{pmatrix} k & k & k \\ 0 & k & k \\ 0 & 0 & k \end{pmatrix} \begin{pmatrix} k[\epsilon] & k[\epsilon] \\ 0 & k \end{pmatrix} \begin{pmatrix} k & k[\epsilon] \\ 0 & k[\epsilon] \end{pmatrix} \begin{pmatrix} k[t_{1,2}t_{2,1}] & kt_{1,2} \\ kt_{2,1} & k \end{pmatrix}$$

with the obvious relations, and the k -algebras,

$$k\{t, u\}/(t^2, u^2, ut), \quad k[t]/(t^3)$$

Lemma 5.2. Let H be any object of $\hat{\underline{a}}_r$, and let $R \in \underline{a}_r$. Then $\text{Mor}(H, R)$ has a natural structure of an affine algebraic scheme $\text{Mor}(H, R) = \text{Spec}(A(H, R))$, and there is a universal morphism,

$$\tilde{\phi} : H \longrightarrow A(H, R) \otimes_k R$$

Proof. Put $(E_{i,j}) = \text{rad}(H)/\text{rad}(H)^2$, and consider the affine space,

$$\mathbf{A}^N = \prod_{i,j} E_{i,j}^* \otimes_k R_{i,j}$$

with coordinates $z_{i,j}(l, m) = t_{i,j}(l) \otimes x_{i,j}(m)$, where $\{t_{i,j}(l)\}_{i,j}$ is a basis of $E_{i,j}$ and $\{r_{i,j}(m)\}_{i,j}$ is a basis, and $\{x_{i,j}(m)\}_{i,j}$ is a dual basis of $R_{i,j}$. An element

$$(\alpha_{i,j}(l, m)) \in \mathbf{A}^N$$

corresponds to a morphism $\phi \in \text{Mor}(H, R)$ if, and only if, the corresponding map

$$t_{i,j}(l) \mapsto \sum_m \alpha_{i,j}(l, m) r_{i,j}(m) \in R_{i,j}$$

satisfies the relations of H . Let these, modulo a high enough power of the radical, be polynomials in the generators $t_{i,j}(l)_{i,j}$ of the form

$$f_p(t_{i,j}(l)) = 0, \quad p = 1, \dots, s,$$

and let the relations of R be expressed in terms of,

$$r_{i,j}(m) r_{j,k}(n) = \sum_p \beta_{i,j,k}^p(m, n) r_{i,k}(p), \quad i, j = 1, \dots, r.$$

Then we obtain equations for $\text{Mor}(H, R)$ given by (commutative) polynomial relations of the form,

$$F_p(z_{i,j}(l, m)) = 0, \quad p = 1, \dots, t.$$

But then

$$t_{i,j}(l) \mapsto \sum_m z_{i,j}(l, m) r_{i,j}(m) \in A(H, R) \otimes_k R_{i,j}$$

where the coordinates $z_{i,j}(l, m)$ are subject to the conditions above, defines the universal morphism $\tilde{\phi}$. \square

Proposition 5.3. *Let A be any k -algebra, $\mathcal{V} = \{V_i\}_{i=1}^r$ any swarm of A -modules, i.e. such that,*

$$\dim_k \text{Ext}_A^1(V_i, V_j) < \infty \quad \text{for all } i, j = 1, \dots, r.$$

(i): *Consider an iterated extension E of \mathcal{V} , with directed graph Γ . Then there exists a morphism of k -algebras*

$$\phi : H(\mathcal{V}) \rightarrow k[\Gamma]$$

such that

$$E \simeq k[\Gamma] \otimes_{\phi} \tilde{V}$$

in the above sense.

(ii): The set of equivalence classes of iterated extensions of \mathcal{V} with extension type Γ , is a quotient of the set of closed points of the affine algebraic scheme

$$\underline{A}[\Gamma] = \text{Mor}(H(\mathcal{V}), k[\Gamma])$$

(iii): There is a versal family $\tilde{V}[\Gamma]$ of A -modules defined on $A[\Gamma]$, containing as fibres all the isomorphism classes of iterated extensions of \mathcal{V} s with extension type Γ .

Proof. Any morphism $\varphi : H \rightarrow k[\Gamma]$ in \underline{a}_r correspond to an iterated extension of the V_i 's. This may be expressed in the following way. As vector spaces, we have an isomorphism,

$$k[\Gamma] \otimes_{\varphi} \tilde{V} \simeq V(\Gamma) \simeq V_{i_1} \times V_{i_2} \times \cdots \times V_{i_s}$$

An A -module structure on this vectorspace, corresponding to an iterated extension with extension type Γ , is given by a homomorphism of k -algebras,

$$\psi : A \longrightarrow \text{End}_k(V_{i_1} \times V_{i_2} \times \cdots \times V_{i_s})$$

inducing a family of linear maps

$$\psi_{p,p+1,\dots,p+q} : A \longrightarrow \text{End}_k(V_{i_p}, V_{i_{p+q}})$$

for $0 \leq p < p+q \leq s$.

Consider these maps as 1-cochains in the Hochschild complex

$$HC^*(A, \text{Hom}_k(V(\Gamma), V(\Gamma)))$$

The maps $\psi_{p,p+1}$ correspond to the extensions $\xi_{p,p+1}$ above, and must therefore be 1-cocycles, or derivations. To obtain an A -module structure, corresponding to an iterated extensions of \mathcal{V} with extension type Γ , the conditions on these cochains are: For all $a, b \in A$,

$$\begin{aligned} \psi_{p,p+1}(a)\psi_{p+1,p+2}(b) &= d\psi_{p,p+1,p+2}(a, b) \\ \psi_{p,p+1}(a)\psi_{p+1,p+2,p+3}(b) + \psi_{p,p+1,p+2}(a)\psi_{p+2,p+3}(b) &= d\psi_{p,p+1,p+2,p+3}(a, b) \\ \dots & \\ \sum_{m=2,\dots,s-1} \psi_{1,2,\dots,m}(a)\psi_{m,\dots,s}(b) &= d\psi_{1,2,\dots,s}(a, b), \end{aligned}$$

which means that all Massey products of the form

$$\langle \xi_{i_p, i_{p+1}}, \xi_{i_{p+1}, i_{p+2}}, \dots, \xi_{i_{p+q-1}, i_{p+q}} \rangle$$

are defined and contain zero.

Now (i) follows from the very definition of H , generated as it is by a basis of the dual Ext^1 's, with relations exactly expressing the vanishing of the above Massey products. (ii) and (iii) then follows from deformation theory, together with the Lemma (4.9), above. \square

Example 5.4. Consider the extension E_{ijk} of length 3 given as the composite extension of $\xi_{i,j} : 0 \leftarrow V_i \leftarrow E_{ij} \leftarrow V_j \leftarrow 0$ and $\xi_{i,j,k} : 0 \leftarrow E_{ij} \leftarrow E_{ijk} \leftarrow V_k \leftarrow 0$. Take the pullback $\xi_{i,k}$ of $\xi_{i,j,k}$ via $V_j \rightarrow E_{ij}$ and consider the diagram

$$\begin{array}{ccccccc}
& & 0 & & 0 & & 0 \\
& & \uparrow & & \uparrow & & \uparrow \\
0 & \leftarrow & V_i & \leftarrow & E_{ij} & \leftarrow & V_j & \leftarrow & 0 \\
& & \uparrow & & \uparrow & & \uparrow & & \\
& & E_{ik} & \leftarrow & E_{ijk} & \leftarrow & E_{jk} & & \\
& & \uparrow & & \uparrow & & \uparrow & & \\
& & V_k & = & V_k & = & V_k & & \\
& & \uparrow & & \uparrow & & \uparrow & & \\
& & 0 & & 0 & & 0 & &
\end{array}$$

Let $\psi_{i,j} \in \text{Der}_k(A, \text{Hom}_k(V_i, V_j))$ be a Hochschild cocycle representing the class $\xi_{i,j}$. The multiplication with $a \in A$ on E_{ij} , identified with $V_j \times V_i$ as k -vectorspace, is given by

$$(v_j, v_i)a = (v_j a + \psi_{i,j}(a, v_i), v_i a)$$

and the multiplication with $a \in A$ on E_{ijk} identified with $V_k \times V_j \times V_i$ as k -vectorspace, must be such that,

$$(v_k, v_j, 0)a = (v_k a + \psi_{j,k}(a, v_j), v_j a, 0).$$

If there exists an action of A on E_{ijk} consistent with the above, then one proves the existence of a Hochschild cochain

$$\psi_{i,j,k} \in \text{Hom}_k(A, \text{Hom}_k(V_i, V_k))$$

such that

$$d\psi_{i,j,k}(a, b) = \psi_{i,j}(a)\psi_{j,k}(b).$$

From this follows that the cup-product $\xi_{ij} \cup \xi_{jk}$ is 0 in $\text{Ext}_A^2(V_i, V_k)$. This is also the criterion for the existence of $\xi_{i,j,k}$. Moreover if ξ and ξ' are two extensions

$$\begin{array}{l}
\xi : 0 \leftarrow E_{ij} \leftarrow E_{ijk} \leftarrow V_k \leftarrow 0 \\
\xi' : 0 \leftarrow E_{ij} \leftarrow E'_{ijk} \leftarrow V_k \leftarrow 0
\end{array}$$

with the same pullback ξ_{jk} , then there is an extension

$$\xi_{ik} : 0 \leftarrow V_i \leftarrow E_{ik} \leftarrow V_k \leftarrow 0$$

such that its pullback, via $E_{ij} \rightarrow V_i$, is the difference $\xi - \xi'$.

Consider for the iterated extensions E_{ijk} , the extension diagram

$$\Gamma : \quad \textcircled{i} \longrightarrow \textcircled{j} \longrightarrow \textcircled{k}$$

the first one of the example (5.1) above, then the corresponding k -algebra is given by,

$$k[\Gamma] = \begin{matrix} & & i & j & k \\ \begin{matrix} i \\ j \\ k \end{matrix} & \begin{pmatrix} k & 0 & 0 & 0 & 0 \\ \ddots & \dots & \dots & \dots & \vdots \\ & k & k & k & 0 \\ & & \ddots & \dots & \vdots \\ & & & k & k & 0 \\ & & & & \ddots & \vdots \\ & & & & & k & 0 \\ & & & & & & \ddots & \dots \\ & & & & & & & k \end{pmatrix} \end{matrix}$$

Notice that E_{ijk} then corresponds to a morphism

$$\phi : H \rightarrow k[\Gamma]$$

determined, modulo $Rad^2(k[\Gamma])$ (i.e. the radical squared), by

$$\xi_{i,j} = \phi|_{E_{i,j}} \rightarrow k, \text{ and } \xi_{j,k} = \phi|_{E_{j,k}} \rightarrow k$$

Here

$$Rad^2(k[\Gamma]) = \begin{matrix} & & i & j & k \\ \begin{matrix} i \\ j \\ k \end{matrix} & \begin{pmatrix} 0 & 0 & 0 & 0 & 0 \\ \ddots & \dots & \dots & \dots & \vdots \\ & 0 & 0 & k & 0 \\ & & \ddots & \dots & \vdots \\ & & & 0 & 0 & 0 \\ & & & & \ddots & \vdots \\ & & & & & 0 & 0 \\ & & & & & & \ddots & \dots \\ & & & & & & & 0 \end{pmatrix} \end{matrix}$$

and ϕ is, according to the analysis above, "calibrated" by the morphism $\phi_{i,k} : E_{i,k} \rightarrow k$, i.e. by $Ext_A^1(V_i, V_k)$, as it should.

Corollary 5.5. (i): Given any finitely generated module M on a Noetherian ring A , there is a finite set of primes $E(M)$, containing the set of associated primes $Ass(M)$, such that the module M is an iterated extensions of the corresponding modules A/\mathfrak{p} for $\mathfrak{p} \in E(M)$. The extension type of such an iterated extension is an ordered directed graph $\Gamma(M)$ the nodes of which is $E(M)$.

(ii): For any finite ordered directed graph Γ , with nodes corresponding to a set of primes $\mathcal{P} \subset Spec(A)$, there is an affine versal family of A -modules \tilde{M} with extension type Γ , and $E(M) = \mathcal{P}$.

Proof. Obvious \square

Now, let $H(|\Gamma|)$ be the formal moduli of the family $|\Gamma|$. We know that any iterated extension of the $\{V_i\}_{i=1}^r$ with *extension type*, i.e. graph, Γ corresponds to a morphism in \underline{a}_r ,

$$\alpha : H \longrightarrow k[\Gamma].$$

Moreover the set of isomorphism classes of such modules is parametrized by a quotient space of the affine scheme,

$$\underline{A}(\Gamma) := \text{Mor}_{\underline{a}_r}(H(|\Gamma|), k[\Gamma]).$$

Let $\alpha \in \underline{A}(\Gamma)$, and let $V(\alpha)$ denote the corresponding iterated extension module, then the tangent space of $\underline{A}(\Gamma)$ at α is,

$$T_{\underline{A}(\Gamma), \alpha} := \text{Der}_k(H(|\Gamma|), k[\Gamma]_\alpha),$$

where $k[\Gamma]_\alpha$ is $k[\Gamma]$ considered as a $H(|\Gamma|)$ -bimodule via α . The obstruction space for the deformation functor of α is $HH^2(H(|\Gamma|), k[\Gamma])$, and we may, as is explained in [La 1], compute the complete local ring of $\underline{A}(\Gamma)$ at α . In particular we may decide whether the point is a smooth point of $\underline{A}(\Gamma)$, or not.

The automorphism group G of $k[\Gamma]$, considered as an object of \underline{a}_r , has a Lie algebra which we shall call \mathfrak{g} . Obviously we have,

$$\mathfrak{g} = \text{Der}_k(k[\Gamma], k[\Gamma]).$$

Clearly an iterated extension α with graph Γ will be isomorphic as A -module to $g(\alpha)$, for any $g \in G$. In particular, if $\delta \in \mathfrak{g}$, then $\exp(\delta)(\alpha)$ is isomorphic to α as an iterated extension of A -modules, with the same graph as α .

Consider the map,

$$\alpha^* : \text{Der}_k(k[\Gamma], k[\Gamma]) \rightarrow \text{Der}_k(H(\Gamma), k[\Gamma]_\alpha).$$

The image of α^* is the subspace of the tangent space of $\underline{A}(\Gamma)$ at α along which the corresponding module has constant isomorphism class.

Notice that if α is a smooth point, and α^* is not surjective then there is a positive-dimensional moduli space of iterated extension modules with graph Γ through α .

Clearly the kernel of α^* is contained in the Lie algebra of automorphisms of the module $V(\alpha)$, and should be contained in $\text{End}_A(V(\alpha))$. From this follows that if $V(\alpha)$ is indecomposable then $\ker \alpha^* = 0$. The Euler type derivations, defined by,

$$\delta_E(\gamma_{i,j}) = \rho_{i,j} \gamma_{i,j}, \quad \rho_{i,j} \in k$$

are the easiest to check! Notice however, that there may be discrete automorphisms in G , not of exponential type, leaving α invariant. Notice also that an indecomposable module may have an endomorphism-ring which is a non-trivial local ring.

§6. Noncommutative algebraic geometry.

To any, not necessarily finite, swarm $\underline{c} \subset \underline{\text{mod}}(A)$ of right- A -modules, we may associate two associative k -algebras, see [La 5], $O(|\underline{c}|, \pi) = \varprojlim_{\mathcal{V} \subset |\underline{c}|} O(\mathcal{V})$, and a sub-quotient $\mathcal{O}_\pi(\underline{c})$, together with natural k -algebra homomorphisms,

$$\eta(|\underline{c}|) : A \longrightarrow O(|\underline{c}|, \pi)$$

and,

$$\eta(\underline{c}) : A \longrightarrow \mathcal{O}_\pi(\underline{c})$$

with the property that the A -module structure on \underline{c} is extended to an \mathcal{O} -module structure in an optimal way.

Definition 6.1. A swarm \underline{c} of right A -modules, such that $\eta(\underline{c})$ is an isomorphism will be called an *affine non-commutative scheme* for A .

In particular, for finitely generated k -algebras, the (usually infinite) diagram $Simp_{<\infty}^*(A)$ consisting of the finite dimensional simple A -modules, and the *generic point* A , together with all morphisms between them, is a swarm. This is easily proved. Moreover, consider for some finite family $\mathcal{V} = \{V_i\}_{i=1}^r \subset Simp(A)$ and for any family of morphism $\Phi = \{\phi_i : A \rightarrow V_i\}$ in $Simp_{<\infty}^*(A)$ the corresponding versal lifting,

$$\Phi_{\mathcal{V},\Phi} : A \longrightarrow \tilde{V} = (H_{i,j} \otimes V_j),$$

then we find that,

$$\mathcal{O} := \mathcal{O}_\pi(Simp_{<\infty}^*(A)) \subset End_k(A),$$

and $\alpha \in \mathcal{O}$ if and only if for all finite $\Phi \subset Simp_{<\infty}^*(A)$ there is an $\alpha_{\mathcal{V}} \in End_H(\mathcal{V})$ such that $\alpha_{\mathcal{V}}\Phi_{\mathcal{V},\phi_i} = \Phi_{\mathcal{V},\phi_i}\alpha$.

Now let

$$Simp_{<\infty}(A) = \bigcup_n Simp_n(A)$$

be the set of (iso-classes of) finite dimensional simple right A -modules. An n -dimensional simple A -module $V \in Simp_n(A)$ defines a surjective homomorphism of k -algebras, $\rho : A \rightarrow End_k(V)$, the kernel of which is a two-sided maximal ideal \mathfrak{m}_V , of A . Let $Max_{\leq\infty}(A)$ be the set of all such maximal ideals of A , for $n \geq 1$. To exclude some strange and for our purposes non-interesting cases, we shall assume that our associative k -algebras A have the following property:

$$Rad(A)^\infty := \bigcap_{\mathfrak{m} \in Max_{<\infty}(A), n \geq 0} \mathfrak{m}^n = 0$$

For want of a better name, we shall call such algebras *geometric*. This condition is actually satisfied for most finitely generated k -algebras that we shall be interested in, and in particular it is satisfied for the free k -algebra on d symbols, $A = k \langle x_1, x_2, \dots, x_d \rangle$, see [La 5].

Proposition 6.2. Let A be a geometric k -algebra, then the natural homomorphism,

$$\eta(Simp^*(A)) : A \longrightarrow \mathcal{O}_\pi(Simp_{<\infty}^*(A))$$

is an isomorphism, i.e. $Simp_{<\infty}^*(A)$ is a scheme for A .

Proof. It is an easy consequence of the definition and the next Lemma 6.3 that the morphism $\eta(Simp^*(A))$ is injective for geometric k -algebras. The rest follows from [La 5], (4.1).

□

In particular, $Simp_{<\infty}^*(k \langle x_1, x_2, \dots, x_d \rangle)$, is a scheme for $k \langle x_1, x_2, \dots, x_d \rangle$. To analyze the local structure of $Simp_n(A)$, we need the following, see [La 5],(3.23),

Lemma 6.3. *Let $\mathcal{V} = \{V_i\}_{i=1,\dots,r}$ be a finite subset of $\text{Simp}_{<\infty}(A)$, then the morphism of k -algebras,*

$$A \rightarrow O(\mathcal{V}) = (H_{i,j} \otimes_k \text{Hom}_k(V_i, V_j))$$

is topologically surjective.

Proof. Since the simple modules V_i , $i = 1, \dots, r$ are distinct, there is an obvious surjection, $\eta_0 : A \rightarrow \prod_{i=1,\dots,r} \text{End}_k(V_i)$. Put $\mathfrak{r} = \ker \eta_0$, and consider for $m \geq 2$ the finite-dimensional k -algebra, $B := A/\mathfrak{r}^m$. Clearly $\text{Simp}(B) = \mathcal{V}$, so that by the generalized Burnside theorem, see [La 5], (2.6), we find, $B \simeq O^B(\mathcal{V}) := (H_{i,j}^B \otimes_k \text{Hom}_k(V_i, V_j))$. Consider the commutative diagram,

$$\begin{array}{ccc} A & \longrightarrow & (H_{i,j}^A \otimes_k \text{Hom}_k(V_i, V_j)) =: O^A(\mathcal{V}) \\ \downarrow & & \downarrow \\ B & \longrightarrow & (H_{i,j}^B \otimes_k \text{Hom}_k(V_i, V_j)) \xrightarrow{\alpha} O^A(\mathcal{V})/\mathfrak{m}^m \end{array}$$

where all morphisms are natural. In particular α exists since $B = A/\mathfrak{r}^m$ maps into $O^A(\mathcal{V})/\text{rad}^m$, and therefore induces the morphism α commuting with the rest of the morphisms. Consequently α has to be surjective, and we have proved the contention.

□

Localization and topology. Let $s \in A$, and consider the open subset $D(s) = \{V \in \text{Simp}(A) \mid \rho(s) \text{ invertible in } \text{End}_k(V)\}$. The Jacobson topology on $\text{Simp}(A)$ is the topology with basis $\{D(s) \mid s \in A\}$. It is clear that the natural morphism,

$$\eta : A \rightarrow O(D(s), \pi)$$

maps s into an invertible element of $O(D(s), \pi)$. Therefore we may define the localization $A_{\{s\}}$ of A , as the k -algebra generated in $O(D(s), \pi)$ by $\text{im} \eta$ and the inverse of $\eta(s)$. This furnishes a general method of localization with all the properties one would wish. And in this way we also find a canonical (pre)sheaf, \mathcal{O} defined on $\text{Simp}(A)$.

Definition 6.4. *When the k -algebra A is geometric, such that $\text{Simp}^*(A)$ is a scheme for A , we shall refer to the presheaf \mathcal{O} , defined above on the Jacobson topology, as the structure presheaf of the scheme $\text{Simp}(A)$.*

In the next § we shall see that the Jacobson topology on $\text{Simp}(A)$, restricted to each $\text{Simp}_n(A)$ is the Zariski topology for a classical scheme-structure on $\text{Simp}_n(A)$.

The algebraic (scheme) structure on $\text{Simp}_n(A)$. A standard n -commutator relation in a k -algebra A is a relation of the type,

$$[a_1, a_2, \dots, a_{2n}] := \sum_{\sigma \in \Sigma_{2n}} \text{sign}(\sigma) a_{\sigma(1)} a_{\sigma(2)} \dots a_{\sigma(2n)} = 0$$

where $\{a_1, a_2, \dots, a_{2n}\}$ is a subset of A . Let $I(n)$ be the two-sided ideal of A generated by the subset,

$$\{[a_1, a_2, \dots, a_{2n}] \mid \{a_1, a_2, \dots, a_{2n}\} \subset A\}.$$

Consider the canonical homomorphism,

$$p_n : A \longrightarrow A/I(n) =: A(n).$$

It is known that any homomorphism of k -algebras,

$$\rho : A \longrightarrow \text{End}_k(k^n) =: M_n(k)$$

factors through p_n , see e.g. [Formanek].

Corollary 6.5. (i). Let $V_i, V_j \in \text{Simp}_{\leq n}(A)$ and put $\mathfrak{r} = \mathfrak{m}_{V_i} \cap \mathfrak{m}_{V_j}$. Then we have, for $m \geq 2$,

$$\text{Ext}_A^1(V_i, V_j) \simeq \text{Ext}_{A/\mathfrak{r}^m}^1(V_i, V_j)$$

(ii). Let $V \in \text{Simp}_n(A)$. Then,

$$\text{Ext}_A^1(V, V) \simeq \text{Ext}_{A(n)}^1(V, V)$$

Proof. (i) follows directly from Lemma 4. To see (ii), notice that $\text{Ext}_A^1(V, V) \simeq \text{Der}_k(A, \text{End}_k(V))/\text{Triv} \simeq \text{Der}_k(A(n), \text{End}_k(V))/\text{Triv} \simeq \text{Ext}_{A(n)}^1(V, V)$. The second isomorphism follows from the fact that any derivation maps a standard n -commutator relation into a sum of standard n -commutator relations.

□

Example 6.6. Notice that, for distinct $V_i, V_j \in \text{Simp}_{\leq n}(A)$, we may well have,

$$\text{Ext}_A^1(V_i, V_j) \neq \text{Ext}_{A(n)}^1(V_i, V_j).$$

In fact, consider the matrix k -algebra,

$$A = \begin{pmatrix} k[x] & k[x] \\ 0 & k[x] \end{pmatrix},$$

and let $n = 1$. Then $A(1) = k[x] \oplus k[x]$. Put $V_1 = k[x]/(x) \oplus (0)$, $V_2 = (0) \oplus k[x]/(x)$, then it is easy to see that,

$$\text{Ext}_A^1(V_i, V_j) = k, \quad \text{Ext}_{A(1)}^1(V_i, V_j) = 0, \quad i \neq j,$$

but,

$$\text{Ext}_A^1(V_i, V_i) = \text{Ext}_{A(1)}^1(V_i, V_i) = k, \quad i = 1, 2.$$

Lemma 6.7. *Let B be a k -algebra, and let V be a vector space of dimension n , such that the k -algebra $B \otimes \text{End}_k(V)$ satisfies the standard n -commutator-relations, i.e. such that the ideal, $I(n) \subset B \otimes \text{End}_k(V)$ generated by the standard n -commutators $[x_1, x_2, \dots, x_{2n}]$, $x_i \in B \otimes \text{End}_k(V)$, is zero. Then B is commutative.*

Proof. In fact, if $b_1, b_2 \in B$ is such that $[b_1, b_2] \neq 0$, then the obvious n -commutator,

$$(b_1 e_{1,1})(b_2 e_{1,1})e_{1,2}e_{2,2}\dots e_{n-1,n} \cdot e_{n,n} - (b_2 e_{1,1})(b_1 e_{1,1})e_{1,2}e_{2,2}\dots e_{n-1,n} \cdot e_{n,n}$$

is different from 0. Here $e_{i,j}$ is the $n \times n$ matrix with all elements equal to 0, except the one in the (i, j) position, where the element is equal to 1.

□

Lemma 6.8. *If A is a finite type k -algebra, then any $V \in \text{Simp}_n(A)$ is an $A(n)$ -module. Let $\mathcal{V} \subset \text{Simp}_n(A)$ be a finite family, then $H^{A(n)}(\mathcal{V})$ is commutative. In particular,*

- (1) $\text{Ext}_{A(n)}(V_i, V_j) = 0$, for $V_i \neq V_j$
- (2) $H^{A(n)}(V) \simeq H^A(V)^{\text{com}} := H(V)/[H(V), H(V)]$.

Proof. Since

$$A(n) \rightarrow O(\mathcal{V}) \simeq M_n(H^{A(n)}(\mathcal{V}))$$

is topologically surjective, we find using (Lemma 8), that $H^{A(n)}(\mathcal{V})$ is commutative. This implies (1) and the commutativity of $H^{A(n)}(V)$. Consider for $V \in \text{Simp}_n(A)$, the natural commutative diagram of homomorphisms of k -algebras,

$$\begin{array}{ccc} & A & \\ & \downarrow & \searrow \\ Z(A(n)) & \longrightarrow & A(n) & \longrightarrow & H(V) \otimes_k \text{End}_k(V) \\ & \downarrow & \downarrow \alpha & \swarrow & \\ H(V)^{\text{com}} & \longrightarrow & H(V)^{\text{com}} \otimes_k \text{End}_k(V) & & \end{array}$$

where $Z(A(n))$ is the center of $A(n)$. The existence of α is a consequence of the ideal $I(n)$ of A mapping to zero in $H(V)^{\text{com}} \otimes_k \text{End}_k(V) \simeq M_n(H(V)^{\text{com}})$. Therefore there are natural morphisms of formal moduli,

$$H^A(V) \rightarrow H^{A(n)}(V) \rightarrow H^A(V)^{\text{com}} \rightarrow H^{A(n)}(V)^{\text{com}}.$$

Since $H^{A(n)}(V) = H^{A(n)}(V)^{\text{com}}$ the composition,

$$H^{A(n)}(V) \rightarrow H^A(V)^{\text{com}} \rightarrow H^{A(n)}(V)^{\text{com}},$$

must be an isomorphism. Since by Corollary 4. the tangent spaces of $H^{A(n)}(V)$ and $H^A(V)$ are isomorphic, the lemma is proved.

□

Corollary 6.9. *Let $A = k \langle x_1, \dots, x_d \rangle$ be the free k -algebra on d symbols, and let $V \in \text{Simp}_n(A)$. Then*

$$H^A(V)^{\text{com}} \simeq H^{A(n)}(V) \simeq k[[t_1, \dots, t_{(d-1)n^2+1}]]$$

This should be compared with the results of [Procesi 1], see also [Formanek]. In general the natural morphism,

$$\eta(n) : A(n) \rightarrow \prod_{V \in \text{Simp}_n(A)} H^{A(n)}(V) \otimes_k \text{End}_k(V)$$

is not an injection.

Example 6.10. *In fact, let*

$$A = \begin{pmatrix} k & k & k \\ k & k & k \\ 0 & 0 & k \end{pmatrix}.$$

The ideal $I(2)$ is generated by $[e_{1,1}, e_{1,2}, e_{2,2}, e_{2,3}] = e_{1,3}$. So

$$A(2) = \begin{pmatrix} k & k & k \\ k & k & k \\ 0 & 0 & k \end{pmatrix} / \begin{pmatrix} 0 & 0 & k \\ 0 & 0 & k \\ 0 & 0 & 0 \end{pmatrix} \simeq M_2(k) \oplus M_1(k).$$

However,

$$\prod_{V \in \text{Simp}_2(A)} H^{A(2)}(V) \otimes_k \text{End}_k(V) \simeq M_2(k),$$

therefore $\ker \eta(2) = M_1(k) = k$.

Let $O(n)$, be the image of $\eta(n)$, then,

$$O(n) \subseteq \prod_{V \in \text{Simp}_n(A)} H^{A(n)}(V) \otimes_k \text{End}_k(V)$$

and for every $V \in \text{Simp}_n(A)$,

$$H^{O(n)}(V) \simeq H^{A(n)}(V).$$

Put $B = \prod_{V \in \text{Simp}_n(A)} H^{A(n)}(V)$. Choosing bases in all $V \in \text{Simp}_n(A)$, then

$$\prod_{V \in \text{Simp}_n(A)} H^{A(n)}(V) \otimes_k \text{End}_k(V) \simeq M_n(B),$$

Let $x_i \in A, i = 1, \dots, d$ be generators of A , and consider their images $(x_{p,q}^i) \in M_n(B)$. Now, B is commutative, so the k -sub-algebra $C(n) \subset B$ generated by the elements $\{x_{p,q}^i\}_{i=1, \dots, d; p, q=1, \dots, n}$ is commutative. We have an injection,

$$O(n) \rightarrow M_n(C(n)),$$

and for all $V \in \text{Simp}_n(A)$ there is a natural composition of homomorphisms of k -algebras,

$$\alpha : M_n(C(n)) \rightarrow M_n(H^{A(n)}(V)) \rightarrow \text{End}_k(V),$$

inducing a corresponding composition of homomorphisms of the centers,

$$Z(\alpha) : C(n) \rightarrow H^{A(n)}(V) \rightarrow k$$

This sets up a set theoretical injective map,

$$t : \text{Simp}_n(A) \longrightarrow \text{Max}(C(n)),$$

defined by $t(V) := \ker Z(\alpha)$.

Since $A(n) \rightarrow H^{A(n)}(V) \otimes_k \text{End}_k(V)$ is topologically surjective, $H^{A(n)}(V) \otimes_k \text{End}_k(V)$ is topologically generated by the images of x_i , $i = 1, \dots, d$. It follows that we have a surjective homomorphism,

$$\hat{C}(n)_{t(V)} \rightarrow H^{A(n)}(V).$$

Categorical properties implies, that there is another natural morphism,

$$H^{A(n)}(V) \rightarrow \hat{C}(n)_{t(V)},$$

which composed with the former is an automorphism of $H^{A(n)}(V)$. Since

$$M_n(C(n)) \subseteq \prod_{V \in \text{Simp}_n(A)} H^{A(n)}(V) \otimes_k \text{End}_k(V),$$

it follows that for $\mathfrak{m}_v \in \text{Max}(C(n))$, corresponding to $V \in \text{Simp}_n(A)$, the finite dimensional k -algebra $M_n(C(n)/\mathfrak{m}_v^2)$ sits in a finite dimensional quotient of some,

$$\prod_{V \in \mathcal{V}} H^{A(n)}(V) \otimes_k \text{End}_k(V).$$

where $\mathcal{V} \subset \text{Simp}_n(A)$ is finite. However, by Lemma 4. the morphism,

$$A(n) \longrightarrow \prod_{V \in \mathcal{V}} H^{A(n)}(V) \otimes_k \text{End}_k(V)$$

is topologically surjectiv. Therefore the morphism,

$$A(n) \longrightarrow M_n(C(n)/\mathfrak{m}_v^2)$$

is surjectiv, implying that the map

$$H^{A(n)}(V) \rightarrow \hat{C}(n)_{\mathfrak{m}_v},$$

is surjectiv, and consequently, $H^{A(n)}(V) \simeq \hat{C}(n)_{\mathfrak{m}_v}$.

We now have the following theorem, see Chapter VIII, §2, of the book [Procesi 2], where part of this theorem is proved.

Theorem 6.11. *Let $V \in \text{Simp}_n(A)$, correspond to the point $\mathfrak{m}_v \in \text{Max}(C(n))$. Then there exist a Zariski neighborhood U_v of v in $\text{Max}(C(n))$ such that any closed point $\mathfrak{m}'_v \in U$ corresponds to a unique point $V' \in \text{Simp}_n(A)$. Let $U(n)$ be the open subset of $\text{Max}(C(n))$, the union of all U_v for $V \in \text{Simp}_n(A)$. $O(n)$ defines a non-commutative structure sheaf $\mathcal{O}(n) := \mathcal{O}_{\text{Simp}_n(A)}$ of Azumaya algebras on the topological space $\text{Simp}_n(A)$ (Jacobson topology). The center $\mathcal{S}(n)$ of $\mathcal{O}(n)$, defines a scheme structure on $\text{Simp}_n(A)$. Moreover, there is a morphism of schemes,*

$$\kappa : U(n) \longrightarrow \text{Simp}_n(A),$$

Such that for any $v \in U(n)$,

$$\hat{\mathcal{S}}(n)_{\kappa(v)} \simeq H^{A(n)}(V)$$

Proof. Let $\rho : A \longrightarrow \text{End}_k(V)$ be the surjective homomorphism of k -algebras, defining $V \in \text{Simp}_n(A)$. Let, as above $e_{i,j} \in \text{End}_k(V)$ be the elementary matrices, and pick $y_{i,j} \in A$ such that $\rho(y_{i,j}) = e_{i,j}$. Let us denote by σ the cyclical permutation of the integers $\{1, 2, \dots, n\}$, and put,

$$s_k := [y_{\sigma^k(1), \sigma^k(2)}, y_{\sigma^k(2), \sigma^k(3)} \cdots y_{\sigma^k(n), \sigma^k(1)}], \quad s := \sum_{k=0,1,\dots,n-1} s_k \in A.$$

Clearly $s \in I(n-1)$. Since $[e_{\sigma^k(1), \sigma^k(2)}, e_{\sigma^k(2), \sigma^k(3)} \cdots e_{\sigma^k(n), \sigma^k(1)}] = e_{\sigma^k(1), \sigma^k(n)} \in \text{End}_k(V)$, $\rho(s) := \sum_{k=0,1,\dots,n-1} \rho(s_k) \in \text{End}_k(V)$ is the matrix with non-zero elements, equal to 1, only in the $(\sigma^k(1), \sigma^k(n))$ position, so the determinant of $\rho(s)$ must be $+1$ or -1 . The determinant $\det(s) \in C(n)$ is therefore nonzero at the point $v \in \text{Spec}(C(n))$ corresponding to V . Put $U = D(\det(s)) \subset \text{Spec}(C(n))$, and consider the localization $O(n)_{\{s\}} \subseteq M_n(C(n)_{\{\det(s)\}})$, the inclusion following from general properties of the localization. Now, any closed point $v' \in U$ corresponds to a n -dimensional representation of A , for which the element $s \in I(n-1)$ is invertible. But then this representation cannot have a $m < n$ dimensional quotient, so it must be simple.

Since $s \in I(n-1)$, the localized k -algebra $O(n)_{\{s\}}$ does not have any simple modules of dimension less than n , and no simple modules of dimension $> n$. In fact, for any finite dimensional $O(n)_{\{s\}}$ -module V , of dimension m , the image \hat{s} of s in $\text{End}_k(V)$ must be invertible. However, the inverse \hat{s}^{-1} must be the image of a polynomial (of degree $m-1$) in s . Therefore, if V is simple over $O(n)_{\{s\}}$, i.e. if the homomorphism $O(n)_{\{s\}} \rightarrow \text{End}_k(V)$ is surjective, V must also be simple over A . Since now $s \in I(n-1)$, it follows that $m \geq n$. If $m > n$, we may construct, in the same way as above an element in $I(n)$ mapping into a nonzero element of $\text{End}_k(V)$. Since, by construction, $I(n) = 0$ in $A(n)$, and therefore also in $O(n)_{\{s\}}$, we have proved what we wanted. By a theorem of M. Artin, see [Artin], $O(n)_{\{s\}}$ must be an Azumaya algebra with center, $\mathcal{S}(n)_{\{s\}} := Z(O(n)_{\{s\}})$. Therefore $O(n)$ defines a presheaf $\mathcal{O}(n)$ on $\text{Simp}_n(A)$, of Azumaya algebras with center $\mathcal{S}(n) := Z(\mathcal{O}(n))$. Clearly, any $V \in \text{Simp}_n(A)$, corresponding to $\mathfrak{m}_v \in \text{Max}(C(n))$ maps to a point $\kappa(v) \in \text{Simp}(\mathcal{O}(n))$. Let $\mathfrak{m}_{\kappa(v)}$ be the corresponding maximal ideal of $\mathcal{S}(n)$. Since $O(n)$ is locally Azumaya, it follows that,

$$\hat{\mathcal{S}}(n)_{\mathfrak{m}_{\kappa(v)}} \simeq H^{O(n)}(V) \simeq H^{A(n)}(V).$$

The rest is clear.

□

$\text{Spec}(C(n))$ is, in a sense, a compactification of $\text{Simp}_n(A)$. It is, however not the correct *completion* of $\text{Simp}_n(A)$. In fact, the points of $\text{Spec}(C(n)) - \text{Simp}_n(A)$ may correspond to semi-simple modules, which do not deform into simple n -dimensional modules. We shall in the next § return to the study of the (notion of) completion, together with the degeneration processes that occur, at *infinity* in $\text{Simp}_n(A)$.

Example 6.12. Let us check the case of $A = k \langle x_1, x_2 \rangle$, the free non-commutative k -algebra on two symbols. Pick $V \in \text{Simp}_2(A)$, and let us compute $\text{Ext}_A^1(V, V)$. We would like to find a basis $\{t_i^*,\}_{i=1}^5$, represented by derivations $\psi_i \in \text{Der}_k(A, \text{End}_k(V))$, $i=1,2,3,4,5$. This is easy, since for any two A -modules V_1, V_2 , we have the exact sequence,

$$\begin{aligned} 0 \rightarrow \text{Hom}_A(V_1, V_2) \rightarrow \text{Hom}_k(V_1, V_2) \rightarrow \text{Der}_k(A, \text{Hom}_k(V_1, V_2)) \\ \rightarrow \text{Ext}_A^1(V_1, V_2) \rightarrow 0 \end{aligned}$$

proving that, $\text{Ext}_A^1(V_1, V_2) = \text{Der}_k(A, \text{Hom}_k(V_1, V_2))/\text{Triv}$, where Triv is the sub-vector space of trivial derivations. Pick $V \in \text{Simp}_2(A)$ defined by the homomorphism $A \rightarrow M_2(k)$ mapping the generators x_1, x_2 to the matrices

$$X_1 := \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix} =: e_{1,2}, \quad X_2 := \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix} =: e_{2,1}.$$

Notice that

$$X_1 X_2 = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} =: e_{1,1} = e_1, \quad X_2 X_1 = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} =: e_{2,2} = e_2,$$

and recall also that for any 2×2 -matrix $(a_{p,q}) \in M_2(k)$, $e_i(a_{p,q})e_j = a_{i,j}e_{i,j}$. The trivial derivations are generated by the derivations $\{\delta_{p,q}\}_{p,q=1,2}$, defined by,

$$\delta_{p,q}(x_i) = X_i e_{p,q} - e_{p,q} X_i.$$

Clearly $\delta_{1,1} + \delta_{2,2} = 0$. Now, compute and show that the derivations ψ_i , $i = 1, 2, 3, 4, 5$, defined by,

$$\psi_i(x_1) = 0, \text{ for } i = 1, 2, \quad \psi_i(x_2) = 0, \text{ for } i = 4, 5,$$

by,

$$\psi_1(x_2) = e_{1,1}, \psi_2(x_2) = e_{1,2}, \psi_3(x_1) = e_{1,2}, \psi_4(x_1) = e_{2,1}, \psi_5(x_1) = e_{2,1}$$

and by,

$$\psi_3(x_2) = e_{2,1},$$

form a basis for $\text{Ext}_A^1(V, V) = \text{Der}_k(A, \text{End}_k(V))/\text{Triv}$. Since $\text{Ext}_A^2(V, V) = 0$ we find $H(V) = k \langle\langle t_1, t_2, t_3, t_4, t_5 \rangle\rangle$ and so $H(V)^{\text{com}} \simeq k[[t_1, t_2, t_3, t_4, t_5]]$. The formal versal family \tilde{V} , is defined by the actions of x_1, x_2 , given by,

$$X_1 := \begin{pmatrix} 0 & 1 + t_3 \\ t_5 & t_4 \end{pmatrix}, \quad X_2 := \begin{pmatrix} t_1 & t_2 \\ 1 + t_3 & 0 \end{pmatrix}.$$

One checks that there are polynomials of X_1, X_2 which are equal to $t_i e_{p,q}$, modulo the ideal $(t_1, \dots, t_5)^2 \subset H(V)$, for all $i, p, q = 1, 2$. This proves that $\hat{C}(2)_v$ must be isomorphic to $H(V)$, and that the composition,

$$A \longrightarrow A(2) \longrightarrow M_2(C(2)) \subset M_2(H(V))$$

is topologically surjective. By the construction of $C(n)$ this also proves that

$$C(2) \simeq k[t_1, t_2, t_3, t_4, t_5].$$

Computing we find the following formulas,

$$\begin{aligned} \operatorname{tr} X_1 &= t_4, \quad \operatorname{tr} X_2 = t_1, \\ \det X_1 &= -t_5 - t_3 t_5, \quad \det X_2 = -t_2 - t_2 t_3, \\ \det(X_1 X_2) &= 1 + 2t_3 + t_3^2 + t_2 t_5 \end{aligned}$$

Moreover, the Formanek center, in this case is cut out by the single equation:

$$f := \det[X_1, X_2] = -((1 + t_3)^2 - t_2 t_5)^2 + (t_1(1 + t_3) + t_2 t_4)(t_4(1 + t_3) + t_1 t_5).$$

From this follows that

$$\operatorname{Simp}_2(A) = \mathbf{A}^5 - V(f).$$

§7. The smooth locus of an affine noncommutative scheme.

Definition 7.1. Let $V \in \operatorname{Simp}_n(A)$, then V is called formally smooth if,

$$\operatorname{Ext}_A^2(V, V) = HH^2(A, \operatorname{End}_k(V)) = 0$$

and smooth (or regular is better), if the natural k -linear map,

$$\kappa : \operatorname{Der}_k(A, A) \longrightarrow \operatorname{Ext}_A^1(V, V)$$

is surjective.

Notice that if,

$$HH^2(A, \underline{\mathfrak{m}}) = 0$$

for all maximal two-sided ideals of A , then $\operatorname{Simp}_n(A)$ is smooth for all n .

Let $V \in \operatorname{Simp}_n(A)$, and let $v \in \operatorname{Simp}(Z(A))$ be the point corresponding to V . Denote by \mathfrak{m}_v the corresponding maximal ideal of $Z(A)$. Clearly $Z(A)$ operate naturally on the Hochschild cohomology, $HH^1(A, A)$, and the map κ factors through, $HH^1(A, A)/\mathfrak{m}_v HH^1(A, A)$, so that if V is smooth, we obtain a surjectiv k -linear map,

$$\kappa_0 : HH^1(A, A)/\mathfrak{m}_v HH^1(A, A) \longrightarrow \operatorname{Ext}_A^1(V, V).$$

It follows that $\max_{V \in \operatorname{Simp}_n(A)} \{ \dim_k HH^1(A, A)/\mathfrak{m}_v HH^1(A, A) \}$ is an upper bound for the dimensions of the smooth locus of $\operatorname{Simp}_n(A)$ for all $n \geq 1$.

Clearly the definition of (formal) smoothness also works for any representation V .

Proposition 7.2. *If $V \in \text{Simp}_n(A)$ is smooth or formally smooth, then the corresponding point $v \in \text{Spec}(C(n))$ is also smooth.*

Proof. Assume that $V \in \text{Simp}_n(A)$ is formally smooth, then obviously the completion of the local ring of $\text{Simp}_n(A)$ at V is $H(V)^{\text{com}}$, which since $H(V)$ has no obstructions and therefore must be the completion of the free non-commutative k -algebra, is a formal power series algebra, and thus V is a smooth point of $\text{Simp}_n(A)$.

Now, assume V is smooth, and consider the natural commutative diagram,

$$\begin{array}{ccc}
 & \text{Der}_k(A, A) & \\
 & \downarrow \rho & \\
 & \text{Der}_k(A(n), A(n)) & \\
 & \downarrow \kappa & \searrow \gamma \\
 & \text{Der}_k(O(n), O(n)) & \xrightarrow{\delta} \text{Ext}_A^1(V, V) \\
 & \downarrow \lambda & \nearrow \epsilon \\
 & \text{Der}_k(O(n)_{\{s\}}, O(n)_{\{s\}}) & \\
 & \uparrow \alpha & \uparrow \beta \\
 \text{Der}_k(S(n), S(n)) & \longrightarrow & \text{Der}_k(S(n), k(v)).
 \end{array}$$

Notice that β is an isomorphism. This has been proved above. That ρ exists is easily seen, since for any derivation $\delta \in \text{Der}_k(A)$, and for any standard commutator $[x_1, x_2, \dots, x_{2n}] \in I(n)$, we must have $\delta([x_1, x_2, \dots, x_{2n}]) \in I(n)$. Notice that the kernel of the homomorphism, $A(n) \rightarrow O(n)$ is the image in $A(n)$ of

$$\mathfrak{n} = \bigcap_{\mathfrak{m} \in \text{Max}_n(A), m \geq 1} \mathfrak{m}^m.$$

Clearly any derivation will map an element of \mathfrak{n} into \mathfrak{n} , proving the existence of κ . λ is defined by localization at the point $v \in \text{Spec}(C(n))$, as in the proof of Theorem 12. We may assume $O(n)_{\{s\}}$ is a matrix algebra $M_n(S(n))$, and use the fact that any derivation of a matrix algebra is given by a derivation of the centre and an inner derivation, (HH^1 is Morita invariant). The inner derivation will map to zero in $\text{Ext}_A^1(V, V)$, and so the composition of α and ϵ is surjective.

□

Notice that for a smooth point $v \in \text{Spec}(C(n))$, $\text{Ext}_A^2(V, V)$ may well be different from 0.

Examples 7.3. 1. Let S be any commutative algebra, and denote by $\mathfrak{b} \subseteq \mathfrak{a} \subset S$ two ideals of S . Consider the k -algebra,

$$A := \left\{ \begin{pmatrix} a_{1,1} & a_{1,2} \\ a_{2,1} & a_{2,2} \end{pmatrix} \mid a_{i,j} \in S, a_{1,1} - a_{2,2} \in \mathfrak{a}, a_{1,2}, a_{2,1} \in \mathfrak{b} \right\}.$$

Clearly the centre of $A = A(2) = O(2)$, is $S(2) = C(2) = S$ and a simple calculation shows that,

$$A(1) = \left\{ \left(\begin{array}{cc} \tilde{a}_{1,1} & \tilde{a}_{1,2} \\ \tilde{a}_{2,1} & \tilde{a}_{2,2} \end{array} \right) \mid \tilde{a}_{i,j} \in \mathfrak{b}/\mathfrak{ab}, i \neq j, \tilde{a}_{1,1}, \tilde{a}_{2,2} \in S/\mathfrak{b}^2, \tilde{a}_{1,1} - \tilde{a}_{2,2} \in \mathfrak{a}/\mathfrak{b}^2 \right\}.$$

Then $A(1)$ is the commutative k -algebra expressed by Nagata rings, i.e.

$$A(1) = ((S/\mathfrak{b}^2)[(\mathfrak{a}/\mathfrak{b}^2)])[(\mathfrak{b}/\mathfrak{ab})^2].$$

Consider the subschemes $V(\mathfrak{a}) \subset V(\mathfrak{b}) \subset \text{Spec}(S)$. Then, $\text{Simp}_2(A) = \text{Spec}(S) - V(\mathfrak{b})$ and a simple calculation shows that $\text{Simp}_1(A) = \text{Spec}(A(1))$ is a thickening of the affine scheme $\text{Spec}((S/\mathfrak{b}^2)[(\mathfrak{a}/\mathfrak{b}^2)])$. In the special case,

$$S = k[t_1, t_2], \mathfrak{a} = (f, g), \mathfrak{b} = (f)$$

where $f, g \in S$, correspond to two curves, $V(f), V(g)$ that intersect in a finite set U , one finds that $\text{Simp}_2(A)$ is an open affine subscheme of $\text{Spec}(S)$, and that $\text{Simp}_1(A) = \text{Spec}(A(1))$ is the disjoint union of the curve $V(f)$ with itself, amalgamated at the points of U . If both $V(f)$ and $V(g)$ are smooth, and intersect normally at the points of U , then the embedding-dimension of $\text{Simp}_1(A) = \text{Spec}(A(1))$ at a point not in U , is 2, and at the points of U , 6!

2. Let in the above example, $\mathfrak{b} = \mathfrak{a} = (t_1, t_2)$, then $\text{Simp}_2(A) = \text{Spec}(S) - \{(0, 0)\}$, therefore not affine, and $\text{Simp}_1(A) = \text{Spec}(A(1))$ is a thick point situated at the origin of the affine 2-space $\text{Spec}(S)$.

Example 7.4. Let us try to compute the $\text{Simp}_2(A)$ for the non-commutative cusp, i.e. for the k -algebra,

$$A = k \langle x, y \rangle / (x^3 - y^2).$$

We first notice that the center $Z(A) \subset A$ is the subalgebra of A generated by $t := x^3 - y^2$. Put

$$u_1 = x^2y, v_1 = yx^2.$$

Then there is a surjective morphism,

$$k[t, t^{-1}] \langle u, v \rangle / (uvu - vuv) \longrightarrow A(t^{-1})$$

mapping u to u_1 and v to v_1 . In fact, $u_1v_1 = t^2x$ and $v_1u_1v_1 = t^3y$, and finally $u_1v_1u_1 = t^3y = v_1u_1v_1$. (The relations with the equation of Yang-Baxter, if any, will have to be discovered.)

Now let us compute the $\text{Simp}_n(A)$. It is clear that any surjectiv homomorphism of k -algebras,

$$\rho_v : A \longrightarrow \text{End}_k(V)$$

will map $Z(A) = k[t]$ into $Z(\text{End}_k(V)) = k$, inducing a point $v \in \text{Simp}(k[t]) = \mathbf{A}^1$. This means that $\text{Simp}_n(A)$ is fibred over the affine line $\text{Spec}(k[t]) = \mathbf{A}^1$. Let $\rho_v(x)^3 = \rho_v(y)^2 = \kappa(v)\mathbf{1}$, where $\mathbf{1}$ is the identity matrix, and where $\kappa(v)$ is a

parameter of the cusp. Then either $v = origin =: \underline{0}$ or we may assume $\kappa(v) \neq 0$. Consider now the diagram:

$$\begin{array}{ccc}
 k[t = x^3 = y^2] & & \\
 \downarrow & \searrow & \\
 A & \xrightarrow{\rho_v} & End_k(V) \\
 \downarrow & \nearrow & \\
 k[x]/(x^3 - \kappa(v)) * k[y]/(y^2 - \kappa(v)) & &
 \end{array}$$

Clearly, if $\kappa(v) \neq 0$ the simple representations of A are fibered on the cusp with fibres being the simple representations of $U := k[x]/(x^3 - \kappa(v)) * k[y]/(y^2 - \kappa(v))$, isomorphic to the group algebra of the modular group $Sl_2(\mathbf{Z})$. Since the representation theory of $Sl_2(\mathbf{Z})$ is well known at least in small dimensions, we could hope to use classical representation theory to describing the open subscheme of $Simp_n(A)$ corresponding to $\kappa(v) \neq 0$, for all $n \geq 1$. We shall, however see that deformation theory can be very usefull to understand the representations of $Sl_2(\mathbf{Z})$.

Moreover we shall have to work a little to find the fibre of $Simp_n(A)$ corresponding to the singular point of the cusp. When $n = 2$ it is clear that we have no choice, but to fix the Jordan form of $\rho_v(y)$ equal to the Jordan form of

$$\rho_v(x) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}.$$

Let $I(\rho_v(x))$ be the isotropy subgroup of the action of $Gl_n(k)$ on $M_n(k)$, at $\rho_v(x)$. Set theoretically, the fiber is then the double quotient,

$$I(\rho_v(x)) \backslash Gl_n(k) / I(\rho_v(x))$$

To find the scheme structure we may compute the formal moduli of the simple module given by,

$$\rho_v(x) = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}, \rho_v(y) = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}.$$

We find the following, see [Jø-La-Sl].

Proposition 7.5. *Let A be the non-commutative cusp. Then*

- (i) $Simp_1(A) = Spec(k[x, y]/(x^3 - y^2))$
- (ii) $Simp_2(A)$ is fibered on the cusp minus the origin, with fiber $E(\underline{t}) = U_2/T^2$ where U_2 is an open subscheme of the 3-dimensional scheme of all pairs of 2-vectors, with vector product equal 1, and T^2 is a two dimensional torus, acting naturally on U_2 .
- (iii) $S(2) = k[t^2, t^3, u]$.
- (iv) The fiber $E(\underline{0})$ over $\underline{0}$ is given by,

$$\tilde{\rho}(x) = \begin{pmatrix} t & 1 \\ 0 & t \end{pmatrix}, \tilde{\rho}(y) = \begin{pmatrix} u & 0 \\ 1+v & -u \end{pmatrix}$$

parametrized by the k -algebra $k[t, u, v]/(t^2, u^2, (1+v)t)$, i.e. it is the open subscheme of the double line parametrized by v , with the point $v = -1$ removed.

- (v) In particular we find that $E(\underline{0})$ is a component of $Simp_2(A)$.

We end these notes with some usefull references:

REFERENCES

- [Artin] Artin, M.: On Azumaya Algebras and Finite Dimensional Representations of Rings. *Journal of Algebra* 11, 532-563 (1969).
- [A-T-vdB] Artin, M. and Tate, J. and Van den Bergh, M.: Some algebras associated to automorphisms of elliptic curves. *Grothendieck Festschrift, Vol 1. Progress in Math. Vol 86, Birkhäuser*, pp. 33-85, (1990).
- [Connes] Connes, A.: *Géométrie Noncommutative*. InterEditions, (1990)
- [Curtis and Reiner] Curtis, C.W. and Reiner, I.: *Representation Theory of finite Groups and associative Algebras*. Pure and applied Mathematics, Interscience Publishers, (1962).
- [Eis] Eisenbud, D.: Homological algebra on a complete intersection with application to group representation. *Trans.Amer.Math.Soc.* vol 260, (1980), pp. 35-64.
- [Formanek] Formanek, E.: *The polynomial identities and invariants of $n \times n$ Matrices*. Regional Conferenc Series in Mathematics, Number 78. Published for the Conference Board of the Mathematical Sciences by the American Mathematical Society, Providence, Rhode Island, (1990)
- [G] Gabriel, P.: Unzerlegbare Darstellungen I, II. *Mauscripta Math.* 6, (1972), pp. 71-103. *Symposia Math.Inst.Naz.Alt.Mat.* 11, (1973), pp.81-104.
- [Gr-K] Greuel, G.M. and Knörrer, H.: Einfache kurvensingularitäten und torsionsfreie modulen. *Math.Ann.* vol 270, (1985), pp. 417-425.
- [J.L.S.] Jndrup, S., Laudal, O.A., Sletsje, A.B.: *Noncommutative plane curves*. Institute Mittag-Leffler. Report No. 20, 2003/2004..
- [Ile] Ile, R.: *noncommutative deformations. $ab \neq ba$* . Master Thesis, University of Oslo, 1990.
- [K] Kadison L.: *New Examples of Frobenius Extensions*, AMS-University Lecture Series, Vol 14., (1999).
- [Lang] Lang, S.: *Algebra*, Addison-Wesley, (1965).
- [La 0] Laudal, O.A.: Sur les limites projectives et inductives. *Ann.Sci.Ecole Normale Sup.*, Paris, t. 82, (1965).
- [La 1] Laudal, O.A.: Formal moduli of algebraic structures. *Lecture Notes in Mathematics*, Springer Verlag, vol. 754, (1979)
- [La 2] Laudal, O.A.: Matric Massey products and formal moduli I. in (Roos, J.E.ed.) *Algebra, Algebraic Topology and their interactions* *Lecture Notes in Mathematics*, Springer Verlag, vol 1183, (1986) pp. 218-240.
- [La-Pf] Laudal, O.A. and Pfister, G.: *Local moduli and singularities*. *Lecture Notes in Mathematics*, Springer Verlag, vol. 1310, (1988)
- [La 3] Laudal, O.A.: Noncommutative deformations of modules. *Homology, Homotopy and Applications*. Vol 4 (2), 2002, pp. 357-396.
- [La 4] Laudal, O.A.: *Noncommutative Algebraic Geometry*. Max-Planck-Institut für Mathematik, Preprint Series 2000 (115).
- [La 5] Laudal, O.A.: Noncommutative algebraic geometry. *Rev. Mat. Iberoamericana* 19 (2003), 509-580.
- [La 6] Laudal, O.A.: The structure of $Simp_{<\infty}(A)$ for finitely generated k -algebras. Manuscript.
- [Manin] Manin, Y.I.: *Topics in Noncommutative Geometry*. Princeton University Press, (1991).
- [Procesi] Procesi, C.: Non-commutative affine rings. *Atti Accad. Naz. Lincei Rend.Cl. Sci. Fis. Mat. Natur.* (8)(1967), 239-255
- [R] Reiten, I.: An introduction to representation theory of Artin algebras. *Bull.London Math. Soc.* 17, (1985), pp. 209-233.
- [Romstad] Romstad, T.: *Non-commutative Algebraic Geometry Applied to Invariant Theory of Finite Group Actions*. Cand.Scient.Thesis. Institute of Mathematics, University of Oslo (2000).
- [Rosenberg] Rosenberg, A.L.: *Noncommutative algebraic geometry and representations of quantized algebra*. Klüwer, Mathematics and its applications. vol. 330, (1995)
- [Sch] Schlessinger, M.: Functors of Artin rings. *Trans.Amer.Math. Soc.* vol.130, (1968), pp.208-222.
- [Serre] Serre, J.P.: *Cours d'Arithmétique*. PUF, Paris, 1970.
- [Siv] Siveland, A.: *Matric Massey products and formal moduli*. Thesis. Institute of Mathematics, University of Oslo (1995)