

SYNCHROTRON RADIATION

Lenny Rivkin
Paul Scherrer Institute, Switzerland

ICTP SCHOOL ON SYNCHROTRON RADIATION AND APPLICATIONS
Trieste, Italy, May 2006

Useful books and references

- A. Hofmann, *The Physics of Synchrotron Radiation*
Cambridge University Press 2004
- H. Wiedemann, *Synchrotron Radiation*
Springer-Verlag Berlin Heidelberg 2003
- H. Wiedemann, *Particle Accelerator Physics I and II*
Springer Study Edition, 2003
- A. W. Chao, M. Tigner, *Handbook of Accelerator Physics and Engineering*, World Scientific 1999

CERN Accelerator School Proceedings

Synchrotron Radiation and Free Electron Lasers

Grenoble, France, 22 - 27 April 1996

(A. Hofmann's lectures on synchrotron radiation)

CERN Yellow Report 98-04

http://cas.web.cern.ch/cas/CAS_Proceedings-DB.html


Brunnen, Switzerland, 2 – 9 July 2003

CERN Yellow Report 2005-012

<http://cas.web.cern.ch/cas/BRUNNEN/lectures.html>

SYNCHROTRON RADIATION

Curved orbit of electrons in magnet field



Accelerated charge →


Electromagnetic radiation

Crab Nebula
6000 light years away




First light observed
1054 AD

GE Synchrotron
New York State



First light observed
1947

60 000 users world-wide





THEORETICAL UNDERSTANDING →

1873 Maxwell's equations

→ made evident that changing charge densities would result in electric fields that would radiate outward

1887 Heinrich Hertz demonstrated such waves:



..... this is of no use whatsoever !

Maxwell equations (poetry)

*War es ein Gott, der diese Zeichen schrieb
Die mit geheimnisvoll verborg'nem Trieb
Die Kräfte der Natur um mich enthüllen
Und mir das Herz mit stiller Freude füllen.*


Ludwig Boltzman

*Was it a God whose inspiration
Led him to write these fine equations
Nature's fields to me he shows
And so my heart with pleasure glows.*


translated by John P. Blewett

Why do they radiate?


Charge at rest: Coulomb field, no radiation




Uniformly moving charge
does not radiate (but! Cerenkov!)



Accelerated charge



Bremsstrahlung or breaking radiation



1898 Liénard:

ELECTRIC AND
MAGNETIC FIELDS
PRODUCED BY A POINT
CHARGE MOVING ON AN
ARBITRARY PATH
(by means of retarded potentials)

L'Éclairage Électrique

REVUE HEBDOMADAIRE D'ÉLECTRICITÉ

DIRECTION SCIENTIFIQUE

A. CORNU, Professeur à l'École Polytechnique, Membre de l'Institut. — A. D'ARSONVAL, Professeur au Collège de France, Membre de l'Institut. — G. LIPPMANN, Professeur à la Sorbonne, Membre de l'Institut. — D. MONNIER, Professeur à l'École centrale des Arts et Manufactures. — H. POINCARÉ, Professeur à la Sorbonne, Membre de l'Institut. — A. POTIER, Professeur à l'École des Mines, Membre de l'Institut. — J. BLONDIN, Professeur agrégé de l'Université.

CHAMP ÉLECTRIQUE ET MAGNÉTIQUE

PRODUIT PAR UNE CHARGE ÉLECTRIQUE CONCENTRÉE EN UN POINT ET ANIMÉE D'UN MOUVEMENT QUELCONQUE

Admettons qu'une masse électrique en mouvement de densité ρ et de vitesse u en chaque point produit le même champ qu'un courant de conduction d'intensité $u\rho$. En conservant les notations d'un précédent article (1) nous obtiendrons pour déterminer le champ, les équations

$$\frac{1}{4\pi} \left(\frac{d\gamma}{dy} - \frac{d\beta}{dz} \right) = \rho u_x + \frac{df}{dt} \quad (1)$$

$$V^2 \left(\frac{dh}{dy} - \frac{dg}{dz} \right) = - \frac{1}{4\pi} \frac{dx}{dt} \quad (2)$$

avec les analogues déduites par permutation tournante et en outre les suivantes

$$\rho = \left(\frac{df}{dx} + \frac{dg}{dy} + \frac{dh}{dz} \right) \quad (3)$$

$$\frac{dx}{dx} + \frac{d\beta}{dy} + \frac{d\gamma}{dz} = 0. \quad (4)$$

De ce système d'équations on déduit facilement les relations

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) f = V^2 \frac{d\rho}{dx} + \frac{d}{dt} (\rho u_x) \quad (5)$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) \alpha = 4\pi V^2 \left[\frac{d}{dz} (\rho u_y) - \frac{d}{dy} (\rho u_z) \right] \quad (6)$$

(1) La théorie de Lorentz, *L'Éclairage Électrique*, t. XIV, p. 417. α, β, γ , sont les composantes de la force magnétique et f, g, h , celles du déplacement dans l'éther.

Soient maintenant quatre fonctions ψ, F, G, H définies par les conditions

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) \psi = - 4\pi V^2 \rho. \quad (7)$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) F = - 4\pi V^2 \rho u_x$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) G = - 4\pi \rho u_y$$

$$\left(V^2 \Delta - \frac{d^2}{dt^2} \right) H = - 4\pi V^2 \rho u_z \quad (8)$$

On satisfera aux conditions (5) et (6) en prenant

$$4\pi f = - \frac{d\psi}{dx} - \frac{1}{V^2} \frac{dF}{dt} \quad (9)$$

$$\alpha = \frac{dH}{dy} - \frac{dG}{dz}. \quad (10)$$

Quant aux équations (1) à (4), pour qu'elles soient satisfaites, il faudra que, en plus de (7) et (8), on ait la condition

$$\frac{d\psi}{dt} + \frac{dF}{dx} + \frac{dG}{dy} + \frac{dH}{dz} = 0. \quad (11)$$

Occupons-nous d'abord de l'équation (7). On sait que la solution la plus générale est la suivante :

$$\psi = \int \frac{\rho \left[x', y', z', t - \frac{r}{V} \right]}{r} d\omega \quad (12)$$

Fig. 1. First page of Liénard's 1898 paper.

Liénard-Wiechert potentials

$$\varphi(\mathbf{t}) = \frac{1}{4\pi\epsilon_0} \frac{q}{[\mathbf{r}(1 - \mathbf{n} \cdot \vec{\beta})]_{ret}}$$
$$\vec{\mathbf{A}}(\mathbf{t}) = \frac{q}{4\pi\epsilon_0 c^2} \left[\frac{\vec{\mathbf{v}}}{\mathbf{r}(1 - \mathbf{n} \cdot \vec{\beta})} \right]_{ret}$$

and the electromagnetic fields:

$$\nabla \cdot \vec{\mathbf{A}} + \frac{1}{c^2} \frac{\partial \varphi}{\partial t} = 0 \quad (\text{Lorentz gauge})$$

$$\vec{\mathbf{B}} = \nabla \times \vec{\mathbf{A}}$$

$$\vec{\mathbf{E}} = -\nabla \varphi - \frac{\partial \vec{\mathbf{A}}}{\partial t}$$


Fields of a moving charge

$$\vec{\mathbf{E}}(t) = \frac{q}{4\pi\epsilon_0} \left[\frac{\vec{\mathbf{n}} - \vec{\beta}}{(1 - \vec{\mathbf{n}} \cdot \vec{\beta})^3 \gamma^2} \cdot \frac{1}{r^2} \right]_{ret} +$$


$$\frac{q}{4\pi\epsilon_0 c} \left[\frac{\vec{\mathbf{n}} \times [(\vec{\mathbf{n}} - \vec{\beta}) \times \vec{\beta}]}{(1 - \vec{\mathbf{n}} \cdot \vec{\beta})^3 \gamma^2} \cdot \frac{1}{r} \right]_{ret}$$

$$\vec{\mathbf{B}}(t) = \frac{1}{c} [\vec{\mathbf{n}} \times \vec{\mathbf{E}}]$$


Transverse acceleration



Radiation field quickly separates itself from the Coulomb field




Longitudinal acceleration




**Radiation field cannot
separate itself from the
Coulomb field**

Moving Source of Waves



Time compression

Electron with velocity β emits a wave with period T_{emit} while the observer sees a different period T_{obs} because the electron was moving towards the observer



$$T_{\text{obs}} = (1 - \mathbf{n} \cdot \boldsymbol{\beta}) T_{\text{emit}}$$

The wavelength is shortened by the same factor

$$\lambda_{\text{obs}} = (1 - \beta \cos \theta) \lambda_{\text{emit}}$$


in ultra-relativistic case, looking along a tangent to the trajectory

$$\lambda_{\text{obs}} = \frac{1}{2\gamma^2} \lambda_{\text{emit}}$$



since

$$1 - \beta = \frac{1 - \beta^2}{1 + \beta} \approx \frac{1}{2\gamma^2}$$

Radiation is emitted into a narrow cone



$$\theta = \frac{1}{\gamma} \cdot \theta_e$$



$$v \ll c$$

$$v \approx c$$

Synchrotron radiation power

Power emitted is proportional to:

$$P \propto E^2 B^2$$

$$P_{\gamma} = \frac{c C_{\gamma}}{2\pi} \cdot \frac{E^4}{\rho^2}$$

$$C_{\gamma} = \frac{4\pi}{3} \frac{r_e}{(m_e c^2)^3} = 8.858 \cdot 10^{-5} \left[\frac{\text{m}}{\text{GeV}^3} \right]$$

The power is all too real!




Fig. 12. Damaged X-ray ring front end gate valve. The power incident on the valve was approximately 1 kW for a duration estimated to 2–10 min and drilled a hole through the valve plate.

Synchrotron radiation power

Power emitted is proportional to:

$$P \propto E^2 B^2$$

$$P_\gamma = \frac{c C_\gamma \cdot E^4}{2\pi \rho^2}$$

$$P_\gamma = \frac{2}{3} \alpha \hbar c^2 \cdot \frac{\gamma^4}{\rho^2}$$

$$C_\gamma = \frac{4\pi}{3} \frac{r_e}{(m_e c^2)^3} = 8.858 \cdot 10^{-5} \left[\frac{\text{m}}{\text{GeV}^3} \right]$$

$$\alpha = \frac{1}{137}$$

Energy loss per turn:

$$U_0 = C_\gamma \cdot \frac{E^4}{\rho}$$


$$\hbar c = 197 \text{ Mev} \cdot \text{fm}$$

$$U_0 = \frac{4\pi}{3} \alpha \hbar c \frac{\gamma^4}{\rho}$$

Typical frequency of synchrotron light

Due to extreme collimation of light observer sees only a small portion of electron trajectory (**a few mm**)

$$l \sim \frac{2\rho}{\gamma}$$




Pulse length:
difference in times it
takes an electron
and a photon to
cover this distance

$$\Delta t \sim \frac{l}{\beta c} - \frac{l}{c} = \frac{l}{\beta c} (1 - \beta)$$

$$\omega \sim \frac{1}{\Delta t} \sim \gamma^3 \omega_0$$

$$\Delta t \sim \frac{2\rho}{\gamma c} \cdot \frac{1}{2\gamma^2}$$




Spectrum of synchrotron radiation

- Synchrotron light comes in a series of flashes every T_0 (revolution period)

- the spectrum consists of harmonics of

$$\omega_0 = \frac{1}{T_0}$$



- flashes are extremely short: harmonics reach up to very high frequencies

$$\omega_{\text{typ}} \cong \gamma^3 \omega_0$$

- At high frequencies the individual harmonics overlap

$$\omega_0 \sim 1 \text{ MHz}$$

$$\gamma \sim 4000$$

$$\omega_{\text{typ}} \sim 10^{16} \text{ Hz !}$$


continuous spectrum !

$$\frac{dP}{d\omega} = \frac{P_{\text{tot}}}{\omega_c} S\left(\frac{\omega}{\omega_c}\right)$$


$$S(x) = \frac{9\sqrt{3}}{8\pi} x \int_x^\infty K_{5/3}(x') dx' \quad \int_0^\infty S(x') dx' = 1$$

$$P_{\text{tot}} = \frac{2}{3} \hbar c^2 \alpha \frac{\gamma^4}{\rho^2}$$

$$\omega_c = \frac{3c\gamma^3}{2\rho}$$




Synchrotron radiation flux for different LEP energies




$$\text{Flux} \left[\frac{\text{photons}}{\text{s} \cdot \text{mrad} \cdot 0.1\% \text{BW}} \right] = 2.46 \cdot 10^{13} E[\text{GeV}] I[\text{A}] G_1(x)$$

Polarisation

Synchrotron radiation observed in the plane of the particle orbit is horizontally polarized, i.e. the electric field vector is horizontal



Observed out of the horizontal plane, the radiation is elliptically polarized




Polarisation: spectral distribution

$$\frac{dP}{d\omega} = \frac{P_{tot}}{\omega_c} S(x) = \frac{P_{tot}}{\omega_c} [S_\sigma(x) + S_\pi(x)]$$


$$S_\sigma = \frac{7}{8} S$$

$$S_\pi = \frac{1}{8} S$$




Angular divergence of radiation

•at the critical frequency




•well below



$$\omega = 0.2 \omega_c$$

•well above

$$\omega = 2 \omega_c$$



Angular divergence of radiation

The rms opening angle R'

- at the critical frequency:

$$\omega = \omega_c \quad R' \approx \frac{0.54}{\gamma}$$

- well below

$$\omega \ll \omega_c \quad R' \approx \frac{1}{\gamma} \left(\frac{\omega_c}{\omega} \right)^{1/3} \approx 0.4 \left(\frac{\lambda}{\rho} \right)^{1/3}$$

independent of γ !

- well above

$$\omega \gg \omega_c \quad R' \approx \frac{0.6}{\gamma} \left(\frac{\omega_c}{\omega} \right)^{1/2}$$

END