



Sixth International Conference on Perspectives in Hadronic Physics

12 - 16 May 2008

Medium Modifications from $4\text{He}(e,e'p)3\text{H}$

Simona Malace (Postdoc)
University of South Carolina
USA

Michael Paolone (Ph.D. Student)
University of South Carolina
USA

Steffen Strauch
University of South Carolina
USA

Medium Modifications from ${}^4\text{He}(e,e'p){}^3\text{H}$

- Nucleons in the Nuclear Medium and in-medium electromagnetic form factors
- Preliminary results from JLab experiment E03-104 (Hall A Collaboration)
 - ▶ Polarization transfer
 - ▶ Induced polarization
- Momentum dependence of bound nucleon wave function

Simona Malace (Postdoc), Michael Paolone (Ph.D. Student), Steffen Strauch
University of South Carolina


Sixth International Conference on Perspectives in Hadronic Physics
ICTP, Trieste, Italy – May 12-16, 2008

Nucleons in the Nuclear Medium

- **Conventional Nuclear Physics:**
 - ▶ Nuclei are effectively and well described as point-like protons and neutrons (+ form factor) and interaction through effective forces (meson exchange).
 - ▶ Medium effects arise through non-nucleonic degrees of freedom.
 - ▶ Are free nucleons and mesons, under every circumstance, the best quasi-particle to chose?
- **Nucleon Medium Modifications:**
 - ▶ Nucleons and mesons are not the fundamental entities in QCD.
 - ▶ Medium effects arise through changes of fundamental properties of the nucleon.
 - ▶ Do nucleons change their quark-gluon structure in the nuclear medium?

The EMC Effect

- The European Muon Collaboration used muon scattering to measure nuclear structure functions and observed a **depletion of the nuclear structure function** $F_2^A(x)$ in the valence-quark regime $0.3 \leq x \leq 0.8$.
- J. Smith and G. Miller: **chiral quark-soliton model of the nucleon**
Conventional nuclear physics does not explain EMC effect.



J.R. Smith and
G.A. Miller, Phys.
Rev. Lett. **91**,
212301 (2003)


- → **Nucleon structure is modified** in the nuclear medium
- Note: prelim. E03-103 ^4He data consistent with SLAC A=12 param.

Dave Gaskell, NuINT07, May 31 2007

y - Scaling Function

- y - scaling analysis of quasielastic scattering data
- Deviation of the **cross-section** from scattering from **free nucleons** scales to a function of a single variable y , the longitudinal momentum distribution.
- y-scaling property very sensitive to **change of nucleon radius**
- **Limits:** $Q^2 > 1 \text{ (GeV/c)}^2 : \Delta G_M < 3\%$

$$F(y) = \frac{\sigma(q, \omega)}{Z\sigma_{ep} + N\sigma_{en}} \cdot \frac{d\omega}{dy}$$




I. Sick, D. Day and J.S. McCarthy, Phys. Rev. Lett. **45**, 871 (1980);
Limit on radius from I. Sick, in: H. Klapdor (Ed.), Proc. Int. Conf. on Weak and Electromagnetic Interactions in Nuclei, Springer-Verlag, Berlin, 1986, p. 415

Coulomb Sum Rule

- **CSR:** Integral of the quasi-elastic electric response Response $R_L(q, \omega)$

$$S_L(q) = \frac{1}{Z} \int_{0+}^{\infty} \frac{R_L(q, \omega)}{\tilde{G}_E^2} d\omega \rightarrow 1$$

- **Experimental findings** controversial:
 - ▶ No quenching in the data observed [2]
 - ▶ Quenching of S_L is experimentally established [3]
 - ▶ Good agreement between theory and experiment for ${}^4\text{He}$ when using free-nucleon form factors [4]




- **Limits:** $Q^2 \leq 0.5 \text{ (GeV/c)}^2$:
 $\Delta G_E < 15\%$ or even $< 5\%$
- New data expected from JLab E05-110
 [Choi, Chen, and Meziani]


[1] I. Sick, Phys. Lett. B **157**, 13 (1985)
 [2] J. Jourdan, Nucl. Phys. A **603**, 117 (1996)
 [3] J. Morgenstern, Z.-E. Meziani, Phys. Lett. B **515**, 269 (2001)
 [4] J. Carlson, J. Jourdan, R. Schiavilla, and I. Sick, Phys. Lett. B **553**, 191 (2003)

Quark Meson Coupling Model (QMC)

- **Structure of the nucleon** described by valence quarks in a bag (Cloudy-bag model).
- **Nuclear system** described using effective scalar (σ) and vector (ω) meson fields.



intermediate baryon restricted to N or Δ





- Scalar and vector fields of nuclear matter couple directly to confined quarks.

→ Modification of **internal structure** of bound nucleon

D.H. Lu, A.W. Thomas, K. Tsushima, A.G. Williams, K. Saito, Phys. Lett. B **417**, 217 (1998)

D.H. Lu *et al.*, Phys. Rev. C **60**, 068201 (1999)


QMC: Bound Proton EM Form Factors



- Electromagnetic rms radii and magnetic moments of the bound proton are increased.
- At low Q^2 : **Charge form factor** much more sensitive to the nuclear medium than the **magnetic** ones.

D.H. Lu *et al.*, Phys. Rev. C **60**, 068201 (1999)

Chiral Quark Soliton Model (CQSM)



- Chiral-soliton model provides the quark and antiquark substructure of the proton, embedded in nuclear matter.
- Medium modifications:
 - ▶ significant for G_E , only moderate for G_M
 - ▶ no strong enhancement of the magnetic moment

CQSM: J.R. Smith and G.A. Miller, Phys. Rev. C **70**, 065205 (2004)

Other Models

- **Extended Skyrme Model**

U. Yakhshiev, U. Meißner, A. Wirzba, Eur. Phys. J. A **16**, 569 (2003)

- ▶ Model of the nucleon based on **Skyrme Lagrangian**
- ▶ Results for ${}^4\text{He}$ comparable to QMC, but differ in detail
- ▶ $(G_E/G_M)_{\text{medium}}/(G_E/G_M)_{\text{free}} \approx 1$ for $R = 1$ fm

- **Nambu–Jona-Lasinio model**

T. Horikawa, W. Bentz, Nucl. Phys. A **762**, 102 (2005)

- ▶ Nucleon as **quark-diquark** bound state + **nuclear matter** in the mean field approximation.
- ▶ Medium modifications: increase of the electric size in the medium
- ▶ **Medium modifications decrease with increasing Q^2** for both, spin and orbital form factors.

- **Generalized Parton Distributions**

S. Liuti, hep-ph/0608251, hep-ph/0601125

- ▶ Connection between the modifications induced by the nuclear medium of the nucleon form factors and of the deep inelastic structure functions, obtained using the concept of **generalized parton distributions**.

Medium-modified form factors are not an experimental observable. How can we test these predictions?

Strategy:

- Choose an **observable** with high sensitivity to nucleon structure while being at the same time least sensitive to conventional medium effects.
- Choose a dense yet simple nuclear **target**, which allows for microscopic calculations.
- Provide high-precision data to put Nuclear Physics models to **rigorous test**.

Polarization-Transfer Technique

- **Free** electron-nucleon scattering

$$\frac{G_E}{G_M} = -\frac{P'_x}{P'_z} \cdot \frac{(E_i + E_f)}{2m} \tan\left(\frac{\theta_e}{2}\right) \quad {}^1\text{H}(\vec{e}, e'\vec{p})$$

- **Bound** nucleons → evaluation within model
Reaction-mechanism effects predicted to be small and minimal for


- ▶ Quasielastic scattering
- ▶ Small missing momenta
- ▶ Symmetry about $\mathbf{p}_m = 0$

$$A(\vec{e}, e'\vec{p})B$$


R. Arnold, C. Carlson, and F. Gross, Phys. Rev. C **23**, 363 (1981); for reaction-mechanism effects, e.g., J.M. Laget, Nucl. Phys. A **579**, 333 (1994), J.J. Kelly, Phys. Rev. C **59**, 3256 (1999), A. Meucci, C. Guisti, and F.D. Pacati, Phys. Rev. C **66**, 034610 (2002).

Proton Recoil Polarization in ${}^4\text{He}(\vec{e}, e' \vec{p}){}^3\text{H}$

- **Kinematics:** low missing momentum, quasielastic scattering



- **Channel identification** by missing mass (Mike Paolone)



- Polarization-transfer ratio P'_x/P'_z : sensitive to G_E/G_M

$$R = \left(\frac{P'_x}{P'_z} \right)_{\text{bound}} / \left(\frac{P'_x}{P'_z} \right)_{\text{free}}$$

- Induced polarization P_y : sensitive to final-state interactions

Thomas Jefferson National Accelerator Facility




JLab in Newport News, VA



Hall A Counting House

E93-049 and E03-104 at Jefferson Lab Hall A


${}^4\text{He}(e, e'p){}^3\text{H}$ in quasielastic kinematics $Q^2 = 0.5 - 2.6 \text{ (GeV/c)}^2$




S. Strauch, *et al.*, Phys. Rev. Lett. **91**, 052301(2003);
JLab E03-104, R. Ent, R. Ransome, S. Strauch, P. Ulmer (spokespersons)

Polarization Measurement

Spin-dependent scattering




Focal-Plane Polarimeter



Observed angular distribution

$$\begin{aligned}
 I(\vartheta, \varphi) &= I_0(\vartheta) (1 + \epsilon_y \cos \varphi + \epsilon_x \sin \varphi) \\
 &= I_0(\vartheta) [1 + A_C (P_y \cos \varphi - P_x \sin \varphi)]
 \end{aligned}$$


Observed Angular Distribution



- Excellent control of systematic uncertainties for **polarization transfer**

- Instrumental asymmetries complicate the extraction of **induced polarization**
 - ▶ Detector misalignment
 - ▶ Detector inefficiencies
 - ▶ Tracking problems
- (Simona Malace)

Free Proton Form-Factor Ratio $\mu_p G_E/G_M$




- Preliminary results from E03-104 in good agreement with previous data.
- Small statistical uncertainties 0.7% from E03-104.
- Final data will have reduced systematic uncertainties.

^2H and $^4\text{He}(e,e'p)$ Polarization-Transfer Ratios

$$R = \left(\frac{P'_x}{P'_z} \right)_{\text{bound}} / \left(\frac{P'_x}{P'_z} \right)_{\text{free}}$$

- ^2H and ^1H polarization-transfer data are similar.
- ^4He data are significantly different than ^2H , ^1H data.
- Small effect for less dense nucleus, larger for denser.
- RDWIA and RMSGA models cannot describe ^4He data.




^2H Model: H. Arenövel; see: B. Hu *et al.*, Phys. Rev. C **73**, 064004 (2006).

RDWIA: J.M. Udias *et al.*, Phys. Rev. Lett. **83**, 5451 (1999).

Relativistic Multiple-Scattering Glauber Approximation (RMSGGA):



P. Lava *et al.*, Phys. Rev. C **71**, 014605 (2005), D. Debruyne *et al.*, Phys. Rev. C **62**, 024611 (2000).

${}^4\text{He}(e,e'p){}^3\text{H}$ - Polarization-Transfer Ratio




- Previous and preliminary high-statistics data from E03-104 are also low compared to RDWIA
- R^{RDWIA} reduced by 3% compared to R^{RPWIA} due to Enhancement of lower components (spinor distortions) in RDWIA
- Small sensitivity to
 - ▶ bound-state wave function
 - ▶ current operator
 - ▶ optical potential (not including charge exchange terms)

Role of MEC in ${}^4\text{He}(e,e'p){}^3\text{H}$



direct knockout contribution




seagull (contact) diagram with one-pion exchange

- The seagull diagram effects generally **small** and **visible only at high missing momenta**; MEC expected to give more significant effect in the induced polarization
 Relativistic mean-field calculation: A. Meucci, C. Giusti, and F.D. Pacati, *Phys. Rev. C* **66**, 034610 (2002)
- R is suppressed by about 4%** with respect to that obtained with one-body currents only
 R. Schiavilla, O. Benhar, A. Kievsky, L.E. Marcucci, and M. Viviani, *Phys. Rev. Lett.* **94**, 072303 (2005)

Polarization Transfer in ${}^4\text{He}(\vec{e}, e' \vec{p}){}^3\text{H}$

$$G(Q^2, \rho) = G(Q^2) \frac{G_{\text{QMC}}(Q^2, \rho)}{G_{\text{QMC}}(Q^2)}$$



- In-medium form factors: density-dependent form factors are evaluated at the local density $\rho(r)$.
- R is reduced by an additional 6% (QMC).
- Data effectively described by proton medium modified form factors

Interpretation of Polarization-Transfer Data


- Data consistent with:

- ▶ RDWIA
- ▶ Density-dependent **medium modified form factors.**

OR


- ▶ Free form factors
- ▶ MEC
- ▶ **Spin-dependent charge-exchange FSI** (not well constrained \Rightarrow need P_y from E03-104)

R. Schiavilla, O. Benhar, A. Kievsky, L.E. Marcucci, and M. Viviani, Phys. Rev. Lett. **94**, 072303 (2005)



- The modeling of final-state interactions can be tested by measuring the induced polarization, P_y .

Induced Polarization in ${}^4\text{He}(e, e' \vec{p})$




Note: Data are acceptance corrected; inner uncertainties are statistical only; full analysis of E03-104 will have reduced systematic uncertainties

- P_y is a measure of **final-state interactions (FSI)**
- P_y is insensitive to **in-medium form factors.**
- Observed final-state interaction **small** and with only **very weak Q^2 dependence**
- Results from RDWIA and Laget consistent with data
- Spin-dependent charge exchange terms not well constrained by N-N scattering and possibly overestimated

Bound Nucleon Wave Function

Pointlike Configurations (PLC)

- Smaller average interaction strength
- PLC suppressed in the bound state
- Contribution of PLCs exhibit a strong momentum dependence (arising from the reduction of the interaction strength)



Ciofi degli Atti *et al.* argue that **medium modifications should strongly depend on the nucleon momentum** (nucleon virtuality)


$$v = p^2 - m_N^2$$

$$= \left(M_A - \sqrt{(M_A - m_N + E)^2 + \mathbf{p}^2} \right)^2 - \mathbf{p}^2 - m_N^2$$

- At $v = 0$, modification should vanish.
- “Would be nice to study modification of the nucleon form factors as a function of the nucleon momentum.” [\[Mark Strikman\]](#)

C. Ciofi degli Atti, L.L. Frankfurt, L.P. Kaptari, M.I. Strikman, Phys. Rev. C **76**, 055206 (2007)
M.R. Frank, B.K. Jennings, G.A. Miller, Phys. Rev. C **54**, 920 (1996)

Proton Virtuality – Suppression of PLCs?



- Polarization-transfer double ratio shows (linear) **dependence on proton virtuality** with the trend of $R \approx 1$ for $p^2 = m^2_N$
- Excellent description with the **RDWIA + QMC** model.

Summary

- **Models predict change of the internal structure of bound nucleon**
- **Recoil-polarization in $^4\text{He}(e,e'p)^3\text{H}$**
 - ▶ Two polarization observables act together to constrain the interpretation of the data
 - **Polarization transfer**: sensitive to in-medium form factors
 - **Induced polarization**: sensitive to final-state interactions, not sensitive to in-medium form factors
- **Preliminary results**
 - ▶ Data effectively described **by in-medium electromagnetic form factors** or strong **charge-exchange FSI**
 - ▶ Induced polarization crucial to clarify role of FSI and new results from E03-104 will provide needed constraints