



The Abdus Salam
International Centre for Theoretical Physics



2139-22

**School on Synchrotron and Free-Electron-Laser Sources and their
Multidisciplinary Applications**

26 April - 7 May, 2010

**Surface, Interface, and Materials Studies Using Photoelectron Spectroscopy,
Diffraction, and Holography**

Charles S. Fadley
*Lawrence Berkeley National Laboratory
USA*

International Center for Theoretical Physics, Trieste, Italy
Fuggle-Fonda School on Synchrotron Radiation and Applications

LECTURES FOR 30 APRIL AND 3 MAY, 2010

***SURFACE, INTERFACE, AND MATERIALS STUDIES USING
PHOTOELECTRON SPECTROSCOPY,
DIFFRACTION, AND HOLOGRAPHY***

Lecturers:


Chuck Fadley, Department of Physics, University of California, Davis &
Materials Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, USA

*Introduction to surface and interface science, vuv/soft x-ray spectroscopies, photoelectron
spectroscopy/diffraction/holography, hard x-ray photoemission*

Andrea Goldoni, Elettra, Trieste, Italy

*Valence bands, dispersion, and Fermi surface mapping by photoemission,
angle-resolved photoemission, many-body effects,
complex materials, spin-resolved studies*

Chuck Fadley
Dept. of Physics, University of California Davis
Davis California
and
Lawrence Berkeley National Laboratory
Berkeley, California





Fugle-Fonda School on Synchrotron Radiation and Applications

International Center for Theoretical Physics

April-May, 2010

With acknowledgments to:



The photoelectric effect: Annalen der Physik ¹⁷, 146 (1905)

$$h\nu = E_{initial} - E_{final} = E_{binding} + E_{kinetic}$$

(Nobel Prize for it in 1921 – But no mention of it in his Nobel lecture)



*“I would like to tell
you how pleased I
am that you have
given up your
light-quantum
theory”*


von Laue to Einstein, 1907 letter



In his 1913 letter nominating Einstein for the membership of Prussian Academy of Science, Max Planck et al. wrote:

- “*In sum, one can say there is hardly one among the great problems in which modern physics is so rich to which Einstein has not made a remarkable contribution. That he may sometimes has missed the target in his speculations, as, for example, in his hypothesis of light quanta, cannot really be held too much against him, for it is not possible to introduce really new ideas even in the most exact sciences without sometimes taking a risk.*”

Typical experimental geometry for energy- and angle-resolved photoemission measurements



Outline

Surface, interface, and nanoscience—short introduction

Some surface concepts and techniques→photoemission

Synchrotron radiation: experimental aspects

Electronic structure—a brief review

**The basic synchrotron radiation techniques:
more experimental and theoretical details**

Valence-level photoemission

Core-level photoemission

**Photoemission with high ambient pressure
around the sample**

***SURFACE, INTERFACE, AND MATERIALS STUDIES USING
PHOTOELECTRON SPECTROSCOPY, DIFFRACTION, HOLOGRAPHY, AND MICROSCOPY;
(X-RAY FLUORESCENCE HOLOGRAPHY)***

Chuck Fadley

Department of Physics, University of California-Davis, Davis, CA, &
Materials Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, CA

---OUTLINE OF FADLEY LECTURES FOR 16 APRIL THROUGH 18 APRIL, 2008---

(With complementary coverage of related/additional material by Andrea Goldoni, Elettra)

References below are to papers handed out as in format: Paper [No.], section no. or page nos, or to other lectures in this School as appropriate. See also original literature references referred to directly on many slides.

•INTRODUCTION:

--Surface and interface phenomena: (*"Modern Techniques of Surface Science"*, D.P. Woodruff and T.A. Delchar, (Cambridge Univ. Press, 1994), 2nd Edition,; *"Surface Physics"*, A. Zangwill (Cambridge Univ. Press, 1990))

What are they? Why study them?

Applications in technology: semiconductor ICs, spintronics, et al.

Nanoscience/nanotechnology

Ultrahigh vacuum

Some basic concepts and characterization techniques: TEM, LEED and STM

Electron escape and surface sensitivity

Typical experimental systems

--Synchrotron radiation experiments: (*Other lectures in this School*)

Basic considerations—brief review

X-ray emission and nomenclature

Synchrotron radiation

X-ray interactions with matter and basic techniques

Photoelectron spectroscopy =

photoemission(PS, PES)

X-ray absorption spectroscopy (XAS,

NEXAFS(=XANES) + EXAFS =

XAES (*Other lectures in this School*)

X-ray emission/x-ray fluorescence spectroscopy (XES, XFS)

and resonant inelastic x-ray scattering (RIXS)

X-ray scattering and diffraction (XRD, *other lectures in this School*)

X-ray optical measurements (refraction, reflection and penetration depth,

Standing waves,...)

• **ELECTRONIC STRUCTURE:** (*Zangwill book, Paper [1], Chap.III*):

Basics of electronic structure and bonding
Hartree Fock Method, Koopmans Theorem and corrections to it
The exchange interaction and magnetism
Atomic orbitals, spin-orbit splitting
Molecular orbitals
Electrons in solids, bands

• **THE BASIC SR SPECTROSCOPIES—MORE EXPERIMENTAL AND THEORETICAL DETAILS:**

(*Paper [1], Chaps.I and III, Paper [15]*)

Photoelectron spectroscopy (PES, PS, XPS)
Auger electron spectroscopy (AES)
X-ray absorption spectroscopy (XAS, NEXAFS, XANES)
X-ray emission and resonant inelastic scattering (XES, RIXS)
Instrumentation for PES
Spectrometers and detectors
Electron spin detection
Measuring electron kinetic and binding energies:
Work function, inner potential
Sample charging

•CORE-LEVEL SPECTROSCOPY (PART 1):

--Core intensities (the 3-step model) and quantitative surface analysis:

(Papers [1], Chap. VI, Paper [2], 1-4, Papers [15] and [16])

Quantitative formulas for surface analysis

Surface sensitivity enhancement at grazing emission

--Differential photoelectric cross sections and selection rules

Basic forms and tabulations

Cooper minima

Resonant photoemission:

Intraatomic single atom resonant photoemission (RPE, SARPE)--

Well known

Interatomic multi-atom resonant photoemission (MARPE)--

a new effect in molecules, solids (*Paper [8]*)

Non-dipole effects at higher energies

--Inelastic attenuation length tabulations and estimates

--Elastic scattering effects in surface analysis

--Electron refraction in escape from surface

- **PHOTOELECTRON DIFFRACTION (CORE LEVELS):**

(Papers [1], D; [2], 5; [3]-[5], [15], [16])

--Basic diffraction and measurement process: scanned-angle and scanned-energy

--Energy dependence of scattering:

 Forward-dominated at high energies

 Back and forward at low energies

--Basic theory:

 Scattering factors: plane-wave and spherical-wave

 Vibrational effects and Debye-Waller factors

--Determination of structures from:

 Forward scattering peaks—adsorbed molecules

 More complex diffraction patterns

 (incl. full-solid -angle data and R-factor analysis)

Analysis via single-scattering and multiple scattering theory--review of

theoretical approaches and computer exercises for those

interested (*Paper [9] plus program EDAC discussed in lecture and exercises*)

--Fingerprint diffraction patterns

--Some example applications: adsorbates, clean surface core-level shifts, epitaxial overlayers, Moiré structures, time-dependent surface reactions

--Fourier transforms of scanned-energy data: path-length differences

- **PHOTOELECTRON HOLOGRAPHY:**

(Papers [3], 5.4; [4], 5.3; [5]; [6]; [7];[11]; [15])

--Basic process of hologram formation and image reconstruction:

 ~a Fourier-like transform of several types

--Applications in single-energy and multiple-energy form to
adsorbates and multilayer substrates

--Comparison of methods, including new approaches

- **VALENCE-LEVEL SPECTROSCOPY: (Paper [15], and Goldoni lectures)**

- The low-energy (UPS) limit: (*Goldoni lectures*)
 - Selection rules on wave vector
 - Band-structure mapping
 - Fermi-surface mapping
- Vibrational/phonon effects: UPS↔XPS limits (*Paper [2], [6], [14], [15], [16]*)
- The high-energy (XPS) limit: (*Papers [2], [6], [15], [16]*)
 - Density-of-states measurements
- Hard x-ray photoemission in the 5-15 keV range: a new direction (*Papers [14], [16]*)

- **CORE-LEVEL SPECTROSCOPY (PART 2):**

- X-ray optical effects: resonant and non-resonant, standing waves (*Papers [12], [13], [15]*,
 - Probing buried interfaces with soft x-ray standing waves (*Paper [12]*)
 - Chemical shifts in core binding energies (*Paper [1], Chap. IV*)
 - Potential model*
 - Equivalent-core approx. and relationship to thermochemical energies
 - Multiplet splittings & spin-polarized spectra (*Paper [1], Chap. V, A-D*)
 - Spin-polarized photoelectron diffraction and holography
 - Spin polarization via spin-orbit-split levels excited with circular polarized Radiation—the Fano effect
 - Magnetic circular dichroism in core photoemission
 - Non-magnetic circular dichroism in core photoemission
 - (circular dichroism in angular distributions--CDAD)
 - Shake-up/shake-off and Sudden Approx. sum rules
 - Final-state screening and relaxation effects, satellites (*Paper [1], Chap. V, A-D*)
 - Vibrational effects in spectra (*Paper [1], Chap. V, E*)

- **PHOTOEMISSION WITH A HIGH AMBIENT PRESSURE AT THE SAMPLE::**

General references on various aspects of photoelectron spectroscopy, diffraction, holography (available at School website):

Paper [1] "Basic Concepts of X-ray Photoelectron Spectroscopy", C.S.F, in Electron Spectroscopy, Theory, Techniques, and Applications, Brundle and Baker, Eds. (Pergamon Press, 1978) Vol. II, Ch. 1.

Paper [2] "Angle-Resolved X-ray Photoelectron Spectroscopy", C.S.F., Progress in Surface Science 16, 275 (1984).

Paper [3] "The Study of Surface Structures by Photoelectron Diffraction and Auger Electron Diffraction", C.S.F., in Synchrotron Radiation Research: Advances in Surface and Interface Science, Bachrach, Ed. (Plenum,1992)

Paper [4] "Photoelectron Diffraction: New Dimensions in Space, Time, and Spin", C.S. Fadley, M.A. Van Hove, Z. Hussain, and A.P. Kaduwela, J. Electron Spectrosc. 75, 273, (1995).

Paper [5] "Diffraction and Holography with Photoelectrons and Fluorescent X-Rays", C. S. Fadley et al., Progress in Surface Science 54, 341 (1997).

Paper [6] "Atomic Holography with Electrons and X-rays", P.M. Len, C.S. Fadley, and G. Materlik, invited paper appearing in X-ray and Inner-Shell Processes: 17th International Conference, R.L. Johnson, H. Schmidt-Böcking, and B.F. Sonntag, Eds., American Institute of Physics Conference Proceedings, No. 389 (AIP, New York, 1997) pp. 295-319.

Paper [7] "Theoretical Aspects of Electron Emission Holography", L. Fonda, Phys. Stat. Sol. (b) 188, 599 (1995). (Theoretical study by founder of this school.)

Paper [8] "Multi-Atom Resonant Photoemission", A.W. Kay, F.J. Garcia de Abajo, S.-H. Yang, E. Arenholz, B.S. Mun, N. Mannella, Z. Hussain, M.A. Van Hove, and C.S. Fadley, Physical Review B 63, 115119 (2001).

Paper [9] "Multiple Scattering of Electrons in Solids and Molecules: a Novel Cluster-Model Approach", F. J. Garcia de Abajo, C.S. Fadley, and M.A. Van Hove, Physical Review B63, 075404 (2001). (Paper describing the new "EDAC" multiple scattering program available for online usage at <http://maxwell.optica.csic.es/software/edac/a.html> in course tutorials and for anyone wishing to try it at home. See also downloadable "MSCD" program at <http://electron.lbl.gov/~mscd/>.)

Paper [10] "Fermi Surface Mapping by Angle-Resolved Photoemission", J. Osterwalder, Surface Review and Letters 4, 391 (1997). (Covered in greater detail in Osterwalder lectures.)

Paper [11] "Photoelectron and X-ray Holography by Contrast: Enhancing Image Quality and Dimensionality", C.S. Fadley, M.A. Van Hove, A. Kaduwela, S. Omori, L. Zhao, and S. Marchesini, J. Phys. Cond. Mat. 13, 10517 (2001).

Paper [12] "Probing Multilayer Spintronic Structures with Photoelectron and X-Ray Emission Spectroscopies Excited by X-Ray Standing Waves", S.-H. Yang, B.C. Sell, and C. S. Fadley, *J. Appl. Phys.* **103**, 07C519 (2008)

Paper [13] "X-ray Optics, Standing Waves, and Interatomic Effects in Photoemission and X-ray Emission", C. S. Fadley, S.-H. Yang, B. S. Mun, J. Garcia de Abajo, invited Chapter in the book "Solid-State Photoemission and Related Methods: Theory and Experiment", W. Schattke and M.A. Van Hove, Editors, (Wiley-VCH Verlag, Berlin GmbH, 2003), ISBN: 3527403345, 38 pp., 17 figs.

Paper [14] "X-Ray Photoelectron Spectroscopy and Diffraction in The Hard X-Ray Regime: Fundamental Considerations and Future Possibilities", C. S. Fadley, *Nuclear Instruments and Methods A* **547**, 24-41 (2005), special issue edited by J. Zegenhagen and C. Kunz.

Paper [15] "Atomic-Level Characterization of Materials with Core- and Valence-Level Photoemission: Basic Phenomena and Future Directions", C.S. Fadley, *Surf. Interface Anal.* **2008**, **40**, 1579–1605.

Paper [16] "X-ray photoelectron spectroscopy: Progress and perspectives"
C.S. Fadley, *Journal of Electron Spectroscopy and Related Phenomena* **178–179** (2010) 2–32

Key Reference [17] "X-ray Data Booklet", Center for X-Ray Optics and the Advanced Light Source, LBNL, January, 2001, available online at: <http://xdb.lbl.gov/>

Additional very useful websites:

X-ray optical calculations: reflectivities, penetration depths for a variety of mirror/surface geometries—

http://www-cxro.lbl.gov/optical_constants/

General properties of the elements and their compounds: <http://www.webelements.com>

Calculation of photoelectron diffraction with program EDAC:



<http://maxwell.optica.csic.es/software/edac/a.html>

[17]

X-RAY DATA BOOKLET
Center for X-ray Optics and Advanced Light Source
Lawrence Berkeley National Laboratory

<http://xdb.lbl.gov/>

- [Introduction](#)
- [X-Ray Properties of Elements](#)
- [Electron Binding Energies](#)
- [X-Ray Energy Emission Energies](#)
- [Fluorescence Yields for K and L Shells](#)
- [Principal Auger Electron Energies](#)
- [Subshell Photoionization Cross-Sections](#)
- [Mass Absorption Coefficients](#)
- [Atomic Scattering Factors](#)
- [Energy Levels of Few Electron Ions](#)
- [Periodic Table of X-Ray Properties](#)
- [Synchrotron Radiation](#)
- [Characteristics of Synchrotron Radiation](#)
- [History of X-rays and Synchrotron Radiation](#)
- [Synchrotron Facilities](#)
- [Scattering Processes](#)
- [Scattering of X-rays from Electrons and Atoms](#)
- [Low-Energy Electron Ranges in Matter](#)
- [Optics and Detectors](#)
- [Crystal and Multilayer Elements](#)
- [Specular Reflectivities for Grazing-Incidence Mirrors](#)
- [Gratings and Monochromators](#)
- [Zone Plates](#)
- [X-Ray Detectors](#)
- [Miscellaneous](#)
- [Physical Constants](#)
- [Physical Properties of the Elements](#)
- [Electromagnetic Relations](#)
- [Radioactivity and Radiation Protection](#)
- [Useful Formulas](#)



| | | | | | | | | | | | |
|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|------------------------|-------------------------|-------------------------|
| H¹ | | | | | | | | | | He² | |
| 1s | | | | | | | | | | 1s ² | |
| Li³ | Be⁴ | | | | | | | | | | |
| 2s | 2s ² | | | | | | | | | | |
| Na¹¹ | Mg¹² | | | | | | | | | | |
| 3s | 3s ² | | | | | | | | | | |
| K¹⁹ | Ca²⁰ | Sc²¹ | Ti²² | V²³ | Cr²⁴ | Mn²⁵ | Fe²⁶ | Co²⁷ | Ni²⁸ | Cu²⁹ | Zn³⁰ |
| 4s | 4s ² | 3d | 3d ² | 3d ³ | 3d ⁵ | 3d ⁵ | 3d ⁶ | 3d ⁷ | 3d ⁸ | 3d ¹⁰ | 3d ¹⁰ |
| | | 4s ² | 4s ² | 4s ² | 4s | 4s ² | 4s ² | 4s ² | 4s ² | 4s | 4s ² |
| Rb³⁷ | Sr³⁸ | Y³⁹ | Zr⁴⁰ | Nb⁴¹ | Mo⁴² | Tc⁴³ | Ru⁴⁴ | Rh⁴⁵ | Pd⁴⁶ | Ag⁴⁷ | Cd⁴⁸ |
| 5s | 5s ² | 4d | 4d ² | 4d ⁴ | 4d ⁵ | 4d ⁶ | 4d ⁷ | 4d ⁸ | 4d ¹⁰ | 4d ¹⁰ | 4d ¹⁰ |
| | | 5s ² | 5s ² | 5s | 5s | 5s | 5s | 5s | 5s | 5s ² | 5s ² |
| Cs⁵⁵ | Ba⁵⁶ | La⁵⁷ | Hf⁷² | Ta⁷³ | W⁷⁴ | Re⁷⁵ | Os⁷⁶ | Ir⁷⁷ | Pt⁷⁸ | Au⁷⁹ | Hg⁸⁰ |
| 6s | 6s ² | 5d | 5d ² | 5d ³ | 5d ⁴ | 5d ⁵ | 5d ⁶ | 5d ⁹ | 5d ⁹ | 5d ¹⁰ | 5d ¹⁰ |
| | | 6s ² | — | 6s | 6s | 6s ² |
| Fr⁸⁷ | Ra⁸⁸ | Ac⁸⁹ | Ce⁵⁸ | Pr⁵⁹ | Nd⁶⁰ | Pm⁶¹ | Sm⁶² | Eu⁶³ | Gd⁶⁴ | Tb⁶⁵ | Dy⁶⁶ |
| 7s | 7s ² | 6d | 4f ² | 4f ³ | 4f ⁴ | 4f ⁵ | 4f ⁶ | 4f ⁷ | 4f ⁷ | 4f ⁸ | 4f ¹⁰ |
| | | 7s ² | 6s ² | 5d | 5d | 6s ² |
| Th⁹⁰ | Pa⁹¹ | U⁹² | Np⁹³ | Pu⁹⁴ | Am⁹⁵ | Cm⁹⁶ | Bk⁹⁷ | Cf⁹⁸ | Es⁹⁹ | Fm¹⁰⁰ | Md¹⁰¹ |
| — | 5f ² | 5f ³ | 5f ⁵ | 5f ⁶ | 5f ⁷ | 5f ⁷ | 5f ⁷ | 6d | 4f ¹¹ | 4f ¹² | 4f ¹³ |
| 6d ² | 6d | 6d | 7s ² | 6s ² | 6s ² | 6s ² |
| 7s ² | 6s ² | 6s ² | 6s ² |

Periodic Table, with the Outer Electron Configurations of Neutral Atoms in Their Ground States

The notation used to describe the electronic configuration of atoms and ions is discussed in all textbooks of introductory atomic physics. The letters *s*, *p*, *d*, . . . signify electrons having orbital angular momentum 0, 1, 2, . . . in units \hbar ; the number to the left of the letter denotes the principal quantum number of one orbit, and the superscript to the right denotes the number of electrons in the orbit.

Table 3 Crystal structures of the elements

The data given are at room temperature for the most common form, or at the stated temperature in deg K. For further descriptions of the elements see Wyckoff, Vol. 1, Chap. 2. Structures labeled complex are described there.

Table 4 Density and atomic concentration

The data are given at atmospheric pressure and room temperature, or at the stated temperature in deg K. (Crystal modifications as for Table 3.)

Table 1 Debye temperature and thermal conductivity^a

| Li | Be | | | | | | | | | | | | | B | C | N | O | F | Ne |
|------|------|---|------|------|------|------|------|------|------|------|------|------|-----------------|------|------|------|----|----|----|
| 344 | 1440 | | | | | | | | | | | | | 2230 | | | | 75 | |
| 0.85 | 2.00 | | | | | | | | | | | | | 0.27 | 1.29 | | | | |
| Na | Mg | | | | | | | | | | | | | Al | Si | P | S | Cl | Ar |
| 158 | 400 | Low temperature limit of θ , in Kelvin | | | | | | | | | | | | 428 | 645 | | | 92 | |
| 1.41 | 1.56 | Thermal conductivity at 300 K, in $\text{W cm}^{-1}\text{K}^{-1}$ | | | | | | | | | | | | 2.37 | 1.48 | | | | |
| K | Ca | Sc | Ti | V | Cr | Mn | Fe | Co | Ni | Cu | Zn | Ga | Ge | As | Se | Br | Kr | | |
| 91 | 230 | 360. | 420 | 380 | 630 | 410 | 470 | 445 | 450 | 343 | 327 | 320 | 374 | 282 | 90 | | 72 | | |
| 1.02 | | 0.16 | 0.22 | 0.31 | 0.94 | 0.08 | 0.80 | 1.00 | 0.91 | 4.01 | 1.16 | 0.41 | 0.60 | 0.50 | 0.02 | | | | |
| Rb | Sr | Y | Zr | Nb | Mo | Tc | Ru | Rh | Pd | Ag | Cd | In | Sn _w | Sb | Te | I | Xe | | |
| 56 | 147 | 280 | 291 | 275 | 450 | | 600 | 480 | 274 | 225 | 209 | 108 | 200 | 211 | 153 | | 64 | | |
| 0.58 | | 0.17 | 0.23 | 0.54 | 1.38 | 0.51 | 1.17 | 1.50 | 0.72 | 4.29 | 0.97 | 0.82 | 0.67 | 0.24 | 0.02 | | | | |
| Cs | Ba | La β | Hf | Ta | W | Re | Os | Ir | Pt | Au | Hg | Tl | Pb | Bi | Po | At | Rn | | |
| 38 | 110 | 142 | 252 | 240 | 400 | 430 | 500 | 420 | 240 | 165 | 71.9 | 78.5 | 105 | 119 | | | | | |
| 0.36 | | 0.14 | 0.23 | 0.58 | 1.74 | 0.48 | 0.88 | 1.47 | 0.72 | 3.17 | | 0.46 | 0.35 | 0.08 | | | | | |
| Fr | Ra | Ac | Ce | Pr | Nd | Pm | Sm | Eu | Gd | Tb | Dy | Ho | Er | Tm | Yb | Lu | | | |
| | | | | | | | | | 200 | | 210 | | | | 120 | 210 | | | |
| | | | 0.11 | 0.12 | 0.16 | | 0.13 | | 0.11 | 0.11 | 0.11 | 0.16 | 0.14 | 0.17 | 0.35 | 0.16 | | | |
| | | | Th | Pa | U | Np | Pu | Am | Cm | Bk | Cf | Es | Fm | Md | No | Lr | | | |
| | | | 163 | | 207 | | 0.06 | 0.07 | | | | | | | | | | | |
| | | | 0.54 | | 0.28 | | | | | | | | | | | | | | |

^aMost of the θ values were supplied by N. Pearlman; references are given the A.I.P. Handbook, 3rd ed; the thermal conductivity values are from R. W. Powell and Y. S. Touloukian, Science 181, 999 (1973).

Outline

→ **Surface, interface, and nanoscience—short introduction**

Some surface concepts and techniques→photoemission

Synchrotron radiation: experimental aspects

Electronic structure—a brief review

**The basic synchrotron radiation techniques:
more experimental and theoretical details**

Valence-level photoemission

Core-level photoemission

**Photoemission with high ambient pressure
around the sample**

- Why surfaces, interfaces, structures at the nanometer scale?

$1 \text{ nm} = 10 \text{ \AA} = 0.001 \text{ micron}$

Cube of 1 nm sides has 75% of its atoms on the surface

Many areas of science/technology

Nobel Prizes in Physics and Chemistry--2007

From Spinwaves to Giant Magnetoresistance (GMR) and Beyond



Peter Grünberg held his Nobel Lecture on 8 December 2007, at Aula Magna, Stockholm University. He was introduced by Professor Per Carlson, Chairman of the Nobel Committee for Physics.

The Origin, the Development and the Future of Spintronics



Albert Fert delivered his Nobel Lecture on 8 December 2007, at Aula Magna, Stockholm University. He was introduced by Professor Per Carlson, Chairman of the Nobel Committee for Physics.

Reactions at Solid Surfaces: From Atoms to Complexity




Gerhard Ertl delivered his Nobel Lecture on 8 December 2007, at Aula Magna, Stockholm University, where he was introduced by Professor Gunnar von Heijne, Chairman of the Nobel Committee for Chemistry.



The Nobel Prize in Physics 1921

"for his services to Theoretical Physics, and especially for his discovery of the law of the photoelectric effect"



Albert Einstein

**Photoelectric effect→
Photoemission or
Photoelectron spectroscopy
(PS, PES)**



The Nobel Prize in Physics 1937

"for their experimental discovery of the diffraction of electrons by crystals"



Clinton Joseph Davisson




George Paget Thomson

**Low energy
electron
diffraction
(LEED)**



The Nobel Prize in Physics 1981

"for his contribution to the development of high-resolution electron spectroscopy"



Kai M. Siegbahn

**X-ray photoelectron
spectroscopy
(XPS) or
Electron spectroscopy
for chemical analysis
(ESCA)**



The Nobel Prize in Physics 1986

"for his fundamental work in electron optics, and for the design of the first electron microscope"



Ernst Ruska



Gerd Binnig




Heinrich Rohrer

Electron and Scanning tunneling microscopy (STM)


**Scientific and technological areas involving
surface/interface/nano science:**

- Integrated circuits—higher speed, higher density




And the Shrink Goes On...

.35 μ Process Technology



intel.


.25 μ Process Technology



Now 0.065 μ = 65 nm = 650 Å ('05)
 \rightarrow 45 nm = 450 Å ('07)

**~few atomic layers—
currently
15 Å SiO₂**

**Some
serious
challenges**



Cross section of a MOS transistor. Electron tunneling through the gate oxide (left inset) and high-concentration dopant interactions (right inset) are posing fundamental limitations to continuing historical transistor scaling trends.

High-k + Metal Gate Transistors

Metal Gate


- Increases the gate field effect

High-k Dielectric



- Increases the gate field effect
- Allows use of thicker dielectric layer to reduce gate leakage

HK + MG Combined

- Drive current increased >20% (>20% higher performance)
- Or source-drain leakage reduced >5x
- Gate oxide leakage reduced >10x



~1 %




World's Smallest Transistor

**IBM
Science
2001**


**Scientific and technological areas related to
surface/interface/nano science:**

- Integrated circuits—higher speed, higher density
- Magnetic storage and circuits—higher density, magnetic logic

“Moore’s Law” for magnetic storage




How far can we go?



The number of atoms used to store one bit of information with different forms of magnetic or optical storage has reduced over the years. The blue region indicates the superparamagnetic regime, below which thermal fluctuations at room temperature could alter the orientation of magnetic bits.

Spin-Valve Read Head




Uses “giant magnetoresistance (GMR)”
and “exchange bias”
--in every high-speed read head now

Crucial surfaces &
buried interfaces
everywhere,
as well as complex
materials
(e.g. colossal
magnetoresistance
(CMR))

Some new directions with magnetic nanolayer structures--”spintronics”

Magnetic Random Access Memory (MRAM-Non Volatile)



Up to 100 Mbit devices in R&D: applications to e.g. cell phone use

Scientific and technological areas related to surface/interface/nano science:

- **Integrated circuits**—higher speed, higher density
- **Magnetic storage**—higher density, magnetic logic
- **Catalysis**—auto catalytic converter, petrochemical processing
- **Corrosion**—major annual economic cost
- **Polymer surface modification**—promote adhesion, fire resistance,...
- **Batteries, fuel cells**—the hydrogen economy?
- **Lubrication (tribology)**—nanometer-scale layers
- **Atmospheric particulates**—ice, carbonaceous,...
- **Nuclear reactors and waste storage**—how long-lasting?
- **Environmental science**—retention of contaminants in soil
- **Biomaterials**—compatibility through surface interactions
- **Sensors**—surface reactions→change in voltage, resistance

Outline

Surface, interface, and nanoscience—short introduction

→ Some surface concepts and techniques→photoemission

Synchrotron radiation: experimental aspects

Electronic structure—a brief review




**The basic synchrotron radiation techniques:
more experimental and theoretical details**

Valence-level photoemission


Core-level photoemission

**Photoemission with high ambient pressure
around the sample**

Electrons as surface probes: the electron inelastic mean free paths in solids



Some Comple- mentary Surface Structure Probes




| <u>Type of order:</u> | Short (< 10 Å) | Short, long and disorder | Long (> 100 Å) |
|----------------------------------|--|--------------------------|--|
| <u>Atom & site specific:</u> | Yes | No | No |
| <u>Sensing depth:</u> | 5-40 Å | Mostly surface D.O.S. | 5-20 Å |
| <u>Lateral resolution:</u> | 1 mm^2 to $(300 \text{\AA})^2$ | Single atom | 1 mm^2 to 1 micron^2 |


● Some growth modes:



LAYER-BY-LAYER (FvdM)
EX. Fe/W(110)
Gd/W(110)




MIXED (SK)
Cu/Ru(001)
Gd/Ru(001)




ISLAND/CLUSTER (VW)
3D → 2D → 1D
Fe/Stepped W



INTERDIFFUSION
Fe/Cu(001)




MIXED-PHASE
EPITAXY/METASTABILITY
fcc & bcc Fe/Cu(001)




SUPERLATTICES IN PLANE
STRAIN
most binaries
FeO/Pt(111)
Gd/W(110)




SURFACE ALLOY
Co/Pt



DEFECTS/STEPS
Fe/Cu
Cr/Fe



ROUGHNESS
Co/Cu
Cr/Fe



FLOATING SURFACTANT
Au/Si(111)-Ag




ALLOYING SURFACTANT
Ga/Si(111)-Sn



TEXTURING
Tb-Fe
(Amorphous?)

Some important structures in nanoscience/nanotechnology



KORTRIGHT
ET AL., J.M.R.
207, 44 ('99')

WHAT DO SURFACES LOOK LIKE? SOME fcc AND bcc SURFACES








Fig. 3.2. Low-index ideal surfaces of a hard-sphere cubic crystal. Vertical and horizontal markings indicate the second and third atom layers, respectively. Cube face is indicated for (100) to set the scale (Nicholas, 1965).





Si(111)-(7x7)—Dimer-adatom-stacking fault model


Low-index fcc metal surfaces




Low-index NaCl surfaces



Fcc(111)
super-
lattice = a
Moiré
pattern:
4 degree
rot'n.




Formation of Moire patterns—two rotated square lattices



FRACTION OF ATOMS ON THE SURFACE

OF A CUBE: $D = \text{ATOMIC DIAM.} \approx 0.2 \text{ nm} = 2 \text{\AA}$



$$\text{SURFACE FRACTION} = \frac{L^3 - (L-2D)^3}{L^3}$$

| <u>L</u> | <u>FRACTION</u> |
|------------------------------------|------------------------|
| $1 \mu\text{m} = 1000 \text{ nm}$ | $0.001 \approx 0.1\%$ |
| $0.1 \mu\text{m} = 100 \text{ nm}$ | $0.012 \approx 1.2\%$ |
| $0.01 \mu\text{m} = 10 \text{ nm}$ | $0.115 \approx 11.5\%$ |
| $0.001 \mu\text{m} = 1 \text{ nm}$ | $0.784 = 78.4\%$ |

Nanoscience
is surface science

SOME UNITS :


1 HAIR \approx 50 microns

1 micron $= 10^{-6} \text{ m} = 1,000 \text{ nm} = 10,000 \text{\AA}$
 $\approx 5,000 \text{ atoms}$

0.001 micron $= 10^{-9} \text{ m} = 1 \text{ nanometer} = 1 \mu\text{m} = 10 \text{\AA}$
 $\approx 5 \text{ atoms}$

WHY IS ULTRAHIGH VACUUM IMPORTANT?

TIME TO BUILD UP A SINGLE
ATOMIC/MOLECULAR LAYER ≈ 1
MONOLAYER $\approx 1 \text{ ml}$ IF EACH ATOM/
MOLECULE FROM GAS PHASE HITTING
SURFACE STICKS: τ_1



$$\tau_1 (\text{sec}) = 2.84 \times 10^{-23} [T(\text{K})M]^{1/2} \rho_s (\text{cm}^{-2}) / P(\text{torr})$$

$$\begin{aligned} & N_2, CO, O_2 \\ & \downarrow \quad \downarrow \\ & \text{WITH TYPICAL MOS. FOR } M \approx 28, 32 \\ & T = 298 \text{ K} \\ & \rho_s = 1-2 \times 10^{15} \text{ cm}^{-2} \end{aligned}$$

| <u>τ_1</u> | <u>P</u> | |
|----------------------------|----------------|--|
| 1 s | 10^{-6} torr | |
| 100 s | 10^{-8} .. | |
| $\sim 2 \text{ min}$ | | |
| $\sim 15 \text{ min}$ | 10^{-9} .. | |
| $\sim 2.8 \text{ hr}$ | 10^{-10} .. | |
| $\sim 27.8 \text{ hr}$ | 10^{-11} .. | |


→ Need to work
at $\sim 10^{-10}-10^{-11}$ torr

Table 4 Density and atomic concentration

The data are given at atmospheric pressure and room temperature, or at the stated temperature in deg K. (Crystal modifications as for Table 3.)



| | | | | | | | | | | | | | | | | |
|---------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-------------------------|-----------|-----------|-----------|----------------------|-----------|
| H 4K | | | | | | | | | | | | | | | He 2K | |
| 0.088 | | | | | | | | | | | | | | | 0.205 (at 37 atm) | |
| Li 78K | Be | | | | | | | | | | | | | | | |
| 0.542 | 1.82 | | | | | | | | | | | | | | | |
| 4.700 | 12.1 | | | | | | | | | | | | | | | |
| 3.023 | 2.22 | | | | | | | | | | | | | | | |
| Na 5K | Mg | | | | | | | | | | | | | | | |
| 1.013 | 1.74 | | | | | | | | | | | | | | | |
| 2.652 | 4.30 | | | | | | | | | | | | | | | |
| 3.659 | 3.20 | | | | | | | | | | | | | | | |
| K 5K | Ca | Sc | Ti | V | Cr | Mn | Fe | Co | Ni | Cu | Zn | Ga | Ge | As | Se | |
| 0.910 | 1.53 | 2.99 | 4.51 | 6.09 | 7.19 | 7.47 | 7.87 | 8.9 | 8.91 | 8.93 | 7.13 | 5.91 | 5.32 | 5.77 | 4.81 | |
| 1.402 | 2.30 | 4.27 | 5.66 | 7.22 | 8.33 | 8.18 | 8.50 | 8.97 | 9.14 | 8.45 | 6.55 | 5.10 | 4.42 | 4.65 | 3.67 | |
| 4.525 | 3.95 | 3.25 | 2.89 | 2.62 | 2.50 | 2.24 | 2.48 | 2.50 | 2.49 | 2.56 | 2.66 | 2.44 | 2.45 | 3.16 | 2.32 | |
| Rb 5K | Sr | Y | Zr | Nb | Mo | Tc | Ru | Rh | Pd | Ag | Cd | In | Sn | Sb | Te | |
| 1.629 | 2.58 | 4.48 | 6.51 | 8.58 | 10.22 | 11.50 | 12.36 | 12.42 | 12.00 | 10.50 | 8.65 | 7.29 | 5.76 | 6.69 | 6.25 | |
| 1.148 | 1.78 | 3.02 | 4.29 | 5.56 | 6.42 | 7.04 | 7.36 | 7.26 | 6.80 | 5.85 | 4.64 | 3.83 | 2.91 | 3.31 | 2.94 | |
| 4.837 | 4.30 | 3.55 | 3.17 | 2.86 | 2.72 | 2.71 | 2.65 | 2.69 | 2.75 | 2.89 | 2.98 | 3.25 | 2.81 | 2.91 | 2.86 | |
| Cs 5K | Ba | La | Hf | Ta | W | Re | Os | Ir | Pt | Au | Hg₂₂₇ | Tl | Pb | Bi | Po | At |
| 1.997 | 3.59 | 6.17 | 13.20 | 16.66 | 19.25 | 21.03 | 22.58 | 22.55 | 21.47 | 19.28 | 14.26 | 11.87 | 11.34 | 9.80 | 9.31 | |
| 0.905 | 1.60 | 2.70 | 4.52 | 5.55 | 6.30 | 6.80 | 7.14 | 7.06 | 6.62 | 5.90 | 4.26 | 3.50 | 3.30 | 2.82 | 2.67 | |
| 5.235 | 4.35 | 3.73 | 3.13 | 2.86 | 2.74 | 2.74 | 2.68 | 2.71 | 2.77 | 2.88 | 3.01 | 3.46 | 3.50 | 3.07 | 3.34 | |
| Fr | Ra | Ac | | | | | | | | | | | | | | |
| | | 10.07 | Ce | Pr | Nd | Pm | Sm | Eu | Gd | Tb | Dy | Ho | Er | Tm | Yb | |
| — | — | 2.66 | 6.77 | 6.78 | 7.00 | — | 7.54 | 5.25 | 7.89 | 8.27 | 8.53 | 8.80 | 9.04 | 9.32 | 6.97 | |
| | | 3.76 | 2.91 | 2.92 | 2.93 | — | 3.03 | 2.04 | 3.02 | 3.22 | 3.17 | 3.22 | 3.26 | 3.32 | 3.02 | |
| | | | 3.65 | 3.63 | 3.66 | — | 3.59 | 3.96 | 3.58 | 3.52 | 3.51 | 3.49 | 3.47 | 3.54 | 3.88 | |
| | | | Th | Pa | U | Np | Pu | Am | Cm | Bk | Cf | Es | Fm | Md | No | |
| | | | 11.72 | 15.37 | 19.05 | 20.45 | 19.81 | 11.87 | — | — | — | — | — | — | — | |
| | | | 3.04 | 4.01 | 4.80 | 5.20 | 4.26 | 2.96 | — | — | — | — | — | — | — | |
| | | | 3.60 | 3.21 | 2.75 | 2.62 | 3.1 | 3.61 | — | — | — | — | — | — | — | |

SOME SURFACE-ANALYTICAL TECHNIQUES






★ = AT UC DAVIS

Why are electrons so useful as probes of surfaces?



COMPILATIONS: Seah & Dench, Surf. Int. Anal. 1, 2 (1979)
Tanuma, Powell, + Penn, Surf. Int. Anal.
13, 919 + 927 (1991); 21, 165 (1994)
Powell & Jablonski, J. Phys. Chem. Ref.
Data 28, 19 (1999); Surf. Int. Anal.
29, 108 (2000)




*More realistic look at
low-energy inelastic
mean free paths--
theory*

**IMFPs do not
necessarily go up
drastically at lower
energy**

S.F. Mao and Z.J.
Ding, Hefei (TBP)

How much deeper do we probe at 5-10 keV?


Au
Inelastic attenuation length
TPP-2M formula
of Tanuma, Powell, Penn




<http://www.ss.teen.setsunan.ac.jp/e-imfp.html>

SOME SURFACE-ANALYTICAL TECHNIQUES


① PHOTOELECTRON SPECTROSCOPY (NOBEL, '21+'81)



③ AUGER ELECTRON SPECTROSCOPY




⑤ SECONDARY ION MASS SPECTROMETRY




② LOW-ENERGY ELECTRON DIFFRACTION (NOBEL, '37)




④ LOW-ENERGY ELECTRON LOSS SPECTROSCOPY



⑥ RUTHERFORD SCATT./ION SCATTERING (NOBEL, '08)




⑦ SCANNING TUNNELING MICROSCOPY (NOBEL, '86)



LOW ENERGY ELECTRON DIFFRACTION


LEED:




Davisson &
Germer (1927):
Electrons are
de Broglie waves

TWO-DIMENSIONAL
SURFACE RECIPRO-
CAL LATTICE


LONG-RANGE
ORDER REQUIRED
OVER $\gtrsim 100\text{\AA}$.



SOME TYPICAL LEED PATTERNS:



$\text{Si}(111)\text{-(}7\times 7\text{)}$





$(\sqrt{3}\times\sqrt{3})\text{R}30^\circ \text{Ag/Si}(111)$

● = spots seen without any
reconstruction or adsorption
of simple $\text{Si}(111)$ surface


LEED: Si(111)7x7

Real Space:

Si surface atoms



- Longer periodicities in real space give closer spots in k-space.
- Higher energy LEED images show spots closer together.
K-Space



SCANNING TUNNELING MICROSCOPY





Figure 4 Scanning tunneling microscopes can be operated in either (a) the constant current mode or (b) the constant height mode. The images of the surface of graphite were made by Richard Sonnenfeld at the University of California at Santa Barbara. The constant height mode was first used by A. Bryant, D. P. E. Smith, and C. F. Quate, *Applied Physics Letters* 48: 832, 1986.

IMAGING, AND
MANIPULATING,
ATOMS AT SURFACES
WITH THE STM




48 iron atoms on a Cu(111) surface—a “quantum corral”


- Some growth modes:



LAYER-BY-LAYER (FvdM)
EX. Fe/W(110)
Gd/W(110)



MIXED (SK)
Cu/Ru(001)
Gd/Ru(001)




ISLAND/CLUSTER (VW)
3D → 2D → 1D
Fe/Stepped W



INTERDIFFUSION
Fe/Cu(001)




MIXED-PHASE
EPITAXY/**METASTABILITY**
fcc & bcc Fe/Cu(001)




SUPERLATTICES IN PLANE
STRAIN
most binaries
FeO/Pt(111)
Gd/W(110)




SURFACE ALLOY
Co/Pt



DEFECTS/STEPS
Fe/Cu
Cr/Fe




ROUGHNESS
Co/Cu
Cr/Fe



FLOATING SURFACTANT
Au/Si(111)-Ag



ALLOYING SURFACTANT
Ga/Si(111)-Sn



TEXTURING
Tb-Fe
(Amorphous?)

Scanning tunneling microscopy: stepped Si(111) surface
















Fig. 2. Tunneling image of silicon (111) surface that shows the 7×7 atomic reconstruction on terraces separated by atomic steps.

Some growth modes:



Scanning tunneling microscopy: metal-on-metal epitaxial growth



**Growth mode depends
strongly on anneal
temperature!**

Tober et al.
Phys. Rev. B
53, 5444 (1996).


- Some growth modes:



LAYER-BY-LAYER (FvdM)
EX. Fe/W(110)
Gd/W(110)



MIXED (SK)
Cu/Ru(001)
Gd/Ru(001)




ISLAND/CLUSTER (VW)
3D → 2D → 1D
Fe/Stepped W



INTERDIFFUSION
Fe/Cu(001)




MIXED-PHASE
EPITAXY/METASTABILITY
fcc & bcc Fe/Cu(001)




SUPERLATTICES IN PLANE
most binaries
FeO/Pt(111)
Gd/W(110)




SURFACE ALLOY
Co/Pt



DEFECTS/STEPS
Fe/Cu
Cr/Fe



ROUGHNESS
Co/Cu
Cr/Fe



FLOATING SURFACTANT
Au/Si(111)-Ag



ALLOYING SURFACTANT
Ga/Si(111)-Sn




TEXTURING
Tb-Fe
(Amorphous?)


**Superlattice =
Moiré structure
in metal-on-
metal
epitaxial
growth**

"WETTING" SINGLE MONOLAYER OF Gd ON W(110)


a) STM:



b) LEED:




c) STM:




(7x14) Moiré pattern
= superlattice

d)




E. Tober et al.
Phys. Rev. B
53, 544 ('96)

A Moiré pattern—Monolayer Gd on W(110)




A typical surface science research system

≥ 1 TECHNIQUE: SURFACE SENSITIVE (e^- , IONS,
ATOMS AS PROBES)
NON-DESTRUCTIVE



Laboratory-based surface science system:



Outline

Surface, interface, and nanoscience—short introduction

Some surface concepts and techniques→photoemission

 **Synchrotron radiation: experimental aspects**

Electronic structure—a brief review


**The basic synchrotron radiation techniques:
more experimental and theoretical details**

Valence-level photoemission

Core-level photoemission



**Photoemission with high ambient pressure
around the sample**

THE ELECTROMAGNETIC SPECTRUM



Typical
surface/materials
science expts.

Producing x-rays: the good old-fashioned way




See Section
1.2 in “X-Ray
Data Booklet”

**X-Ray
Nomenclature
(from “X-Ray
Data Booklet”)**

In general:

$$nl \xrightarrow{\text{Spin-orbit}} nl_{j=l+1/2} \quad nl_{j=l-1/2}$$



See Section
1.2 in “X-Ray
Data Booklet”

Fig. I-1. Transitions that give rise to the emission lines in Table 1-3.



X-Ray energies from the “X-Ray Data Booklet”

Table 1-2. Photon energies, in electron volts, of principal K-, L-, and M-shell emission lines.

| Element | K α_1 | K α_2 | K β_1 | L α_1 | L α_2 | L β_1 | L β_2 | L γ_1 | M α_1 |
|---------|--------------|--------------|-------------|--------------|--------------|-------------|-------------|--------------|--------------|
| 3 Li | 54.3 | | | | | | | | |
| 4 Be | 108.5 | | | | | | | | |
| 5 B | 183.3 | | | | | | | | |
| 6 C | 277 | | | | | | | | |
| 7 N | 392.4 | | | | | | | | |
| 8 O | 524.9 | | | | | | | | |
| 9 F | 676.8 | | | | | | | | |
| 10 Ne | 848.6 | 848.6 | | | | | | | |
| 11 Na | 1,040.98 | 1,040.98 | | 1,071.1 | | | | | |
| 12 Mg | 1,253.60 | 1,253.60 | | 1,302.2 | | | | | |
| 13 Al | 1,486.70 | 1,486.27 | | 1,557.45 | | | | | |
| (14 Si) | 1,739.98 | 1,739.38 | | 1,835.94 | | | | | |
| 15 P | 2,013.7 | 2,012.7 | | 2,139.1 | | | | | |
| 16 S | 2,307.84 | 2,306.64 | | 2,464.04 | | | | | |
| 17 Cl | 2,622.39 | 2,620.78 | | 2,815.6 | | | | | |
| 18 Ar | 2,957.70 | 2,955.63 | | 3,190.5 | | | | | |
| 19 K | 3,313.8 | 3,311.1 | | 3,589.6 | | | | | |
| 20 Ca | 3,691.68 | 3,688.09 | 4,012.7 | 341.3 | 341.3 | 344.9 | | | |
| 21 Sc | 4,090.6 | 4,086.1 | 4,460.5 | 395.4 | 395.4 | 399.6 | | | |


Popular laboratory sources
for photoelectron spectroscopy

Synchrotron Radiation Sources:




Intensity $\propto 1$

BENDING MAGNETS AND WIGGLERS generate fan-shaped beams of synchrotron radiation, whereas undulators emit pencil-thin beams.




Inside the Advanced Light Source

Electron
speed near c:
 $0.99999997 c$




Undulator Radiation



Our primary experimental activity




Synchtron Radiation Sources of the World: ~ 40 operating, 10 planned



<http://www.srs.ac.uk/srs/SRworldwide/>

Beamlines at the ALS 2007







Fig. 2. Average brightness comparisons of the LCLS and other light sources, including proposed FELs at Brookhaven [14] and DESY [15].

"X-Ray Data Booklet"
See Fig. 2.9

Advanced Light Source-- Typical Spectroscopy Beamline Layout



The five ways in which x-rays Interact with Matter:






Fig. 3-1. Total photon cross section σ_{tot} in carbon, as a function of energy, showing the contributions of different processes: τ , atomic photo-effect (electron ejection, photon absorption); σ_{coh} , coherent scattering (Rayleigh scattering—atom neither ionized nor excited); σ_{incoh} , incoherent scattering (Compton scattering off an electron); κ_n , pair production, nuclear field; κ_e , pair production, electric field; σ_{ph} , photonuclear absorption (nuclear absorption usually followed by emission of a neutron or other particle). (From Ref. 3; figure courtesy of J. H. Hubbell.)

"X-Ray Data Booklet"
Section 3.1

The ultraviolet, soft x-ray, hard x-ray measurements:



The Soft X-Ray Spectroscopies




PE = photoemission = photoelectron spectroscopy

XAS = x-ray absorption spectroscopy


AES = Auger electron spectroscopy

XES = x-ray emission spectroscopy

RIXS = resonant inelastic x-ray scattering / x-ray Raman scatt.




Some basic measurements:



**MULTI-TECHNIQUE
PHOTOELECTRON
SPECTROMETER/
DIFFRACTOMETER (MTSD)**


Loadlock
for sample
introduction

Soft x-ray
spectrometer



**MULTI-TECHNIQUE
SPECTROMETER/
DIFFRACTOMETER (MTSD)**

ALS
 $h\nu$ 



**5-axis
sample
manipulator**

**Sample prep.
chamber: LEED,
Knudsen cells,
electromagnet,...**

**Soft x-ray
spectrometer**