



2244-4

Summer School on Particle Physics

6 - 17 June 2011

Higgs and Electroweak Symmetry Breaking - IV

Sally DAWSON Brookhaven National Laboratory USA

EWSB Beyond the Standard Model

Lecture 4, Trieste, 2011 S. Dawson (BNL)

Supersymmetric Models as Alternative to Standard Model

Many New Particles:

- Spin $\frac{1}{2}$ quarks \Rightarrow spin 0 squarks
- Spin $\frac{1}{2}$ leptons \Rightarrow spin 0 sleptons
- Spin 1 gauge bosons \Rightarrow spin $\frac{1}{2}$ gauginos
- Spin 0 Higgs \Rightarrow spin $\frac{1}{2}$ Higgsino

Unbroken supersymmetry \Rightarrow degenerate masses of partners

SUSY must be a broken symmetry

Supersymmetric Theories

- Predict many new undiscovered particles (>29!)
- Very predictive models
 - Can calculate particle masses, interactions, everything you want in terms of a few parameters
 - Solve naturalness problem of Standard Model
- Any Supersymmetric particle eventually decays to the lightest supersymmetric particle (LSP) which is stable and neutral (assuming R parity)
 - Dark Matter Candidate

SUSY Models Unify

- Coupling constants change with energy
- Assume new particles at TeV scale



SUSY....Our favorite model

- Quadratic sensitivity to high scale physics cancelled automatically if SUSY particles at TeV scale
- Cancellation result of *supersymmetry*, so happens at every order



Stop mass should be TeV scale

Supersymmetry (MSSM version)

 Good agreement with EW measurements if SUSY masses are 1-2 TeV



Fermion Masses

• In Standard Model, m_u from $\Phi_c = i\sigma_2 \Phi^*$

$$L_{SM} = -\lambda_u \overline{Q}_L \Phi_c u_R + hc \qquad \Phi_c = \begin{pmatrix} \overline{\phi}^0 \\ -\phi^- \end{pmatrix} \qquad \lambda_u = -\frac{m_u \sqrt{2}}{v_{SM}}$$

- SUSY models don't allow Φ_c interactions
- Supersymmetric models always have at least *two Higgs doublets* with opposite hypercharge in order to give mass to up and down quarks

General 2 Higgs Doublet Model

• 6 free parameters, plus a phase

$$V(H_{1}, H_{2}) = \lambda_{1}(H_{1}^{+}H_{1} - v_{1}^{2})^{2} + \lambda_{2}(H_{2}^{+}H_{2} - v_{2}^{2})^{2} + \lambda_{3}[(H_{1}^{+}H_{1} - v_{1}^{2}) + (H_{2}^{+}H_{2} - v_{2}^{2})]^{2} + \lambda_{4}[(H_{1}^{+}H_{1})(H_{2}^{+}H_{2}) - (H_{1}^{+}H_{2})(H_{2}^{+}H_{1})] + \lambda_{5}[\operatorname{Re}(H_{1}^{+}H_{2}) - v_{1}v_{2}\cos\xi]^{2} + \lambda_{6}[\operatorname{Im}(H_{1}^{+}H_{2}) - v_{1}v_{2}\sin\xi]^{2} \langle H_{1} \rangle = \begin{pmatrix} 0 \\ v_{1} \end{pmatrix} \langle H_{2} \rangle = \begin{pmatrix} 0 \\ v_{2}e^{i\xi} \end{pmatrix}$$

- W and Z masses just like in Standard Model $M_W^2 = \frac{g^2(v_1^2 + v_2^2)}{2}$
- ρ parameter:

$$\rho = \frac{M_W}{M_Z \cos \theta_W} = 1$$

 ρ =1 for any number of Higgs doublets or singlets

Higgs Potential Restricted in SUSY Models

• Two Higgs doublets with opposite hypercharge

$$H_2 = \begin{pmatrix} \phi_2^+ \\ \phi_2^0 \end{pmatrix} \qquad \qquad H_1 = \begin{pmatrix} \phi_1^{0^*} \\ -\phi_1^- \end{pmatrix}$$

- Quartic couplings fixed by supersymmetry
- H_1 has Y=-1/2, H_2 has Y=+1/2 [Q_{em}=(T₃+Y)/2]
- SUSY Potential has "F" and "D" terms:

 $V_{F} = |\mu|^{2} (|H_{1}|^{2} + |H_{2}|^{2})$ $V_{D} = \frac{1}{2} \sum_{a} (g_{a} H_{i}^{*} T^{a} H_{i})^{2} \qquad Sl$

$$U(1): \qquad \left[\frac{g'}{2}\left(\left|H_{2}\right|^{2}-\left|H_{1}\right|^{2}\right)\right]^{2}$$
$$SU(2): \qquad \left[\frac{g}{2}\left(H_{1}^{i*}\sigma_{ij}^{a}H_{1}^{j}+H_{2}^{i*}\sigma_{ij}^{a}H_{2}^{j}\right)\right]^{2}$$

Gauge couplings / generators

SUSY Potential

• SU(2) identity: $\sigma_{ij}^a \sigma_{kl}^a = 2\delta_{il}\delta_{jk} - \delta_{ij}\delta_{kl}$

 $V_{D} = \frac{g^{2}}{8} \left(4 \left| H_{1}^{*} \cdot H_{2} \right|^{2} - 2 \left(H_{1}^{*} \cdot H_{1} \right) \left(H_{2}^{*} \cdot H_{2} \right) + \left(\left| H_{1} \right|^{2} \right)^{2} + \left(\left| H_{2} \right|^{2} \right)^{2} \right) + \frac{g'^{2}}{8} \left(\left| H_{2} \right|^{2} - \left| H_{1} \right|^{2} \right)^{2} \right)$

Scalar potential

$$V = \left|\mu\right|^{2} \left(\left|H_{1}\right|^{2} + \left|H_{2}\right|^{2}\right) + \frac{g^{2} + {g'}^{2}}{8} \left(\left|H_{2}\right|^{2} - \left|H_{1}\right|^{2}\right)^{2} + \frac{g^{2}}{2} \left|H_{1}^{*} \cdot H_{2}\right|^{2}$$

• V is positive definite: minimum at <V>=<H₁>=<H₂>=0

No electroweak symmetry breaking!

The MSSM Philosophy

- Add all soft (dimension 3 or less) terms allowed
 They don't introduce quadratic Λ² contributions
- Potential has 3 free parameters (1 of which is fixed by M_W)

$$V = (m_{1}^{2} + |\mu|^{2})H_{1}H_{1}^{+} + (m_{2}^{2} + |\mu|^{2})H_{2}H_{2}^{+} - m_{12}^{2}(\varepsilon_{ab}H_{1}^{a}H_{2}^{b} + h.c.)$$

+ $(g^{'2} + g^{2})H_{1}H_{1}^{+} - H_{2}H_{2}^{+})^{2} + (g^{2})H_{1}H_{2}^{+}|^{2}$
Gauge Couplings

 If m₁₂=0, potential is positive definite and no symmetry breaking

Minimize Potential

• Minimize potential

$$\frac{\partial V}{\partial h_1^0} = 2\left(m_1^2 + |\mu|^2\right)h_1^0 - 2m_{12}^2h_2^{0*} + \frac{g^2 + {g'}^2}{2}h_1^0\left(\left|h_1^0\right|^2 - \left|h_2^0\right|^2\right) = 0$$

$$\frac{\partial V}{\partial h_2^0} = 2\left(m_2^2 + |\mu|^2\right)h_2^0 - 2m_{12}^2h_1^{0*} - \frac{g^2 + {g'}^2}{2}h_2^0\left(\left|h_1^0\right|^2 - \left|h_2^0\right|^2\right) = 0$$

• Minimization conditions:

$$m_{12}^{2} = \frac{\left(m_{1}^{2} + m_{2}^{2} + 2\mu^{2}\right)\sin 2\beta}{2}$$

$$\mu^{2} = \frac{m_{2}^{2} - m_{1}^{2}\tan^{2}\beta}{\tan^{2}\beta - 1} - \frac{g^{2} + g'^{2}}{4}\left(v_{1}^{2} + v_{2}^{2}\right)}{4} - \frac{M_{Z}^{2}}{2}$$

$$\mu \text{ is naturally of O(M_{Z})}$$

EWSB and SUSY Models

• Electroweak symmetry broken by vevs

$$\langle H_1 \rangle = \begin{pmatrix} v_1 \\ 0 \end{pmatrix} \qquad \langle H_2 \rangle = \begin{pmatrix} 0 \\ v_2 \end{pmatrix}$$

- W gets mass, $M_W^2 = g^2 (v_1^2 + v_2^2)/2$
- 5 Physical Higgs bosons, h⁰, H⁰, H[±], A⁰
- 2 free parameters, typically pick

 M_A , tan $\beta = v_2/v_1$

• Predict M_h , M_H , $M_{H^{\pm}}$

$$M_A^2 = m_{12}^2 \left(\tan \beta + \cot \beta \right)$$
$$M_{H^{\pm}}^2 = M_A^2 + M_W^2$$

Neutral Higgs Masses

$$M_{h,H}^{2} = \frac{1}{2} \left[M_{A}^{2} + M_{Z}^{2} \pm \sqrt{\left(M_{A}^{2} + M_{Z}^{2}\right)^{2} - 4M_{Z}^{2}M_{A}^{2}\cos^{2}2\beta} \right]$$

- $M_h < M_Z \cos 2\beta$
- Theory implies light Higgs boson!
- \bullet Neutral Higgs mass matrix diagonalized with mixing angle α

$$\cos 2\alpha = -\cos 2\beta \left(\frac{M_A^2 - M_Z^2}{M_H^2 - M_h^2}\right)$$

Theoretical Upper Bound on M_h

- At tree level, $M_h < M_Z$
- Large corrections O(G_Fmt²)
 - Predominantly from stop squark loop

$$M_{h}^{2} \le M_{Z}^{2} \cos^{2} 2\beta + \frac{3G_{F}m_{t}^{4}}{\sqrt{2}\pi^{2} \sin^{2} \beta} \ln \left[\frac{\widetilde{m}_{t}^{2}}{m_{t}^{2}}\right]$$

• Stop mass should be TeV scale for naturalness

Theoretical Upper Bound on M_h



- Mt⁴ enhancement
- Logarithmic dependence on stop mass

Higgs Masses in MSSM

$$M_{H^{\pm}}^2 = M_A^2 + M_W^2$$



Large M_A: Degenerate A, H, H[±] and light h

Find Higgs Couplings

• Higgs-fermion couplings:

$$L = -\frac{gm_d}{2M_W \cos\beta} \overline{d}d(H\cos\alpha - h\sin\alpha) + \frac{igm_d \tan\beta}{2M_W} \overline{d}\gamma_5 dA$$
$$-\frac{gm_u}{2M_W \sin\beta} \overline{u}u(H\sin\alpha + h\cos\alpha) + \frac{igm_d \cot\beta}{2M_W} \overline{u}\gamma_5 uA$$

- Couplings given in terms of α , β
- Can be very different from Standard Model
- No new free parameters

Higgs Couplings Different from SM

Lightest Neutral Higgs, h



Higgs Couplings in SUSY Heavier Neutral Higgs, H



Gauge Boson Couplings to Higgs

- $g_{hVV}^2 + g_{HVV}^2 = g_{hVV}^2(SM)$
- Vector boson fusion and Wh production always suppressed in MSSM

$$\frac{g_{hVV}}{g_{h,smVV}} = \sin(\beta - \alpha)$$
$$\frac{g_{HVV}}{g_{HVV}} = \cos(\beta - \alpha)$$
$$g_{h,smVV}$$



Tools

- Calculate SM and MSSM Higgs branching ratios:
 - HDECAY
 - http://people.web.psi.ch/spira/hdecay/
- Calculate MSSM Higgs masses and Higgs branching ratios:
 - FEYNHIGGS
 - http://www.feynhiggs.de/

Both of these programs are very easy to use!

Limits from LEP



Limits on SUSY Higgs from LEP



Remember Higgs Decays in SM

• SM: Higgs branching rates to $b\overline{b}$ and $\tau^+\tau^-$ turn off as rate to W⁺W⁻ turns on (M_h > 160 GeV)



Higgs Decays Changed at Large tan β

• MSSM: At large tan β , rates to bb and $\tau^+\tau^-$ large



Rate to bb and $\tau^+\tau^-$ almost constant in MSSM for H, A

Large tanβ Changes Relative Importance of Production Modes



 $\tan\beta \ge 7$, bb production mode larger than gg

Higgs Production Can be Larger than SM

- SUSY Higgs: tan β enhanced couplings to b and τ for H,A
- Production with b's dominates for large M_H



New Higgs Discovery Channels in SUSY



 $bb\phi$ coupling enhanced for large $tan\beta$



Many Possibilities Beyond SUSY

- Add singlet Higgs and try to evade LEP bounds
- Two Higgs doublets, but not SUSY
 - Same spectrum as SUSY
 - Must measure Higgs couplings
- Little Higgs Models
 - Have extended gauge sectors and new charge 2/3 quarks

Effective Lagrangian approach needed to study EWSB sector if no new particles found at LHC

Possibilities at the LHC

- We find a light Higgs with SM couplings and nothing else
 How to answer our questions?
- We find a light Higgs, but it doesn't look SM like
 - Most models (SUSY, Little Higgs, etc) have other new particles
- We don't find a Higgs (or any other new particles)
 - How can we reconcile precision measurements?
 - This is the hardest case

The fun is just beginning Happy Higgs Hunting!