



2256-16

Workshop on Aerosol Impact in the Environment: from Air Pollution to Climate Change

8 - 12 August 2011

Aerosols and Weather Modification including Geo-Engineering principles

R.T. Bruintjes

National Center for Atmospheric Research, Colorado

USA


CLOUD SEEDING TO ENHANCE RAINFALL: CURRENT STATUS AND FUTURE DIRECTIONS


Workshop on Aerosol Impact in the Environment: from Air Pollution to Climate Change

8-12 August 2111 ICTP, Trieste, Italy

The Paradox

- Operational programs in more than 37 countries worldwide
- At least 69 programs in 11 U.S. states in 2005
 - Limited funding supporting research as part of operational programs
 - Impact of human activity on weather and climate recognized and heavily funded










- What role does aerosols play in warm and mixed phase processes in deep convective clouds?
- How does aerosol effects compare with other effects such as thermodynamics (stable layers and updraft speed) and cloud base temperature?
- What are the implications for cloud seeding experiments?


Total aerosol, cloud condensation nuclei (CCN), and ice nuclei (IN) concentrations as a function of temperature.


■ In order to enhance precipitation the concept of seeding is to seed with appropriate CCN or IN to make precipitation develop more efficiently.




Cloud characteristics changes













Fundamentals

What are clouds made of?





Cloud droplets


- Condensation on hygroscopic particles (a specific kind of water-soluble aerosol)
- Droplet growth by condensation
 - Condensation occurs through diffusion of water vapor
 - Cloud droplets are typically in the size range of 5-25 μ m (i.e., very small)
 - They have a very small terminal velocity
- Result: No precipitation from only cloud droplets. The water stays in the cloud

Warm rain process

NCAR


 Collision and coalescence of droplets falling at different terminal velocities leads to raindrop formation

Raindrops are millimeters in size



Cloud droplets are 100 times smaller in diameter






The cold rain process



 Co-existence of ice crystals and <u>supercooled</u> droplets leads to rapid vapor growth of crystals, followed in suitable convective clouds by accretional growth (riming), fall-out, and melting as rain









The origins of cloud seeding...




- Freeze supercooled droplets by introducing artificial "seeds"
- Langmuir and Schaeffer, Schenectady, NY, Nov 13, 1946: Dropped 3 pounds of dry ice pellets into a stratus deck at 14,000 ft, -20C, flying in a race-track pattern



WEATHER MODIFICATION







The Initial Age


- Spectacular claims (rain, hail, hurricanes, snow, etc)
- Problems with scientific evaluation
 - How to detect seeding effect (10-50%) against a background of natural variability some 10-100 times greater?



- Emergence of the professional statistician into the field
- Physical hypotheses stated; almost none actually checked
- New Age: starting in the 1970s with direct measurements and advanced computer models
 - Airborne instruments, digital data
 - Doppler radar
 - Coordinated experiments
 - 3D cloud modeling
 - Radiometers







- Much has been learned about natural rainfall and seeding
 - Warm vs cold rain
- The fundamental weather mod result: Agl seeding produces lots of ice crystals
- The effects of seeding on precipitation are essentially never as clear-cut as in the stratus cloud photo. Clouds are complex.
- Controversial results regarding Agl seeding for rain increase from convective clouds
 - Re-analysis of experiments; effects less clear-cut
- Better progress with continued efforts studying winter orographic snowfall enhancement and hygroscopic seeding of convective clouds
- Greatly decreased funding since the '80s

Principal Conclusion (NAS 2003)

Advances in atmospheric sciences positions us to mount concerted and sustained effort to delineate scope and expectations of future weather modification research

- Must be directed at answering fundamental scientific questions
- Will yield results that go beyond application to weather modification (e.g., NWP, QPF)
- Emphasis be on understanding processes and results

Scientific approach



- Preliminary study (needs, area, clouds, rainfall, tools, human resources)
- Design (infrastructure, training, tools, conceptual model, operations, evaluation)
- Execution (operations plan, data gathering, processing and archival, quality control, analyses, evaluation)
- Evaluation (assessment, hydrological, economical and social impacts and cost/benefit assessment)


ALL STEPS ARE CRITICAL IN THE EXECUTION OF A SCIENTIFICALLY BASED PROGRAM TO QUANTIFY THE RESULTS


PRELIMINARY STUDIES


WHY?


TEMPORAL AND SPATIAL VARIATIONS IN:


- Climatology of clouds and precipitation in a region
- Thermodynamic structure
- of the atmosphere
- Aerosol and associated
- microphysical variations
















Key Uncertainties



Cloud and precipitation microphysics issues

 Background concentration, sizes, and chemical composition of aerosols participating in cloud processes

Cloud dynamics issues

 Cloud-to-cloud and mesoscale interactions relating to updraft and downdraft structures and cloud evolution and lifetimes

Cloud modeling issues

 Combination of best cloud models with advanced observing systems in carefully designed field tests and experiments

Seeding-related issues

- Targeting of seeding agents, diffusion and transport of seeding material, and spread of seeding effects throughout the cloud volume
- Measurement capabilities and limitations of cell-tracking software, radar, and technologies to observe seeding effects



DESIGN

- Infrastructure
- Conceptual model
- Operations
- Evaluation
- Training

Conceptual Models

Hygroscopic Conceptual Model

Hygroscopic Seeding with Flares

Destabilizes narrow droplet spectra

Broadens cloud droplet spectra near cloud base

& enhances drizzle production

Recirculation helps drizzle enter growth area and spread to other turrets

embryos
Riming rates hig
Increases precipitation at
Potential increas


Riming rates higher
Potential increased secondary
ice generation

May form primary graupel

drizzle above freezing level


Updrafts helps loft

Distribución de gotas uniforme Distribución de gotas no uniforme




cloud base

Colisión/coalescencia mínima



Dynamic seeding conceptual model




DYNAMIC SEEDING CONCEPTUAL MODEL FOR WARM-BASED SUPERCOOLED CHMULI

(REVISED IN JULY 1992)

Water dropfel

Rice cultural

Discrepancy



Good Candidate Clouds

Cumulus Congestus

Or larger... Cumulus clouds deep enough to allow for precipitation development (>2km vertical depth


Narrow Cloud Base Spectra

Formed on a continental aerosol population. Spectra will exhibit high concentration, small mean diameter, and no drops larger than 20 µm

Sustained Updraft


At least 100-300 feet per min

Aerosol, thermodynamic and Cloud




Aerosols, CCN and Cloud droplet concentrations (India)

High concentrations of droplets due to pollution





CCN and aerosol conc.




Broadening of cloud droplet spectra by re-circulation









CCN effect: Difficult to form rain in clouds

Effects on Ice Processes




Large drops freezing


20 00 Timebara 2 1

Concentrations: ~5-10L-1
Similar to concentrations of observed large drops
POTENTIAL INVIGORATION OF CLOUD GROWTH DUE TO LATENT HEAT OF FREEZING


Secondary Ice Formation



Concentrations between 200 to 400L⁻¹





Comparison of "special" natural and "seeded" cases: ICAR



Significant acceleration of the precipitation process results. Cooper et al., 1997

Radar estimate of rainfall within the TITAN framework







Evaluation approaches

- Physical observations
 - Numerical models, observations, theory
- Statistical evaluations
- Radar tracking
- Precipitation gauge stations
- Indirect evaluations
 - Hail: crop damage
 - Rain: River flows, dam levels, etc




Evaluation Issues

- Inadvertent weather modification
- Geographical meteorological differences
- Natural variability
- Inadequate understanding of physical processes
- Controversy: Unsubstantiated claims
- Independent assessment and evaluation
- Old technology

Operation and Execution South African program (1990's) NCAR


- Initiated a randomized experiment using hygroscopic flares
- Some aircraft measurements
- New evaluation methods
 - Used the "storm" as the experimental unit
 - Radar estimated rainfall
 - Many more experimental cases
 - Objective storm-tracking software (TITAN)
 - Allowed for study of time-resolved response to seeding





Results and Questions



 To what extent would weather modification operations be dependent on these background concentrations? (e.g. South African and Mexico experiments)







Response Variables



TIME-SERIES VARIABLES

Radar-estimated precipitation flux (m³ s-1), using the Marshall-Palmer Z-R relationship.

Total storm mass (in kilotons).

Storm mass above 6 km MSL (in kilotons).

Storm area (km²).

(Height of maximum reflectivity) - (Z-weighted vertical centroid).

STORM (time-integrated) VARIABLES

- Total case precipitation in kilotons from decision time until
 60 minutes after decision time.
- •Area-time-integral (ATI) in km²h from decision time until 60 minutes after decision time. (ATI is defined as the integral of storm area over time, computed using discreet volume-scantime intervals.)
- Duration of the experimental unit from decision time to 60 min after decision time.

(a) 7 min. before seeding



(c) 44 min. after seeding



(b) 19 min. after seeding



(d) 50 min. after seeding




Evolution of the seeded cloud on July 23, 1996

STORM DURATION

1997/1998






Precipitation Flux (m³/s)

NCAR


1997/1998




MEXICO EXPERIMENT EXAMPLE



All Storms within Radar Coverage Number of Cases in Each Rain-Out Size Category




Classification of Mexico data using aerosol burden (satellite aerosol optical depth) NCAR



EXAMPLE: USE OF NEW TOOLS CLOUD SEEDING RESEARCH PROGRAM (CSRP) Queensland, AUSTRALIA






Cloud base heights




- Cloud base heights generally got lower as the wet season progressed, leading to deeper warm cloud depths in Feb compared to Nov
 - This may affect the mixed phase microphysics



Aerosol/particles


NCAR



Rain formation: Warm rain process


(Typical in Southeast Queensland clouds)NCAR

Collision and coalescence of droplets falling at different terminal velocities leads to raindrop / formation



Raindrops are millimeters in size

> Cloud droplets are 100 times smaller in diameter




Rain formation: Ice process

NCAR


Ice processes are initiated by freezing of large drizzle/rain drops and subsequent initiation of natural seeding (ice splintering) process rapidly depleting cloud liquid water content

- Large drops freezing at ~-5°C
- Initiation of ice splintering process
 - Rapid depletion of cloud water inhibiting lightning in these cases

Map of lightning



Temperature versus time 27 January 2009




Precipitation Processes:

NCAR

Mixed-phase/ice processes initiated by freezing of large drizzle/rain drops and subsequent initiation of natural seeding (ice splintering) process rapidly depleting cloud liquid water content


- Large drop freezing at ~-5°C
 - Initiation of ice splintering process
- Rapid conversion of LWC to ice
- Rapid depletion of LWC inhibiting lightning in these cases

Temperature versus time 27 January 2009



Dual Polarization Radar







Time height diagrams of radar reflectivity and differential radar reflectivity


Dual Polarization Radar measurements


NCAR







Rain drop size distributions






Cloud base heights and warm cloud depths during Queensland project






13 Feb 2009


- Deep stratiform system
- Little supercooled liquid water
- Lots of ice
- Glaciogenic or hygroscopic seeding not recommended




Monthly lightning maps November 2007 to March 2008












Lower frequency of lightning in coastal regions due to potentially less LWC between -10 and -20°C where naturally most charge buildup occurs






Microphysical relationships





Total peak droplet concentrations (cm⁻³) as a function of standard deviation of the cloud droplet spectra for the 22 cases when penetrations were conducted in deep convection near cloud base in growing non-precipitating parts of the cloud.




Mean diameter of the cloud base doplet spectra as a function of standard deviation of the cloud droplet spectra for the 22 cases when penetrations were conducted in deep convection near cloud base in growing non-precipitating parts of the cloud.

Summary

- Convective clouds produce large amounts of rain
 - Comparable to flow in major rivers
- Agl seeding of cumulus clouds for rain enhancement is a well used technology in many countries
 - However, it is not a well proven technology
- Based on experiments in S. Africa, Mexico, and Thailand, hygroscopic seeding seems promising under specific circumstances
 - Though promising, it is not well understood
- There are many new tools available to establish confidence in seeding technologies


Summary




- Many factors control the amount of rain from convective clouds
- Aerosols (both CCN and IN) clearly affect the cloud particle structure
- Very likely affect the rain also (but positively or negatively?)

However, the exact relationship has proven

difficult to establish



Aerosol change: natural and anthropogenic



And so...

- A program of research should be considered mandatory even for an operational project
 - Preliminary measurements
 - · Aerosol, cloud structure, rainfall climatology
- Should include evaluation of seeding effect
 - Need a well designed, randomized experiment to know if you are spending the sponsor's money wisely
- Need to look at hydrologic aspects: where does the water go?
- Cost/benefit study
- History tells us it's rarely cost beneficial to take a short-cut around the R&D