



**The Abdus Salam
International Centre for Theoretical Physics**



2331-1

Spring School on Superstring Theory and Related Topics

19 - 27 March 2012

A theory of first order dissipative superfluid dynamics

S. Minwalla

Tata Institute of Fundamental Research

A theory of first order dissipative superfluid dynamics

Jyotirmoy Bhattacharya^a, Sayantani Bhattacharyya^b, Shiraz Minwalla^a and Amos Yarom^{c,d}

^a*Dept. of Theoretical Physics, Tata Institute of Fundamental Research
Homi Bhabha Rd, Mumbai 400005, India.*

^b*Harish-Chandra Research Institute, Chhatnag Road, Jhansi, Allahabad-211019.*

^c*Joseph Henry Laboratories, Princeton University, Princeton, NJ 08544*

^d*Department of Physics, Technion, Haifa 32000, Israel*

*Email: jyotirmoy@theory.tifr.res.in, sayanta@hri.res.in,
minwalla@theory.tifr.res.in, ayarom@princeton.edu*

ABSTRACT: We determine the most general form of the equations of relativistic superfluid hydrodynamics consistent with Lorentz invariance, the Onsager principle and the second law of thermodynamics at first order in the derivative expansion. Once parity is violated, either because the $U(1)$ symmetry is anomalous or as a consequence of a different parity-breaking mechanism, our results deviate from the standard textbook analysis of superfluids. Our general equations require the specification of twenty parameters (such as the viscosity and conductivity). In the limit of small relative superfluid velocities we find a seven parameter set of equations. In the same limit, we have used the AdS/CFT correspondence to compute the parity odd contributions to the superfluid equations of motion for a generic holographic model and have verified that our results are consistent.

Contents

1. Introduction	2
2. The theory of charged fluid dynamics	7
2.1 Parity Invariant Charged Fluid Dynamics	11
2.2 Parity non invariant charged fluid dynamics	17
3. Parity invariant Superfluid hydrodynamics	19
3.1 Onshell inequivalent First order independent data	20
3.2 Constructing the entropy current	23
4. Superfluid dynamics without parity invariance	28
4.1 Onshell inequivalent data	28
4.2 Constraints from vanishing of two derivative and curvature terms	29
4.3 Constraints from positivity of the quadratic form	31
5. Summary of results and special limits	39
5.1 The problem addressed	39
5.2 Listing our main result	41
5.3 Weyl invariant superfluid dynamics	45
5.4 The $\zeta \rightarrow 0$ limit	46
6. A holographic computation	49
6.1 A stationary solution	51
6.2 Adding in a small uniform superfluid velocity	53
6.3 Boosting	55
6.4 The gradient expansion	56
7. Discussion	60
A. The linear independence of data for the parity even superfluid	63
A.1 The linear independence of first order terms	63
A.2 The linear independence of the second order terms	67
B. Derivation of thermodynamic identities for normal fluids	69

C. Details relating to parity violating superfluids	71
C.1 All equations of motion for ideal superfluid	71
C.2 Showing the linear independence, of the first derivative scalar data	75
C.3 Showing the linear independence, of the first derivative vector data	75
C.4 Showing the linear independence of the two derivative parity odd scalar data	76
C.5 Relating the two different entropy current	77
C.6 Computation of the divergence of entropy current	78
C.7 Frame transformation formula	79

1. Introduction

In this paper we work out a complete theory of relativistic superfluid hydrodynamics including dissipative terms at first order in a gradient expansion, allowing for parity violation and the presence of triangle anomalies. In other words we work out the most general form of the equations of 3+1 dimensional s -wave superfluid hydrodynamics consistent with Lorentz invariance and the second law of thermodynamics. While we work in a relativistic context throughout this paper, our final results admit a straightforward non relativistic limit, and are easily formulated in a non relativistic context.

The theory of superfluid hydrodynamics has a long history. The equations of ideal superfluidity (i.e. superfluid dynamics in the absence of dissipative terms) were worked out over 60 years ago by Landau and Tisza [1, 2] in a non relativistic setting. They were generalized to a relativistic superfluid in the early 80's by Israel and Khalatnikov and Lebedev [3, 4, 5] and later reformulated by Carter and Khalatnikov [6, 7] and by Son [8]. In most of this work we will use a formulation of superfluid dynamics close to that of [8]. In a beautiful recent work, Sonner and Withers [9] (see also [10]) have used the equations of Einstein gravity to rederive the Landau-Tisza equations for superfluids that admit a dual gravitational description via the AdS/CFT correspondence of string theory. This development yields independent evidence for the correctness and completeness of the Landau-Tisza theory of ideal superfluidity.

The focus of the current paper is on the one derivative (dissipative) corrections to the Landau-Tisza equations. Dissipative corrections to relativistic superfluids have been extensively studied in the literature [11, 12, 13, 14, 15, 16, 10] and they generalize (and extend) the textbook derivation of such corrections in the non-relativistic limit [17, 18, 19]. While it is straightforward to list the most general dissipative corrections to the

stress tensor, charged current and Josephson relations allowed by Lorentz invariance, it turns out that such a listing also allows for many unphysical possibilities. It is a physical requirement that any hydrodynamic flow be equipped with an entropy current J_S^μ . The second law of thermodynamics requires that the increase in the entropy in any compact, spacelike, region be larger than the incoming entropy flux through the surface of that region; this requires that at every point in spacetime and for every conceivable fluid flow

$$\partial_\mu J_S^\mu \geq 0. \quad (1.1)$$

An interesting and important fact about fluid dynamics, whether superfluid or not, is that the requirement that the divergence of the entropy current always be positive gives rise to important constraints on dissipative corrections to the equations of motion.

As far as we are aware, all previous studies of dissipative corrections to the equations of superfluid hydrodynamics assume, on intuitive grounds, that the entropy current of superfluid hydrodynamics takes a particular canonical form, $J_{S_{\text{canon}}}^\mu$. These studies then determine the dissipative corrections to the energy momentum tensor and charged current (and Josephson condition) consistent with the positivity of the divergence of this canonical entropy current and with covariance under time reversal. The latter restriction is usually called the Onsager relations.

In [20], Son and Surówka observed that, in the presence of triangle anomalies, the entropy current of ordinary fluids (non superfluids) deviates from its canonical form. This observation makes clear that the intuition that fixes the entropy current to its canonical form is not infallible. For this reason, the starting point of our analysis in this paper is the assumption that an entropy current with positive definite divergence exists. However, we make no assumption about the form of this current beyond the requirements of symmetry.

More formally, we allow the (postulated positive divergence) entropy current to take the form

$$J_S^\mu = J_{S_{\text{canon}}}^\mu + \sum_i v_i V_i^\mu \quad (1.2)$$

where $J_{S_{\text{canon}}}^\mu$ is the canonical entropy current referred to above, V_i^μ is a basis of on-shell inequivalent one derivative vectors and v_i are (initially) unconstrained coefficient functions. We then demand that the divergence of J_S^μ be positive semi-definite for any solution of the equations of superfluid hydrodynamics *on an arbitrary background spacetime* and that the Onsager relations are satisfied.¹ These restrictions fix most of

¹Recall that the second law of thermodynamics must apply in any conceivable consistent situation. In particular it must apply when the system is formulated on an arbitrary background spacetime provided the system is free of diffeomorphism anomalies. This condition is true of all experimental superfluids as well as all superfluids obtained via the AdS/CFT correspondence.

the v_i 's in (1.2) and also restrict the possible transport coefficients of the theory.

As we discuss in detail throughout this work, the requirement that the entropy current be of positive divergence *in an arbitrary background spacetime* provides powerful constraints on the form of the entropy current and through it, on the form of the possible dissipative corrections to the hydrodynamic constitutive relations *even in flat space*. For instance, the divergence of the entropy current could contain a term proportional to

$$\nabla_\mu J_S^\mu \propto v_1 R_{\mu\nu} u^\mu u^\nu + \dots \quad (1.3)$$

where u^μ is the fluid velocity, $R_{\mu\nu}$ is the Ricci tensor, and v_1 is one of the coefficient functions in (1.2). The divergence of the entropy current may also contain many other terms proportional to v_i and independent of curvatures. However, for any given fluid flow these other terms can be held fixed while $R_{\mu\nu} u^\mu u^\nu$ is made arbitrarily negative by tuning the curvature tensor.² It follows that the divergence of the entropy current is positive for an arbitrary fluid flow on an arbitrary spacetime only if $v_1 = 0$. Thus, we find a constraint on the entropy current for fluid motion in a flat space background, even though we needed to move to a curved spacetime in order to obtain this constraint.

The result of our extended analysis is as follows. For superfluids which are parity invariant we find that up to a physically unimportant ambiguity (see sections 3 and 4 for details) all the v_i 's in (1.2) should be set to zero—the entropy current agrees with the canonical entropy current.³ This result provides a check of the intuition reviewed in, say, [17], that the entropy current should take its canonical form. It follows that the most general structure of dissipative terms in parity preserving superfluids is given by the 14 parameter family described in [10] which generalized the 13 parameter construction of Clark and Putterman [18, 19]).

Let us now turn to the case of superfluids that do not respect parity symmetry. In this case we find that the entropy current is not constrained to take the canonical form. Not only are the v_i 's non zero, two of them remain undetermined (meaning that they are completely free functions of local thermodynamical variables). In addition there are two physically unimportant ambiguities which we describe in detail in sections 3 and 4.

It turns out that an 18 parameter set of equations of superfluid hydrodynamics are consistent with the positivity of any given choice of entropy current. 14 of these parameters multiply parity invariant structures; these are the parameters that parameterize superfluid hydrodynamics in parity preserving systems. The remaining 4

²Note that curvature tensors do not contribute to the fluid equations at first order, so it is consistent to hold fluid flows fixed while taking curvatures to be very large.

³As a warm up for this analysis we show in section 2 that a similar result is true of parity invariant charged fluid dynamics in the absence of superfluidity

parameters multiply parity odd structures, and are completely new. Thus the most general equations of parity odd superfluid hydrodynamics are parameterized by $14 + 4 + 2$ parameters; the last two parameters being the undetermined v_i s in the entropy current.

At this point it would be useful to clarify some of the terminology we use. We have stated that the equations of motion of superfluid dynamics require the specification of twenty unknown functions of the thermodynamic variables. This may be contrasted with the equations of motion of a normal, charged conformal fluid which requires the specification of two parameters, the shear viscosity and conductivity. For the normal charged conformal fluid, the shear viscosity and conductivity also control the amount of entropy produced by the fluid and so, these parameters are also called dissipative parameters. In the case of parity violating superfluids, five of the twenty parameters are not associated with entropy production so, only fifteen of the twenty parameters are dissipative in the sense described above. In normal charged conformal fluids the conductivity controls not only the response of the current to an external electric field, but also the response of the current to changes in temperature and changes in chemical potential. The situation in superfluids is similar: the twenty parameters, and in particular the two undetermined parameters associated with the entropy current, control the response of the system to various changes in the hydrodynamic variables.

When the superfluid velocity is too large superfluidity breaks down. For this reason, when considering experimentally accessible superfluids such as liquid Helium it is often particularly interesting to study dissipative corrections in the limit that the normal and superfluid velocities are collinear. In this limit the 14 parameter family of dissipative corrections of parity invariant superfluids reduce to the five parameter family described, for instance, in the classic text book of Landau and Lifshitz [17].⁴ When parity is violated, the twenty parameter family of solutions reduces to seven. The two additional parameters determine (in an appropriate frame) the response of the charged current to a magnetic field and to “chiral vorticity”, i.e., changes in $\omega^\mu = \frac{1}{2}\epsilon^{\mu\nu\alpha\beta}u_\nu\partial_\alpha u_\beta$.

When the superfluid density is set to zero we are in the normal phase. In this limit the two undetermined parameters of the superfluid theory reduce to an integration constant and the contributions to the charge current proportional to the magnetic field and chiral vorticity are completely determined by the triangle anomaly [20] (up to an integration constant). In such a limit, we recover the results of [21, 22, 20] (see also [23]).

We emphasize that, as opposed to the configuration at zero superfluid density, the 6 new arbitrary parameters that appear in parity non invariant superfluid hydrodynamics

⁴These reduce further to only three parameters in the case of a conformally invariant theory.

may be non vanishing even in the absence of an anomaly. This difference may be of significance; practically speaking, anomalies are intrinsically relativistic phenomena and always vanish in a nonrelativistic setting. Consequently, the results of this paper suggest that a term in the particle number current proportional to the vorticity could possibly show up in non relativistic table top experiments involving superfluids or superconductors which violate parity (e.g. non centro symmetric superconductors).

We were led to the study of dissipative effects in superfluids in order to understand the results of certain holographic computations using the AdS/CFT correspondence of string theory. We emphasize, however, that our eventual derivation of the general equations of superfluid hydrodynamics makes no use of the AdS/CFT correspondence and so, applies to all superfluids, not just those that admit a dual gravitational description.

In the second part of our paper we test the collinear limit of our general formalism using the AdS/CFT correspondence. In the case of parity preserving superfluids, the authors of [16, 10] showed that it is possible to use the fluid gravity map [24, 25, 26, 27, 28, 21, 22, 29] to derive dissipative corrections to the equations of superfluid dynamics by extending the analytically tractable superfluid model of [30] which is valid close to the phase transition point. The constitutive relations so obtained fit within the 14 parameter fluid dynamical framework spelt out in [10] which generalize the text book predictions of [18, 19]). In particular, it was explicitly verified that the entropy current obtained by holographic methods matches the canonical entropy current in agreement with the intuition of [17].

In this paper we consider a generic asymptotically AdS_5 gravitational system which involves a Chern Simons term for a single bulk $U(1)$ field. Such a Chern Simons term signifies the presence of a $U(1)^3$ triangle anomaly in the dual field theory and so, in particular, introduces parity violating terms into the effective superfluid dynamical description. Particular truncations of type IIB supergravity for which our analysis is valid can be found in [31, 32]. We find that, in the collinear limit, it is possible to obtain integral expressions for all the parity odd transport coefficients in terms of the background solution. Our results are in perfect agreement with the collinear limit of the modified theory of parity violating superfluid dynamics which we present in section 4. We regard this agreement as a nontrivial (though as yet limited) check of the general results of section 4. It should be possible to use the AdS/CFT correspondence away from the collinear limit to find much more extensive test our results; we leave this to future work.

Note Added: After the work presented in this paper was completed, we received a paper [33] that has substantial overlap with section 4 of this paper. While the approach of [33] is similar to the one adopted in this paper, our final results differ in several qualitative and quantitative respects. We present a brief comparison with [33]

in section 5.

2. The theory of charged fluid dynamics

In this pedagogical section we construct the most general equations of Lorentz invariant charged fluid dynamics consistent with the second law of thermodynamics. Our goal is to illustrate our method for determining the most general form of fluid-dynamical equations of motion in a simple and familiar context before tackling the slightly more complicated case of superfluids. The final results of this section are well known; the novelty of this section lies in our method of computation.

The long-wavelength degrees of freedom of a locally equilibrated system with a single global $U(1)$ charge can be taken to be the velocity field $u_\mu(x)$ (normalized so that $u^\mu u_\mu = -1$), the temperature field $T(x)$ and a chemical potential field $\mu(x)$. Both the energy momentum tensor and the charged current can be written in terms of these five fields and their gradients. The equations of motion of charged fluid dynamics are the conservation of the stress tensor and charge current

$$\begin{aligned}\nabla_\mu T^{\mu\nu} &= F^{\nu\mu} J_\mu \\ \nabla_\mu J^\mu &= -\frac{c}{8} \epsilon^{\mu\nu\rho\sigma} F_{\mu\nu} F_{\rho\sigma}\end{aligned}\tag{2.1}$$

which provides the five equations for the five hydrodynamic fields. In these equations we have allowed for the possibility that the current in question has a $U(1)^3$ anomaly. We call the coefficient c the anomaly coefficient. We have also allowed the current to be coupled to an external source with field strength $F_{\mu\nu}$. To completely determine the equations of motion it remains to determine the dependence of $T^{\mu\nu}$ and J^μ on the fields $u^\mu(x)$, $T(x)$, $\mu(x)$ and their derivatives.

By considering a stationary fluid for which $u^\mu = (1, 0, 0, 0)$ and using boost invariance one can argue that the stress tensor and charge current take the form

$$\begin{aligned}T^{\mu\nu} &= (\rho + P)u^\mu u^\nu + P\eta^{\mu\nu} + T_{diss}^{\mu\nu} \\ J^\mu &= qu^\mu + J_{diss}^\mu\end{aligned}\tag{2.2}$$

where $T_{diss}^{\mu\nu}$ and J_{diss}^μ are the contributions to the stress tensor and charge current that involve derivatives of μ , T and u^μ . The equations that express $T_{diss}^{\mu\nu}$ and J_{diss}^μ in terms of fluid dynamical fields and their derivatives are termed constitutive relations. In the long wavelength fluid dynamical limit it is sensible to expand the constitutive relations in powers of derivatives of the fluid dynamical fields u^μ , T and μ . We will refer to such an expansion as a derivative expansion and refer to the terms which are linear in gradients as first order terms. In this paper we work only to first order in the

derivative expansion. The electromagnetic source term $F^{\mu\nu}$ is taken to be of first order in derivatives in this counting.

Field Redefinitions and frame choices

Note that the fluid temperature T , chemical potential μ and velocity u^μ are thermodynamical concepts that are well defined in equilibrium but have no microscopic definitions in dynamical situations. In other words, we are always free to redefine the thermodynamic variables into primed ones according to the equations

$$\begin{aligned} u^\mu &= u'^\mu + \delta u^\mu \\ T &= T' + \delta T \\ \mu &= \mu' + \delta \mu \end{aligned} \tag{2.3}$$

where δu^μ is an arbitrary one derivative vector that obeys $\delta u^\mu u_\mu = \delta u^\mu u'_\mu = 0$ and δT and $\delta \mu$ are arbitrary one derivative scalars. The primed and unprimed fields are each equally good definitions of the velocity, temperature and chemical potential fields. Physically meaningful assertions, such as the constitutive relations for $T_{diss}^{\mu\nu}$ and J_{diss}^μ , must only involve field redefinition invariant quantities.

Thus, let us determine the field redefinition invariant combinations of $T_{diss}^{\mu\nu}$ and J_{diss}^μ . Under the field redefinition (2.3),

$$\begin{aligned} \delta T_{diss}^{\mu\nu} &= (u^\mu \delta u^\nu + u^\nu \delta u^\mu)(P + \rho) + u^\mu u^\nu d(P + \rho) + \eta^{\mu\nu} dP \\ \delta J_{diss}^\mu &= q \delta u^\mu + dqu^\mu \end{aligned} \tag{2.4}$$

where

$$\begin{aligned} \delta T_{diss}^{\mu\nu} &= T_{diss}'^{\mu\nu} - T_{diss}^{\mu\nu} \\ \delta J_{diss}^\mu &= J_{diss}'^\mu - J_{diss}^\mu \end{aligned}$$

and $df(\mu, T)$ represents the change in the function f under the first order variable change (2.3). It useful to decompose $T_{diss}^{\mu\nu}$ and J_{diss}^μ into $SO(3)$ invariant tensors, vectors and scalars. The $SO(3)$ that we are referring to is the group of rotations orthogonal to u^μ . To this end we introduce the projection matrix

$$P^{\mu\nu} = \eta^{\mu\nu} + u^\mu u^\nu. \tag{2.5}$$

We find that there is one tensor, two vectors and three scalars. The unique tensor

$$P^\mu_\alpha P^\nu_\beta T_{diss}^{\alpha\beta} - \frac{P^{\mu\nu}}{3} P_{\alpha\beta} T_{diss}^{\alpha\beta} \tag{2.6}$$

is automatically field redefinition invariant. The two vectors $P^\mu_\alpha T_{diss}^{\alpha\beta} u_\beta$ and $P^\mu_\alpha J^\alpha$ transform under field redefinitions as

$$\begin{aligned}\delta\left(P^\mu_\alpha T_{diss}^{\alpha\beta} u_\beta\right) &= -(P + \rho)\delta u^\mu \\ \delta\left(P^\mu_\alpha J^\alpha\right) &= q\delta u^\mu\end{aligned}\tag{2.7}$$

so that the unique invariant combination of vectors is given by

$$P^\mu_\alpha J^\alpha + \frac{q}{P + \rho}\left(P^\mu_\alpha T_{diss}^{\alpha\beta} u_\beta\right).\tag{2.8}$$

The three scalars transform under field redefinitions as

$$\begin{aligned}\delta\left(P_{\alpha\beta} T_{diss}^{\alpha\beta}\right) &= 3dP \\ \delta\left(u_\alpha T_{diss}^{\alpha\beta} u_\beta\right) &= dq \\ \delta\left(u_\alpha J^\alpha\right) &= -dq\end{aligned}\tag{2.9}$$

so that the unique invariant scalar is given by

$$\frac{1}{3}\left(P_{\alpha\beta} T_{diss}^{\alpha\beta}\right) - \frac{\partial P}{\partial \rho}\left(u_\alpha T_{diss}^{\alpha\beta} u_\beta\right) + \frac{\partial P}{\partial q}\left(u_\alpha J^\alpha\right)\tag{2.10}$$

where $\frac{\partial P}{\partial \rho}$ is taken at constant q and $\frac{\partial P}{\partial q}$ is taken at constant ρ .

Instead of working in a manifest field redefinition invariant manner, it is sometimes convenient to ‘fix’ the field redefinition ambiguity by imposing five additional conditions on the thermodynamic fields so that they are well defined. Different choices of fixing the ambiguity are referred to as frames. One often used frame is the so called Landau frame, in which the velocity and temperature fields are defined to obey the conditions $T_{diss}^{\mu\nu} u_\nu = 0$ and $J_{diss}^\mu u_\nu = 0$. This gives one vector and two scalar conditions, matching the field redefinition degrees of freedom. Another choice of frame is the Eckart frame which is defined by the conditions $J_{diss}^\mu = 0$ together with $u_\mu T_{diss}^{\mu\nu} u_\nu = 0$. The expressions for the invariant vector (2.8) and the invariant scalar (2.10) greatly simplify in either of these frames. In this paper we adopt no such ‘gauge’ choice but work in a fully field redefinition invariant manner.

The strategy for the rest of this section

In order to complete our specification of the equations of charged fluid dynamics, we need to specify $T_{diss}^{\mu\nu}$ and J_{diss}^μ (or more precisely the field redefinition invariant parts (2.6), (2.8) and (2.10) of these expressions) as a function of first derivatives of fluid dynamical fields. Of course, in any particular dynamical system, the explicit form

of the constitutive relations for $T_{diss}^{\mu\nu}$ and J_{diss}^μ can be determined only by a detailed dynamical computation. In this paper we will be interested not in computing the precise form of these quantities in any particular system, but in parameterizing the most general form that the constitutive relationship can take in any system. As we will see below, it will prove possible to completely determine the form of the first order constitutive relations up to three undetermined dissipative parameters, each of which is an arbitrary function of T and μ .

We proceed as follows. As in any effective field theory, we start by writing down all possible expressions which may contribute to $T_{diss}^{\mu\nu}$ and J_{diss}^μ . We then eliminate those that do not satisfy the symmetries of the theory, Lorentz invariance in this case. In addition, since we are dealing with a hydrodynamic theory, we must ensure that the second law of thermodynamics is satisfied. As explained in the introduction, we demand the existence of an entropy current of positive semi-definite divergence even when the theory is formulated on a curved background. As alluded to in section 1, the entropy current is defined to be a four vector J_S^μ satisfying two requirements. The first is that in a configuration where the fluid is in uniform motion,⁵

$$J_S^\mu = su^\mu \quad (\text{for a spacetime independent configuration}) \quad (2.11)$$

with s the entropy density which is related to ρ , P , q , μ and T through

$$\rho + P = sT + \mu q. \quad (2.12)$$

Our second requirement of the entropy current is that its divergence is positive semi-definite in an arbitrary curved background,

$$\nabla_\mu J_S^\mu \geq 0 \quad (2.13)$$

implying that the entropy increase in any region is always greater than the entropy inflow into that region.

For a perfect fluid level (i.e. a fluid in which all gradient terms have been neglected— $T_{diss}^{\mu\nu} = J_{diss}^\mu = 0$) the entropy current is given by (2.11). At this order it is not difficult to verify that $\nabla_\mu(su^\mu) = 0$ using (2.12) and $dP = sdT + qd\mu$.

Once the gradients of u^μ , T and μ/T are non vanishing the divergence of the entropy current no longer vanishes. Indeed, the divergence of the entropy current at the one derivative level will be the focus of much of the rest of this paper. We will demand, on

⁵Such a configuration is a stationary, dissipation free solution to the equations of fluid dynamics. Indeed it may be obtained by boosting a uniform fluid at rest (by which we mean a uniform fluid with velocity field $u^\mu = (1, 0, 0, 0)$).

physical grounds, that it is possible to modify (2.11) by first order corrections so that (2.13) will be satisfied. This requirement will turn out to constrain the possible forms of $T_{diss}^{\mu\nu}$ and J_{diss}^μ .

We start our analysis in section 2.1 by considering parity conserving charged fluids. In section 2.2 we move on to describe parity-violating fluid dynamics.

2.1 Parity Invariant Charged Fluid Dynamics

Consider a hydrodynamic theory in the presence of external electromagnetic fields satisfying (2.1) with $c = 0$. Following the general prescription described at the beginning of this section, we would like to write the most general parity-invariant and Lorentz invariant contributions to J_{diss}^μ , $T_{diss}^{\mu\nu}$ and J_S^μ which involves a single derivative of the hydrodynamic fields u^μ , T and μ . This is carried out in section 2.1.1. We then work out the the restrictions on these terms by requiring that the entropy current has positive semi-definite divergence. This is described in section 2.1.2.

2.1.1 Classification of one and two derivative data

We begin our analysis on a technical point. The tangent space about any point in our spacetime manifold has an $SO(3, 1)$ rotational invariance. However, the fluid velocity vector, $u^\mu(x)$, takes a definite value at that point and breaks this rotational group down to $SO(3)$. It is useful to decompose all derivatives of fluid dynamical fields, at any given spacetime point, into representations of this residual $SO(3)$ rotational group.

In the first column of table 1 we have classified all expressions formed from a single derivative of any of $u^\mu(x)$, $T(x)$ and $\mu(x)$ according to their $SO(3)$ and parity transformation properties. We refer to these expressions as one derivative fluid dynamical data. We have also classified one derivative expressions constructed out of the background electromagnetic fields according to their $SO(3)$ and parity transformation properties. We will refer to these as background data. As fluid and background field data enter our analysis on an even footing, we have listed these expressions together in the first column of table 1.⁶

Not all the expressions in the first column of table 1 are independent under the equations of motion. The equations of motion can be used to solve for some pieces of data in terms of other data. The classification of the equations of motion according to their $SO(3)$ and parity transformation properties can be found in the middle column of table 1. Note that there are no tensor equations of motion.

⁶Since all curvature invariants built out of the background metric have at least two derivatives, there is no one derivative data associated with the metric.

In the last column of table 1 we have listed a choice of independent data. By this we mean a choice of independent one derivative fluid dynamical expressions and one derivative field expressions in terms of which all others can be solved for.

$SO(3)$ and P classification	All data	Equations of motion	Independent data
Scalars	$u^\mu \partial_\mu T$ $u^\mu \nabla_\mu \frac{\mu}{T}$ $\partial_\mu u^\mu$	$u_\mu \nabla_\nu T^{\mu\nu} = 0$ $\nabla_\mu J^\mu = 0$	$S_1 = \partial_\mu u^\mu$
Vectors	$P^{\mu\nu} \partial_\nu T$ $u^\nu \partial_\nu u^\mu$ $P^{\mu\nu} \partial_\nu \frac{\mu}{T}$ $F^{\mu\nu} u_\nu$	$P^\mu{}_\nu \partial_\rho T^{\rho\nu} = 0$	$V_1 = -P^{\mu\nu} \partial_\nu \frac{\mu}{T} + \frac{F^{\mu\nu} u_\nu}{T}$ $V_2 = u^\nu \nabla_\nu u^\mu$ $V_3 = F^{\mu\nu} u_\nu$
Tensors	$\sigma_{\mu\nu}$	–	$T_1 = \sigma_{\mu\nu}$
Pseudo vectors	$\frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu \partial_\alpha u_\beta$ $\frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu F_{\alpha\beta}$	–	$\omega^\mu = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu \partial_\alpha u_\beta$ $B^\mu = \frac{1}{2} \epsilon^{\mu\nu\alpha\beta} u_\nu F_{\alpha\beta}$

Table 1: One-derivative expressions classified according to their transformation laws under the $SO(3)$ residual symmetry and parity. The first column lists all one derivative data. The second column lists the equations of motion. The last column lists a choice of independent data. See (2.14) and (2.15) for the definition of $\sigma_{\mu\nu}$ and $P_{\mu\nu}$ respectively.

While some of the expressions used in table 1 such as

$$\sigma_{\mu\nu} = \frac{1}{2} P^{\mu\alpha} P^{\nu\beta} (\nabla_\alpha u_\beta + \nabla_\beta u_\alpha - P_{\alpha\beta} (\nabla_\lambda u^\lambda)) \quad (2.14)$$

and

$$P^{\mu\nu} = u^\mu u^\nu + \eta^{\mu\nu} \quad (2.15)$$

are standard, some of our notation isn't. The new notation has been introduced in order to prepare the reader for later sections. In particular V_3 is the electric field in the rest frame of a fluid element. In the conventions of Son and Surówka [20] we have

$$V_3 = E_\mu. \quad (2.16)$$

We will soon construct an entropy current that includes terms which are first order in derivatives. The divergence of such an entropy current is of second order in derivatives and includes terms quadratic in first order fluid (and background field) data plus expressions built out of two derivatives acting on fluid fields or single derivatives

of electromagnetic field strengths. We refer to the second class of expressions as two derivative scalar data. When studying the divergence of the entropy current it is useful to have a listing of independent scalar two derivative data.

In the first column of table 2 we list the most general fluid and background field (but not curvature related) two derivative data that transforms as an $SO(3)$ scalar. More explicitly, we list all scalar expressions formed by acting with two derivatives on $u^\mu(x)$, $T(x)$ and $\mu(x)$ together with all scalars formed from the action of a single derivative on electromagnetic field strengths.⁷ In the second column of the same table we list all scalar two derivative equations of motion. In the last column of the same table we list our choice of independent two derivative scalar data (in terms of which we have solved for all the other two derivative scalars).

All data	Equations of motion	Independent data
$u^\mu u^\nu \partial_\mu \partial_\nu T$	$\nabla_\mu \nabla_\nu T^{\mu\nu} = 0$	$P^{\mu\nu} \nabla_\mu \partial_\nu \frac{\mu}{T}$
$P^{\mu\nu} \partial_\mu \partial_\nu T$		
$u^\mu u^\nu \partial_\mu \partial_\nu \left(\frac{\mu}{T}\right)$	$P^\rho_\nu \nabla_\rho \nabla^\mu T^{\mu\nu} = 0$	$u^\mu \nabla_\mu \partial_\nu u^\nu$
$P^{\mu\nu} \partial_\mu \partial_\nu \left(\frac{\mu}{T}\right)$		
$u^\mu \partial_\mu \partial_\nu u^\nu$	$u^\nu \nabla_\nu \nabla^\mu J^\mu = 0$	$\nabla_\mu (F^{\mu\nu} u_\nu)$
$\partial_\mu (F^{\mu\nu} u_\nu)$		

Table 2: Parity even two derivative scalar data for charged fluids. The first column lists all six second order scalars constructed from two derivatives of the hydrodynamic variables and background field strengths. The second column lists the three scalar two derivative equations of motion. The last column lists one choice of a $6-3 = 3$ dimensional basis for the independent two derivative scalar data.

2.1.2 The general entropy current and its divergence

Armed with the listings in tables 1 and 2 we now proceed with our analysis. Traditional studies of first order charged fluid dynamics (see, for example, [17]) assume that the entropy current takes a canonical form,⁸

$$J_{S \text{ canon}}^\mu = s u^\mu - \frac{1}{T} u_\mu T_{diss}^{\mu\nu} - \frac{\mu}{T} J_{diss}^\mu. \quad (2.17)$$

⁷It is also easy to list two derivative fluid data in the 3, 5 and 7 dimensional representations of $SO(3)$, but that will not be required in what follows, so we do not present such a listing.

⁸As explained in [10] the expression in (2.17) is frame invariant, i.e. invariant under a first order field redefinition of T , u^μ and μ . Note that the second term on the right hand side vanishes in the Landau frame while the third term vanishes in the Eckart frame.

As we explained in the introduction, in this work we will not make any prior assumption about the form of the entropy current. According to the analysis of section 2.1.1 the most general parity even first order entropy current is given by

$$J_S^\mu = J_{S\text{ canon}}^\mu + s_1 S_1 u^\mu + \sum_{i=1}^3 v_i V_i^\mu \quad (2.18)$$

where S_1 and V_i are defined in the last column of table 1, and s_1 and the v_i 's are arbitrary functions of $\frac{\mu}{T}$ and T .

We now explore the constraints obtained by enforcing the positivity of the divergence of the entropy current (2.18). It is easily demonstrated (see, for instance, [17, 10]) that the divergence of the canonical part of the entropy current is given by

$$\nabla_\mu J_{S\text{ canon}}^\mu = -\nabla_\mu \left(\frac{u_\nu}{T} \right) T_{diss}^{\mu\nu} - \left(\partial_\mu \left(\frac{\mu}{T} \right) - \frac{F_{\mu\nu} u^\nu}{T} \right) J_{diss}^\mu. \quad (2.19)$$

The right hand side of (2.19) is a quadratic form in one derivative fluid and background electromagnetic field data. The divergence of the non canonical part of the entropy current in (2.18) is also a two derivative expression but is composed of two kinds of terms. The first set of terms are linear in independent two derivative and curvature data. Such terms are always inconsistent with the positivity of the entropy current, and so we must choose s_1 and v_i so that these terms vanish. The second set of terms contains products of one derivative terms. Such terms would modify the quadratic form on the right hand side of (2.19) and do not necessarily vanish. Schematically, we have

$$\partial_\mu J_S^\mu = \left(\begin{array}{c} \text{independent two} \\ \text{derivative and curvature data} \end{array} \right) + \left(\begin{array}{c} \text{quadratic form in} \\ \text{first order data} \end{array} \right). \quad (2.20)$$

The first term on the right hand side of (2.20) must vanish while the second term must be tuned to be positive.

2.1.3 Constraints from positivity of the divergence of the entropy current

We will first explore the constraints that follows from the requirement that no two derivative data appears in the divergence of the entropy current. As explained previously, this implies that the first set of terms on the right hand side of (2.20) must vanish. We will implement this condition separately for two derivative and curvature terms.

Constraints from the vanishing of 2 derivative terms

The two derivative part of the divergence of the entropy is given by

$$-v_1 P^{\mu\nu} \nabla_\mu \partial_\nu \frac{\mu}{T} + (s_1 + v_2) u^\mu \nabla_\mu \partial_\nu u^\nu + \left(v_3 + \frac{v_1}{T} \right) \nabla_\mu (F^{\mu\nu} u_\nu).$$

This expression is a linear combination of the three independent two derivative pieces of data (see Table 2). It follows that the vanishing of two derivative terms requires us to set the coefficients of each of these terms to zero, i.e. to set $v_1 = v_3 = 0$ and $v_2 = -s_1$. Thus the vanishing of two derivative terms in the divergence of the entropy current restricts the entropy current (2.18) to take the form

$$J_S^\mu = J_{S\text{ canon}}^\mu + s_1 (S_1 u^\mu - V_2^\mu) . \quad (2.21)$$

where s_1 is still an arbitrary function of T and μ .

Constraints from vanishing of curvature terms

According to (2.21) the entropy current has a one parameter ambiguity, s_1 . Were we to restrict our attention to a flat space background we would not have been able resolve this ambiguity. Consider a charged fluid propagating on an arbitrary curved background. The cancellation of two derivative terms proportional to s_1 is now incomplete; it is not difficult to check that there is an additional, curvature dependent term in the divergence of the entropy current proportional to $s_1 R_{\alpha\beta} u^\alpha u^\beta$ with $R_{\alpha\beta}$ the Ricci tensor. This term is inconsistent with positivity of the divergence of J_S^μ . Thus, we are forced to set $s_1 = 0$.

We conclude that the requirement that the divergence of the entropy current is positive *in an arbitrary curved background* forces the entropy current to take its canonical form, justifying the assumptions of standard treatments of fluid dynamics e.g. [17].

2.1.4 Constraints on dissipative terms

We have demonstrated that the entropy current takes its canonical form and consequently that its divergence is given by (2.19). It is now not difficult to work out the constraints on dissipative terms that ensure the positivity of the quadratic form on the right hand side of (2.19). We outline the calculation here.

Consider the expansion of $\nabla_\mu \left(\frac{u^\mu}{T} \right)$ and $-V_1 = \partial_\mu \frac{\mu}{T} - \frac{E_\mu}{T}$ which appear on the right hand side of (2.19) into $SO(3)$ invariant tensors vectors and scalars. We find a single tensor, $\sigma_{\mu\nu}$, two vectors,

$$V_1^\mu, \quad \text{and} \quad \left(P^{\mu\nu} \frac{\partial_\nu T}{T} + (u \cdot \partial) u^\mu \right)$$

(see table 1 for a definition of the vector V_1^μ) and three scalars,

$$\frac{(u \cdot \partial) T}{T}, \quad (u \cdot \partial) \nu, \quad \text{and} \quad (\nabla \cdot u) .$$

While the two vectors are completely distinct off-shell, it turns out that the equations of motion imply that they are proportional to each other on-shell. Similarly, the equations

of motion imply that the three scalars are also proportional to each other on-shell. As we demonstrate in Appendix B the explicit relations are

$$\begin{aligned}
\frac{(u.\partial)T}{T} &= - \left[\frac{\partial P}{\partial \rho} \right]_q (\nabla \cdot u) \\
(u.\partial)\nu &= -\frac{1}{T} \left[\frac{\partial P}{\partial q} \right]_\rho (\nabla \cdot u) \\
P^{\mu\nu} \frac{\partial_\nu T}{T} + (u.\partial)u^\mu &= \frac{qT}{\rho + P} V_1^\mu.
\end{aligned} \tag{2.22}$$

Plugging these relations into (2.19), we can rewrite the divergence of the entropy current in the form

$$\begin{aligned}
\nabla_\mu J_S^\mu &= - \frac{(\nabla_\mu u^\mu)}{T} \left[\frac{(T_{diss})_{ab} P^{ab}}{3} - \frac{\partial P}{\partial \rho} (u_\mu u_\nu T_{diss}^{\mu\nu}) + \frac{\partial P}{\partial q} (u_\mu J_{diss}^\mu) \right] \\
&\quad + V_{1\mu} \left[J_{diss}^\mu + \frac{q}{\rho + P} (u_\nu T_{diss}^{\mu\nu}) \right] - \frac{T_{diss}^{\mu\nu} \sigma_{\mu\nu}}{T}
\end{aligned} \tag{2.23}$$

where V_1^μ , B^μ and E^μ were defined in Table 1 and (2.16). We collect their definitions here for convenience:

$$\begin{aligned}
E_\mu &= F_{\mu\nu} u^\nu \\
B_\mu &= \frac{1}{2} \epsilon_{\mu\nu\alpha\beta} u^\nu F^{\alpha\beta} \\
V_{1\mu} &= \frac{E_\mu}{T} - P_\mu^\theta \partial_\theta \nu.
\end{aligned}$$

We will now use (2.23) to constrain the constitutive relations of fluid dynamics, i.e. the expressions for $T_{diss}^{\mu\nu}$ and J_{diss}^μ as a linear expansion in first order scalars, vectors and tensors. To first order in gradients there is only one independent scalar data so the scalar parts of $T_{diss}^{\mu\nu}$ and J_{diss}^μ are necessarily proportional to $\nabla \cdot u$. The vector parts of $T_{diss}^{\mu\nu}$ and J_{diss}^μ must each be expanded as a linear sum of the three independent vectors listed in Table 1. The tensor in Table 1 is proportional to $\sigma_{\mu\nu}$ since there is only one $SO(3)$ invariant tensor. It follows from group theory that positivity of the divergence of the entropy current implies positivity of the scalar, vector and tensor components separately. Thus, we have

$$\begin{aligned}
P_\alpha^\mu P_\beta^\nu T_{diss}^{\alpha\beta} - \frac{P^{\mu\nu}}{3} P_{\alpha\beta} T_{diss}^{\alpha\beta} &= -\eta \sigma^{\mu\nu} \\
P_\alpha^\mu \left(J_{diss}^\alpha + \frac{q}{\rho + P} (u_\nu T_{diss}^{\alpha\nu}) \right) &= \kappa V_1^\mu \\
\frac{(T_{diss})_{ab} P^{ab}}{3} - \frac{\partial P}{\partial \rho} (u_\mu u_\nu T_{diss}^{\mu\nu}) + \frac{\partial P}{\partial q} (u_\mu J_{diss}^\mu) &= -\beta \partial_\alpha u^\alpha
\end{aligned} \tag{2.24}$$

where

$$\eta \geq 0, \quad \kappa \geq 0, \quad \beta \geq 0.$$

These three coefficients are the shear viscosity, η , the heat conductivity, κ , and the bulk viscosity, β . The bulk viscosity is traditionally denoted by ζ but in this work we reserve ζ for different use.⁹

Several aspects of (2.24) deserve comment. First, the requirement of positivity does not individually constrain the three scalar and two vector pieces in $T_{diss}^{\mu\nu}$ and J_{diss}^μ , but only constrains the combinations that appear in (2.10) and (2.8). This is exactly as we would expect: only field redefinition invariant data can be constrained in a physical way. The vectors and scalars that are left undetermined are unphysical; they can be changed, or chosen arbitrarily, by a field redefinition. Despite appearances, (2.24) constitutes a complete determination of the constitutive relations of our system.

We also note that we could have used the fact that the divergence of the entropy current is frame invariant (see [10]) to determine the frame invariant scalar, vector and tensor combinations in (2.6), (2.8) and (2.10); the expression on the right hand side of (2.23) must arrange itself into such frame invariant combinations.

The third aspect to note is that the constraints of positivity are relatively mild in the scalar and tensor sector. The expansion of scalars and tensors is the most general one permitted by symmetry; the requirement of positivity merely imposes inequalities in the coefficients of this expansion. However, the constraint on vectors is much stronger. Symmetry alone would have allowed the expansion of the second line in (2.24) as an arbitrary linear combination of the 3 vectors V_1 , V_2 and V_3 . However the requirement of positivity sets the coefficients V_2 and V_3 to zero,¹⁰ apart from imposing an inequality on the coefficient of the third. We will see this pattern repeated and magnified in the study of superfluid dynamics in sections 3 and 4 in the scalar, vector and tensor sector.

2.2 Parity non invariant charged fluid dynamics

Let us now turn to the dynamics of fluids that are not invariant under parity transformations. According to table 1 we should allow the entropy current to depend on an additional arbitrary pseudo vector. Thus, the most general entropy current for such a

⁹Note that the speed of sound, c_s , is related to the variation of the pressure with respect to energy density through $\frac{\partial P}{\partial \rho} = c_s^2$. Using dimensional analysis one can conclude that $\frac{\partial P}{\partial q} = 0$ in a scale invariant theory. It then becomes clear that in a conformal theory the left hand side of the last equality in (2.24) vanishes as it should.

¹⁰The origin of this constraint is the observation that the quadratic form $ax^2 + bxy + cxz$ is positive only when $b = c = 0$ and $a \geq 0$. The role of x is played by the vector V , while the roles of y and z are played by the other two vectors

fluid takes the form

$$J_S^\mu = J_{S\text{ canon}}^\mu + s_1 S_1 u^\mu + \sum_{i=1}^3 v_i V_i^\mu + \sigma_\omega \omega^\mu + \sigma_B B^\mu. \quad (2.25)$$

In the parity even sector the divergence of this entropy current is identical to the one discussed in subsection 2.1; the arguments in 2.1 go through unchanged and in particular the cancellation of two derivative and scalar terms set $s_1 = v_i = 0$. In the parity odd sector the divergence of the entropy current receives contributions involving the dot product of the pseudo vectors ω^μ and B^μ with ordinary vectors. Positivity of the divergence of the entropy current implies that such products vanish.¹¹ This restriction was analyzed in detail by Son and Surówka [20] who found that it leads to

$$\begin{aligned} P_\alpha^\mu P_\beta^\nu T_{diss}^{\alpha\beta} - \frac{P^{\mu\nu}}{3} P_{\alpha\beta} T_{diss}^{\alpha\beta} &= -\eta \sigma^{\mu\nu} \\ P_\alpha^\mu \left(J_{diss}^\alpha + \frac{q}{\rho + P} (u_\nu T_{diss}^{\alpha\nu}) \right) &= \kappa V^\mu + \tilde{\kappa}_\omega \omega^\mu + \tilde{\kappa}_B B^\mu \\ \frac{(T_{diss})_{ab} P^{ab}}{3} - \frac{\partial P}{\partial \rho} (u_\mu u_\nu T_{diss}^{\mu\nu}) + \frac{\partial P}{\partial q} (u_\mu j_{diss}^\mu) &= -\beta \partial_\alpha u^\alpha \end{aligned} \quad (2.26)$$

where

$$\begin{aligned} \sigma_\omega &= c \frac{\mu^3}{3T} + T\mu k_2 + T^2 k_1 \\ \sigma_B &= c \frac{\mu^2}{2T} + \frac{T}{2} k_2 \\ \tilde{\kappa}_\omega &= c \left(\mu^2 - \frac{2}{3} \frac{q}{\rho + P} \mu^3 \right) + T^2 \left(1 - \frac{2q}{\rho + P} \mu \right) k_2 - \frac{2q}{\rho + P} k_1 \\ \tilde{\kappa}_B &= c \left(\mu - \frac{1}{2} \frac{q}{\rho + P} \mu^2 \right) - \frac{T^2}{2} \frac{q}{\rho + P} k_2 \end{aligned} \quad (2.27)$$

and k_1 and k_2 are integration constants. We will now argue that the requirement of CPT invariance forces k_2 to vanish.¹² The argument goes as follows. Consider the CPT transformation $x^\mu \rightarrow -x^\mu$, $q \rightarrow -q$ (and so $\mu \rightarrow -\mu$). Under this transformation $T_{diss}^{\mu\nu} \rightarrow T_{diss}^{\mu\nu}$ and $J_{diss}^\mu \rightarrow -J_{diss}^\mu$. Also $u^\mu \rightarrow -u^\mu$ so that $\omega^\mu \rightarrow -\omega^\mu$ and $B^\mu \rightarrow B^\mu$. Thus under a CPT transformation it must be that $\tilde{\kappa}_\omega \rightarrow \tilde{\kappa}_\omega$ while $\tilde{\kappa}_B \rightarrow -\tilde{\kappa}_B$. Consistency of this requirement with (2.27) sets $k_1 = 0$. Nothing in our argument

¹¹We will see later that products of vectors and pseudo vectors do not necessarily need to vanish in the case of superfluid dynamics. In the current setup vanishing of such bilinear terms follows from the fact that the divergence has no squares of pseudo vectors and contains only a single squared vector.

¹²We thank D. Son for pointing this out to us.

requires that k_2 vanish (although it would be interesting to find a specific system with $k_2 \neq 0$; k_2 vanishes in all AdS/CFT computations performed so far).

The results (2.26) and (2.27) have several interesting features. First, the presence of an anomaly forces the entropy current to depart from the canonical form (i.e. σ_B and σ_ω are never zero if c is nonzero). Second, it induces new terms in the vector part of the constitutive relations, proportional to the vorticity and the magnetic field. Third, the new contributions to both the entropy current and the vector part of the constitutive relations are completely determined (up to an integration constant that is independent of T and μ) by the anomaly. In other words, although the constitutive relations take a different form from the parity even case, this change in form is completely determined by the anomaly, and we have no new free parameters apart from the integration constant k_2 .

3. Parity invariant Superfluid hydrodynamics

By definition, a superfluid is a fluid phase of a system with a spontaneously broken global symmetry. When discussing superfluids this forces us to consider the gradient of the Goldstone boson as an extra hydrodynamical degrees of freedom in addition to the standard variables u^μ , T and μ . More precisely, if we denote the Goldstone Boson by ψ (ψ is the phase of the condensate of the charged scalar operator) and we also wish turn on a background gauge field A_μ then

$$\xi_\mu = -\partial_\mu \psi + A_\mu \tag{3.1}$$

represents the covariant derivative of the Goldstone Boson and is an extra hydrodynamic degree of freedom.¹³ According to the Landau-Tisza two fluid model the superfluid should be thought of as a two component fluid: a condensed component and a non condensed or normal component. The velocity field of the normal fluid is given by u_μ and the velocity of the condensed phase is proportional to ξ_μ . It is often convenient to define the component of ξ orthogonal to u ,

$$\zeta^\mu = P^{\mu\nu} \xi_\nu. \tag{3.2}$$

The equations of motion of the superfluid are given by

$$\begin{aligned} \partial_\mu T^{\mu\nu} &= F^{\nu\mu} J_\mu \\ \partial_\mu J^\mu &= c E_\mu B^\mu \\ \partial_\mu \xi_\nu - \partial_\nu \xi_\mu &= F_{\mu\nu} \end{aligned} \tag{3.3}$$

¹³In [34, 16] ξ was defined with an opposite sign.

together with the constitutive relations

$$\begin{aligned}
T^{\mu\nu} &= (\rho + P)u^\mu u^\nu + P\eta^{\mu\nu} + f\xi^\mu \xi^\nu + T_{diss}^{\mu\nu} \\
J^\mu &= qu^\mu - f\xi^\mu + J_{diss}^\mu \\
u \cdot \xi &= \mu + \mu_{diss}
\end{aligned} \tag{3.4}$$

where we have chosen to work in an arbitrary ‘fluid frame’ (see [10] for an explanation of this terminology and a fuller introduction to dissipative superfluid dynamics).

As was the case for the theory of charged fluids which we described in the previous section, superfluids also allow for a simple ‘canonical’ entropy current [10]

$$J_S^\mu{}_{\text{canon}} = su^\mu - \frac{\mu}{T}J_{diss}^\mu - \frac{u_\nu T_{diss}^{\mu\nu}}{T} \tag{3.5}$$

where s is the thermodynamical entropy density of our fluid and is related to ρ and P through the Gibbs-Duhem relation

$$\rho + P = sT + \mu q \tag{3.6}$$

and

$$dP = sdT + qd\mu + \frac{1}{2}fd\xi^2 \tag{3.7}$$

where

$$\xi = \sqrt{-\xi^\mu \xi_\mu}. \tag{3.8}$$

It has been demonstrated in [10] that the entropy current (3.5) is invariant under field redefinitions. It was also shown in [10] that the divergence of this entropy current is given by

$$\partial_\mu J_s^\mu = -\partial_\mu \left(\frac{u_\nu}{T} \right) T_{diss}^{\mu\nu} - \left(\partial_\mu \left(\frac{\mu}{T} \right) - \frac{E_\mu}{T} \right) J_{diss}^\mu + \frac{\mu_{diss}}{T} \partial_\mu (f\xi^\mu) \tag{3.9}$$

The rest of this section closely follows section 2. In 3.1 we list the independent first order data and second order scalar data, in section 3.2 we construct the most general positive divergence parity conserving entropy current consistent with Lorentz invariance. We find that up to a certain ambiguous term which is physically trivial, the entropy current agrees with its canonical form (3.5) and therefore the analysis of [10] follows.

3.1 Onshell inequivalent First order independent data

In the case of superfluid dynamics, the $SO(3, 1)$ tangent space symmetry at any point is generically broken down to $SO(2)$ by the nonzero velocity fields u^μ and ξ^μ . In the

special case that u^μ and ξ^μ are collinear, $SO(2)$ is enhanced to $SO(3)$. This special case is physically interesting since it implies that the superfluid component is motionless relative to the normal component—once the superfluid velocity is too large superfluidity breaks down. We will find it convenient to decompose all first order fluid dynamical data into representations of $SO(2)$ and treat the collinear limit as a special point in parameter space.

Representations of $SO(2)$ are all one dimensional. We refer to fluid dynamical data that is invariant under $SO(2)$ as scalar data. All other fluid data has charge $\pm m$ under $SO(2)$, where m is an integer. There is always as much $+m$ as $-m$ data. We will find it useful to group together $+1$ and -1 charge data into a two column which we refer to as vector data; similarly we group $+2$ and -2 data together into tensor data.

Now consider a vector \mathcal{V}^μ whose $m = 1$ and $m = -1$ components are (a, b) . The vector

$$\tilde{\mathcal{V}}^\mu = \epsilon^{\mu\nu\alpha\beta} u_\nu \xi_\alpha \mathcal{V}_\beta \equiv * \mathcal{V}_\mu \quad (3.10a)$$

is a pseudo vector. Its components are proportional to $(a, -b)$. Thus, when considering representations of $SO(2)$, the same data can be packaged into either a vector or into a pseudo vector. The story is similar for all nonscalar representations. For instance, a traceless symmetric tensor $\mathcal{T}_{\mu\nu}$, whose $m = 2$ and $m = -2$ components are (a, b) is simply related to a traceless symmetric pseudo tensor

$$\tilde{\mathcal{T}}_{\mu\nu} = \epsilon_{\mu\alpha\beta\gamma} u^\alpha \xi^\beta \mathcal{T}^{\gamma\nu} \equiv * \mathcal{T}_{\mu\nu} \quad (3.10b)$$

with components proportional to $(a, -b)$. Hence all tensor data can be packaged into pseudo tensors.

We now turn to a listing of the one derivative fluid dynamical and field data for superfluids. In Table 3 we explicitly list all one derivative data, one derivative equations of motion, and then eventually independent one derivative data. The scalar ξ used in this table is given by (3.8). We do not list pseudo vectors and pseudo tensors independently from vectors and tensors as they are isomorphic and contain the same data. In Table 4 we assign labels to our independent data. In the same table we also present a second listing of a basis for independent scalar data which will be more convenient at places. In Appendix A we demonstrate that both sets of seven scalars and the seven vectors listed are independent data, i.e. that we can solve for all other scalars and all other vectors in terms of the chosen basis.

As can be seen from Tables 3 and 4, after imposing the equations of motion we have six first order scalars and one first order pseudo scalar built out of fluid data, one first order scalar and one first order pseudo scalar built out of background field strengths, five first order vectors built out of fluid data, two first order vectors built from

Classification	All data	Equations of motion	Independent data
Scalars (set 1)	$\partial_\mu u^\mu$ $\partial_\mu(T\xi^\mu)$ $\xi^\mu \xi^\nu \partial_\mu u_\nu$ $\xi^\mu \partial_\mu \left(\frac{\xi}{T}\right)$ $\xi^\mu \partial_\mu \left(\frac{\mu}{T}\right)$ $\xi^\mu \partial_\mu T$ $E \cdot \xi$ $u^\mu \partial_\mu \left(\frac{\mu}{T}\right)$ $u^\mu \partial_\mu T$ $u^\mu \partial_\mu \left(\frac{\xi}{T}\right)$ $\xi^\mu u^\nu \partial_\nu u_\nu$	$\partial_\mu J^\mu = c E^\mu B_\mu$ $\xi^\mu u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) = \xi \cdot E$ $\xi_\mu \partial_\nu T^{\mu\nu} = \xi^\mu F_{\mu\nu} J^\nu$ $u_\mu \partial_\nu T^{\mu\nu} = -E_\mu J^\mu$	$\tilde{P}^{\mu\nu} \partial_\mu u_\nu$ $\tilde{P}^{\mu\nu} \partial_\mu (T\xi_\nu)$ $\xi^\mu \xi^\nu \partial_\mu u_\nu$ $\xi^\mu \partial_\mu \left(\frac{\mu}{T}\right)$ $\xi^\mu \partial_\mu \left(\frac{\xi}{T}\right)$ $\xi^\mu \partial_\mu T$ $E \cdot \xi$
Scalars (set 2)	$\partial_\mu u^\mu$ $\partial_\mu(T\xi^\mu)$ $\xi^\mu \xi^\nu \partial_\mu u_\nu$ $\xi^\mu \partial_\mu \left(\frac{\xi}{T}\right)$ $\xi^\mu \partial_\mu \left(\frac{\mu}{T}\right)$ $\xi^\mu \partial_\mu T$ $E \cdot \xi$ $u^\mu \partial_\mu \left(\frac{\mu}{T}\right)$ $u^\mu \partial_\mu T$ $u^\mu \partial_\mu \left(\frac{\xi}{T}\right)$ $\xi^\mu u^\nu \partial_\nu u_\nu$	$\partial_\mu J^\mu = c E^\mu B_\mu$ $\xi^\mu u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) = \xi \cdot E$ $\xi_\mu \partial_\nu T^{\mu\nu} = \xi^\mu F_{\mu\nu} J^\nu$ $u_\mu \partial_\nu T^{\mu\nu} = -E_\mu J^\mu$	$\tilde{P}^{\mu\nu} \partial_\mu u_\nu$ $\tilde{P}^{\mu\nu} \partial_\mu (T\xi_\nu)$ $u^\mu \xi^\nu \partial_\mu u_\nu$ $u^\mu \partial_\mu \left(\frac{\mu}{T}\right)$ $u^\mu \partial_\mu \left(\frac{\xi}{T}\right)$ $u^\mu \partial_\mu T$ $E \cdot \xi$
Pseudo scalars	$\omega \cdot \xi$ $B \cdot \xi$ $\epsilon^{\mu\nu\alpha\beta} u_\mu \xi_\nu \partial_\alpha \xi_\beta$	$e^{\alpha\beta} (\partial_\alpha \xi_\beta - \partial_\beta \xi_\alpha) = e^{\alpha\beta} F_{\alpha\beta}$	$\omega \cdot \xi$ $B \cdot \xi$
Vectors	$\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho u_\nu$ $\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \xi_\nu$ $\tilde{P}^{\mu\nu} E_\nu$ $\tilde{P}^{\mu\nu} F_{\nu\beta} \xi^\beta$ $\tilde{P}^{\mu\nu} \partial_\nu \frac{\mu}{T}$ $\tilde{P}^{\mu\nu} \partial_\nu \frac{\xi}{T}$ $\tilde{P}^{\mu\nu} \partial_\nu T$ $\tilde{P}^{\mu\nu} \xi^\alpha \partial_\nu u_\alpha$ $\tilde{P}^{\mu\nu} u^\rho \partial_\rho u_\nu$ $\tilde{P}^{\mu\nu} u^\rho \partial_\rho \xi_\nu$	$\tilde{P}^{\mu\nu} \partial_\beta T^\beta_\nu = \tilde{P}^{\mu\nu} F_{\nu\beta} J^\beta$ $\tilde{P}^{\alpha\mu} u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) = \tilde{P}^{\alpha\mu} E_\mu$ $\tilde{P}^{\alpha\mu} \xi^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) = \tilde{P}^{\alpha\mu} F_{\mu\nu} \xi^\nu$	$\tilde{P}^{\mu\nu} u^\rho \partial_\rho u_\nu$ $\tilde{P}^{\mu\nu} u^\rho \partial_\rho \xi_\nu$ $\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho u_\nu$ $\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \xi_\nu$ $\tilde{P}^{\mu\nu} E_\nu$ $\tilde{P}^{\mu\nu} F_{\nu\beta} \xi^\beta$ $\tilde{P}^{\mu\nu} \partial_\nu \frac{\mu}{T}$
Tensors	$\sigma_{\mu\nu}^u$ $\sigma_{\mu\nu}^\xi$	<p style="text-align: center;">–</p> <p style="text-align: center;">–</p>	$\sigma_{\mu\nu}^u$ $\sigma_{\mu\nu}^\xi$

Table 3: One derivative data for superfluids. The first column lists all quantities formed from the action of a single derivative on fluid and background fields. The second column lists all one derivative equations of motion. The last columns lists a choice of independent data. The tensors $\sigma_{\mu\nu}^\xi$ and $\sigma_{\mu\nu}^u$ are defined in (3.12)-(3.11). We also used $e^{\alpha\beta} = \epsilon^{\mu\nu\alpha\beta} u_\mu \xi_\nu$.

background fields and two independent tensors. The first tensor is simply the usual

i	1	2	3	4	5	6	7
\mathcal{S}_i^a	$\tilde{P}^{\mu\nu}\partial_\mu u_\nu$	$\tilde{P}^{\mu\nu}\partial_\mu(T\xi_\nu)$	$\xi^\mu\xi^\nu\partial_\mu u_\nu$	$\xi^\mu\partial_\mu\left(\frac{\xi}{T}\right)$	$\xi^\mu\partial_\mu\left(\frac{\mu}{T}\right)$	$\xi^\mu\partial_\mu T$	$E.\xi$
\mathcal{S}_i^b	$\tilde{P}^{\mu\nu}\partial_\mu u_\nu$	$\tilde{P}^{\mu\nu}\partial_\mu(T\xi_\nu)$	$u^\mu\xi^\nu\partial_\mu u_\nu$	$u^\mu\partial_\mu\left(\frac{\xi}{T}\right)$	$u^\mu\partial_\mu\left(\frac{\mu}{T}\right)$	$u^\mu\partial_\mu T$	$E.\xi$
\mathcal{V}_i^a	$\tilde{P}^{\mu\nu}u^\rho\partial_\rho u_\nu$	$\tilde{P}^{\mu\nu}u^\rho\partial_\rho\xi_\nu$	$\tilde{P}^{\mu\nu}\xi^\rho\partial_\rho u_\nu$	$\tilde{P}^{\mu\nu}\xi^\rho\partial_\rho\xi_\nu$	$\tilde{P}^{\mu\nu}\partial_\nu\frac{\mu}{T}$	$\tilde{P}^{\mu\nu}\partial_\nu E_\mu$	$\tilde{P}^{\mu\nu}F_{\nu\beta}\xi^\beta$

Table 4: Labels for the two sets of independent one derivative scalars and one set of independent vectors.

shear tensor $\sigma^{\mu\nu}$ projected orthogonal to the plane formed by the two fluid velocities.

$$\sigma_{\mu\nu}^u = \tilde{P}^{\mu\alpha}\tilde{P}^{\nu\beta}\left(\sigma_{\alpha\beta} - \frac{1}{2}\eta_{\alpha\beta}\tilde{P}_{\gamma\delta}\sigma^{\gamma\delta}\right). \quad (3.11)$$

The second tensor $\sigma_{\mu\nu}^\xi$ is defined by

$$\sigma_{\mu\nu}^\xi = \frac{1}{2}\tilde{P}^{\mu\alpha}\tilde{P}^{\nu\beta}\left(\partial_\alpha\xi_\beta + \partial_\beta\xi_\alpha - \tilde{P}_{\alpha\beta}\tilde{P}^{\gamma\delta}\partial_\gamma\xi_\delta\right). \quad (3.12)$$

The counting of data in the absence of background fields agrees with [10].

As was the case for normal fluids, the divergence of the first order superfluid entropy current is a sum over quadratic one derivative terms and two derivative pieces of data. In order to assist the analysis of the positivity of the divergence of the entropy current we list all the scalar two derivative data, the two derivative equations of motion and a basis for onshell independent two derivative scalars in Table 5. Note that we have nine independent pieces of two derivative fluid dynamical data (as reported in [10]) together with four additional pieces of two derivative data from background field strengths. In Appendix A demonstrate that the scalars listed in the last column of table 5 form a basis of onshell independent scalars.

3.2 Constructing the entropy current

With the independent data at hand we proceed with our analysis. The most general entropy current allowed by symmetries takes the form

$$J_S^\mu = J_{S\text{ canon}}^\mu + u^\mu \sum_{i=1}^7 s_i^a \mathcal{S}_i^a + \xi^\mu \sum_{i=1}^7 s_i^b \mathcal{S}_i^b + \sum_{i=1}^7 v_i \mathcal{V}_i^{a\mu}, \quad (3.13)$$

where the coefficients s_i^a , s_i^b , and v_i are, at the moment, arbitrary functions of $\frac{\mu}{T}$, T , and ξ^2 . Note that we have chosen to expand the terms proportional to u^μ in the basis \mathcal{S}_i^a while terms proportional to ξ^μ are expanded in the basis \mathcal{S}_i^b . This choice will prove

All data	$\tilde{P}^{\mu\nu} u^\rho \partial_\rho \partial_\mu u_\nu$	$\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \partial_\mu u_\mu$	$\tilde{P}^{\mu\nu} u^\rho \partial_\rho \partial_\mu (T \xi_\nu)$	$\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \partial_\mu (T \xi_\nu)$
	$\xi^\mu \xi^\nu u^\rho \partial_\rho \partial_\mu u_\nu$	$u^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\xi}{T}\right)$	$u^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\mu}{T}\right)$	$u^\mu \xi^\rho \partial_\rho \partial_\mu T$
	$\tilde{P}^{\mu\nu} \partial_\mu \partial_\nu \left(\frac{\mu}{T}\right)$	$\tilde{P}^{\mu\nu} \partial_\mu \partial_\nu \left(\frac{\xi}{T}\right)$	$\tilde{P}^{\mu\nu} \partial_\mu \partial_\nu T$	$u^\mu u^\rho \partial_\rho \partial_\mu \left(\frac{\xi}{T}\right)$
	$\xi^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\xi}{T}\right)$	$\xi^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\mu}{T}\right)$	$\xi^\mu \xi^\rho \partial_\rho \partial_\mu T$	$u^\mu u^\rho \partial_\rho \partial_\mu \left(\frac{\mu}{T}\right)$
	$u^\mu u^\rho \partial_\rho \partial_\mu T$	$\xi^\mu \xi^\nu \xi^\rho \partial_\rho \partial_\mu u_\nu$	$u^\mu \xi^\nu u^\rho \partial_\rho \partial_\mu u_\nu$	$\xi^\nu \tilde{P}^{\rho\mu} \partial_\rho \partial_\mu u_\nu$
	$u^\mu \xi^\nu \partial_\mu E_\nu$	$\xi^\mu \xi^\nu \partial_\mu E_\nu$	$\tilde{P}^{\mu\nu} \partial_\mu E_\nu$	$\epsilon^{\mu\nu\lambda\sigma} \xi_\mu u_\lambda \partial_\mu B_\sigma$
Equations of motion	$u^\beta \partial_\beta (\partial_\mu J^\mu) = u^\beta \partial_\beta (c E^\mu B_\mu)$		$\xi^\beta \partial_\beta (\partial_\mu J^\mu) = \xi^\beta \partial_\beta (c E^\mu B_\mu)$	
	$u^\beta \partial_\beta (\xi_\mu \partial_\nu T^{\mu\nu}) = u^\beta \partial_\beta (\xi^\mu F_{\mu\nu} J^\nu)$		$\xi^\beta \partial_\beta (\xi_\mu \partial_\nu T^{\mu\nu}) = \xi^\beta \partial_\beta (\xi^\mu F_{\mu\nu} J^\nu)$	
	$u^\beta \partial_\beta (u_\mu \partial_\nu T^{\mu\nu}) = u^\beta \partial_\beta (-E_\mu J^\mu)$		$\xi^\beta \partial_\beta (u_\mu \partial_\nu T^{\mu\nu}) = \xi^\beta \partial_\beta (-E_\mu J^\mu)$	
	$u^\beta \partial_\beta (\xi^\mu u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu)) = u^\beta \partial_\beta (\xi^\mu E_\mu)$		$\xi^\beta \partial_\beta (\xi^\mu u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu)) = \xi^\beta \partial_\beta (\xi^\mu E_\mu)$	
	$\partial_\alpha \left(\tilde{P}^{\alpha\mu} u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) \right) = \partial_\alpha \left(\tilde{P}^{\alpha\mu} E_\mu \right)$		$\partial_\mu \left(\tilde{P}^{\mu\nu} \partial_\beta T^\beta_\nu \right) = \partial_\mu \left(\tilde{P}^{\mu\nu} F_{\nu\beta} J^\beta \right)$	
	$\partial_\alpha \left(\tilde{P}^{\alpha\mu} \xi^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) \right) = \partial_\alpha \left(\tilde{P}^{\alpha\mu} F_{\mu\nu} \xi^\nu \right)$			
Independent data	$\tilde{P}^{\mu\nu} u^\rho \partial_\rho \partial_\mu u_\nu$	$\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \partial_\mu u_\mu$	$\tilde{P}^{\mu\nu} u^\rho \partial_\rho \partial_\mu (T \xi_\nu)$	$\tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \partial_\mu (T \xi_\nu)$
	$\xi^\mu \xi^\nu u^\rho \partial_\rho \partial_\mu u_\nu$	$u^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\xi}{T}\right)$	$u^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\mu}{T}\right)$	$u^\mu \xi^\rho \partial_\rho \partial_\mu T$
	$\tilde{P}^{\mu\nu} \partial_\mu \partial_\nu \left(\frac{\mu}{T}\right)$	$u^\mu \xi^\nu \partial_\mu E_\nu$	$\xi^\mu \xi^\nu \partial_\mu E_\nu$	$\tilde{P}^{\mu\nu} \partial_\mu E_\nu$
	$\tilde{P}^{\mu\nu} (\partial_\mu F_{\nu\beta}) \xi^\beta$			

Table 5: Two derivative scalar data. The first row gives all two derivative scalar data, the second row lists all the equations of motion. The third row represents a particular choice of independent second order data.

algebraically convenient below. In total we start with twenty one free parameters in the entropy current.

The two derivative terms in the divergence of the entropy current (3.13) are given by

$$\begin{aligned}
\partial_\mu J_S^\mu &= (s_1^a + v_1) \tilde{P}^{\mu\nu} u^\rho \partial_\rho \partial_\mu u_\nu + (s_2^a + v_2) \tilde{P}^{\mu\nu} u^\rho \partial_\rho \partial_\mu (T \xi_\nu) + (s_1^b + v_3) \tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \partial_\mu u_\nu \\
&\quad + (s_2^b + v_4) \tilde{P}^{\mu\nu} \xi^\rho \partial_\rho \partial_\mu (T \xi_\nu) + (s_3^a + s_3^b) \xi^\mu \xi^\nu u^\rho \partial_\rho \partial_\mu u_\nu \\
&\quad + (s_3^a + s_3^b) u^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\xi}{T}\right) + (s_3^a + s_3^b) u^\mu \xi^\rho \partial_\rho \partial_\mu \left(\frac{\mu}{T}\right) \\
&\quad + (s_3^a + s_3^b) u^\mu \xi^\rho \partial_\rho \partial_\mu T + v_5 \tilde{P}^{\mu\nu} \partial_\mu \partial_\nu \left(\frac{\mu}{T}\right) + s_7^a u^\mu \xi^\nu \partial_\mu E_\nu \\
&\quad + s_7^b \xi^\mu \xi^\nu \partial_\mu E_\nu + v_6 \tilde{P}^{\mu\nu} \partial_\mu E_\nu + v_7 \tilde{P}^{\mu\nu} (\partial_\mu F_{\nu\beta}) \xi^\beta + \dots
\end{aligned} \tag{3.14}$$

Following the algorithm of the previous section, we first set the coefficient of each of the thirteen independent two derivative terms listed in Table 5, which appear in the

divergence of the entropy current, to zero. The vanishing of the nine fluid dynamical two derivative terms yields the following nine relations between the s^a 's s^b 's and v 's

$$\begin{aligned} v_1 &= -s_1^a & v_2 &= -s_2^a & v_3 &= -s_1^b & v_4 &= -s_2^b & s_3^b &= -s_3^a & (3.15) \\ s_4^b &= -s_4^a & s_5^b &= -s_5^a & s_6^b &= -s_6^a & v_5 &= 0. \end{aligned}$$

The vanishing of the four electromagnetic field related two derivative scalars yields the additional four relations

$$s_7^a = s_7^b = v_6 = v_7 = 0. \quad (3.16)$$

Apart from the two derivative fluid dynamical and background electromagnetic field data, there are four nontrivial curvature invariants one can form out of the contractions of u^μ , ξ^μ and $g^{\mu\nu}$ with the the Reimann tensor $R_{\alpha\beta\mu\nu}$.¹⁴ After plugging in the constraints in (3.15) and (3.16) into the expression for the entropy current (3.13) we find

$$\begin{aligned} \partial_\mu J_S^\mu &= s_1^a \tilde{P}^{\alpha\beta} u^\mu u^\lambda R_{\lambda\beta\alpha\mu} + s_2^b \tilde{P}^{\alpha\beta} \xi^\mu \xi^\lambda R_{\lambda\beta\alpha\mu} \\ &+ (T s_2^a + s_1^b) \xi^\mu u^\lambda R_{\lambda\beta\alpha\mu} + s_3^a u^\alpha \xi^\beta u^\gamma \xi^\delta R_{\alpha\beta\gamma\delta} + \dots \end{aligned} \quad (3.17)$$

Each of the terms in (3.17) is of indefinite sign. Thus, the coefficients of these four terms must vanish. This implies

$$s_1^a = 0, \quad s_1^b = -T s_2^a, \quad s_2^b = 0, \quad s_3 = 0. \quad (3.18)$$

To summarize, by setting the two derivative and curvature terms that appear in the divergence of the entropy current to zero we have eliminated $9 + 4 + 4 = 17$ of the original 21 coefficients and are left with an entropy current with four undetermined coefficients,

$$\begin{aligned} J_S^\mu &= J_{S \text{ canon}}^\mu + s_2^a \left(u^\mu \tilde{P}^{\alpha\beta} \partial_\alpha (T \xi_\beta) - \tilde{P}^{\mu\beta} u^\nu \partial_\nu (T \xi_\beta) - T \xi^\mu \tilde{P}^{\alpha\beta} \partial_\alpha u_\beta + T \tilde{P}^{\mu\beta} \xi^\nu \partial_\nu u_\beta \right) \\ &+ s_4^a \left(u^\mu \xi^\nu \partial_\nu \frac{\xi}{T} - \xi^\mu u^\nu \partial_\nu \frac{\xi}{T} \right) + s_5^a \left(u^\mu \xi^\nu \partial_\nu \frac{\mu}{T} - \xi^\mu u^\nu \partial_\nu \frac{\mu}{T} \right) + s_6^a (u^\mu \xi^\nu \partial_\nu T - \xi^\mu u^\nu \partial_\nu T). \end{aligned} \quad (3.19)$$

The entropy current (3.19) can be rewritten in a simpler form by introducing the antisymmetric tensor

$$\mathcal{Q}_{\mu\nu} = T(\xi_\mu u_\nu - \xi_\nu u_\mu), \quad (3.20)$$

introducing a unified notation for the three thermodynamical scalar fields

$$\Sigma_i = \left\{ \frac{\mu}{T}, \frac{\xi}{T}, T \right\} \quad i = 1, 2, 3,$$

¹⁴We omit the curvature scalar R in this listing since it is a pure gravitational term and therefore never appears in the divergence of fluid dynamical entropy current.

and also redefining our coefficient functions

$$c_0 = s_2^a \quad c_1 = s_5 - \frac{T\mu}{\mu^2 - \xi^2} \quad c_2 = s_4 + \frac{T\xi}{\mu^2 - \xi^2} \quad c_3 = s_6 - \frac{2}{T}.$$

Then (3.19) takes the form

$$J_S^\mu = J_{S \text{ canon}}^\mu + c_0 \partial_\nu Q^{\nu\mu} + \sum_{i=1}^3 c_i Q^{\mu\nu} \partial_\nu \Sigma_i. \quad (3.21)$$

The divergence of the entropy current (3.21) is given by

$$\begin{aligned} \partial_\mu J_S^\mu = & -\partial_\mu \left(\frac{u_\nu}{T} \right) T_{diss}^{\mu\nu} + V_{1\mu} J_{diss}^\mu + \frac{\mu_{diss}}{T} \partial_\mu (f \xi_\nu) \\ & + (\partial_{\Sigma_i} c_0) (\partial_\mu \Sigma_i) \partial_\nu Q^{\nu\mu} + c_i \partial_\mu Q^{\mu\nu} \partial_\nu \Sigma_i + (\partial_{\Sigma_j} c_i) (\partial_\mu \Sigma_j) Q^{\mu\nu} (\partial_\nu \Sigma_i). \end{aligned} \quad (3.22)$$

The first line of the right hand side of (3.22) corresponds to the divergence of the canonical entropy current and the second line corresponds to the divergence of the new terms. The right hand side of (3.22) can be written as a sum of three classes of quadratic forms: a quadratic form in one derivative scalar data, a quadratic form in one derivative vector data and a quadratic form in one derivative tensor data. Let us first focus our attention on the quadratic form in the vector data. All vector terms that appear on the right hand side of (3.22) are linear combination of the six independent vector pieces of data,

$$\tilde{P}^{\alpha\nu} \xi^\mu \sigma_{\mu\nu}, \quad \tilde{P}_\mu^\alpha \partial_\nu Q^{\mu\nu}, \quad \tilde{P}_\mu^\alpha V_1^\mu, \quad \tilde{P}^{\alpha\nu} (\partial_\nu \Sigma_i) \quad (i = 1, \dots, 3).$$

Independent of the structure of the (as yet undetermined) dissipative terms $T_{diss}^{\mu\nu}$ and J_{diss}^μ , the quadratic form in one derivative vectors does not contain squares of $\partial_\mu \Sigma_i$, and $\tilde{P}_\mu^\alpha \partial_\nu Q^{\mu\nu}$.¹⁵ It is an easily verified fact that a quadratic form that does not contain the square of any given variable is positive if and only if it is independent of that variable. Thus, positivity of the divergence of the entropy current requires that the right hand side of (3.22) does not contain any term proportional to $P_\mu^\alpha \partial_\alpha \Sigma_i$ and $\tilde{P}_\mu^\alpha \partial_\nu Q^{\mu\nu}$.

The term in (3.22) proportional to $\tilde{P}_\mu^\alpha \partial_\alpha \Sigma_i \times \partial_\nu Q^{\mu\nu}$ appears with a coefficient $(c_i + \partial_{\Sigma_i} c_0)$. Setting this coefficient to zero yields an expression for the c_i 's in terms of c_0

$$c_i = -\partial_{\Sigma_i} c_0. \quad (3.23)$$

Inserting (3.23) into (3.21) we find

$$J_S^\mu = J_{S \text{ canon}}^\mu + \partial_\nu (c_0 Q^{\nu\mu}). \quad (3.24)$$

¹⁵This follows because these terms do not appear explicitly on the right hand side of the divergence of the canonical entropy current.

We have ended up with a one parameter set of entropy currents (3.24) specified by the undetermined coefficient c_0 . The divergence of these entropy currents is, however, independent of c_0 . In fact, due to the antisymmetry properties of $\mathcal{Q}^{\mu\nu}$

$$\partial_\mu J_S^\mu = \partial_\mu J_{S_{\text{canon}}}^\mu. \quad (3.25)$$

The parameter c_0 is essentially trivial and is related to a pullback ambiguity as we now explain. It was pointed out in [26] that the following set of operations maps one positive divergence entropy current J_S^μ to another

1. Dualize J^μ to a three-form.
2. Shift this three-form by its Lie derivative with respect to any vector field V^μ
3. Dualize the resultant form back to a current.

The end result of this operation is a shift in the entropy current given by (see eq. 6.6 in [26])

$$\delta J_S^\mu = \nabla_\nu (J_S^\nu V^\mu - V^\nu J_S^\mu) + V^\mu \nabla_\nu J_S^\nu. \quad (3.26)$$

In the current setup we are interested in first order corrections to the entropy current. The right hand side of (3.26) has an explicit derivative. Therefore, the entropy current on the right hand side should be replaced by the perfect fluid entropy current $J_S^\mu = su^\mu$. This implies that the second term on the right hand side of (3.26) is zero (recall that the perfect fluid entropy current is divergence free). Moreover V^μ must be a derivative free vector field.

In ordinary (non superfluid) fluid dynamics there is a unique vector at the zero derivative order—the fluid velocity u^μ . Since $J^\mu \propto u^\mu$ then $V^\mu \propto u^\mu$ implies that the first term on the right hand side of (3.26) also vanishes, and so (3.26) leads to no ambiguity in the entropy current at the first derivative order.

In superfluid dynamics there exist two zero derivative vectors, u^μ and ξ^μ . Consequently (3.26) can be used to generate a shift in the current proportional to $\partial_\nu (c_0 \mathcal{Q}^{\nu\mu})$. We conclude that the freedom to add the total derivative term $\partial_\nu (c_0 \mathcal{Q}^{\nu\mu})$ is precisely the ‘pullback ambiguity’ freedom described in [26].

Going back to (3.25), it follows that the constraints on the dissipative terms $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} from demanding the positivity of the divergence of J_S are identical to the constraints from the positivity of the divergence of the canonical entropy current $J_{S_{\text{canon}}}^\mu$. Such an analysis has been carried out in [10] and we will be carried out in more generality in the next section. We will not repeat it here.

4. Superfluid dynamics without parity invariance

The main goal of this work is to determine the most general expression for the constitutive relations for a Lorentz invariant but parity breaking superfluid. We tackle this problem using the same algorithm which has been spelt out in the previous sections: we determine the most general entropy current which is compatible with the symmetries. The restrictions on the constitutive relations are then determined by demanding positivity of the entropy current.

4.1 Onshell inequivalent data

As we've discussed in section 3.1, the listing of one derivative pseudo vectors and pseudo-tensors of $SO(2)$ can be constructed from the tensors of $SO(2)$ using the ϵ tensor. If \mathcal{V}_μ and $\mathcal{T}_{\mu\nu}$ are vectors and tensors of $SO(2)$ then we define the associated pseudo vectors $\tilde{\mathcal{V}}^\mu$ and pseudo-tensors $\tilde{\mathcal{T}}^{\mu\nu}$ through:

$$\begin{aligned}\tilde{\mathcal{V}}^\mu &= *\mathcal{V}_\mu = \epsilon^{\mu\nu\alpha\beta} u_\nu \xi_\alpha \mathcal{V}_\beta \\ \tilde{\mathcal{T}}^{\mu\nu} &= *\mathcal{T}_{\mu\nu} = \epsilon^{\mu\rho\alpha\beta} u_\rho \xi_\alpha \mathcal{T}_\beta{}^\nu + \epsilon^{\nu\rho\alpha\beta} u_\rho \xi_\alpha \mathcal{T}_\beta{}^\mu.\end{aligned}\tag{4.1}$$

We introduce the following terminology: $*\mathcal{V}_\mu$ is said to be the star of \mathcal{V}_μ , and similarly $*\mathcal{T}_{\mu\nu}$ is said to be the star of $\mathcal{T}_{\mu\nu}$.

Note that $*V_\mu$ is an $SO(2)$ pseudovector for any vector V_μ since the $*$ operation projects onto the subspace orthogonal to both u^μ and ζ^μ where ζ^μ was defined in (3.2),

$$\zeta^\mu = P^{\mu\nu} \xi_\nu.$$

In Table 6 we list the independent one derivative pseudo-scalars, vectors and tensors which we will use in this section. It proves convenient to choose our on-shell independent scalars S_i^c to differ from both the choices made in section 3. The linear independence of this choice of scalars is demonstrated in Appendix C. Our choice of on-shell independent vectors is also different from the one is made in Table 4 of section 3. We have used the superscript c for the pseudo vectors and vectors to flag this difference. We have demonstrated the linear independence of this choice of vectors in Appendix C.

In table 7 we list all two derivative pseudo scalars. The format we use is identical to that of previous sections. We list all possible two derivative pseudo scalars that can be constructed out of fluid and field data in the left column. The equations of motion appear in the second column. For technical reasons we find it convenient to formally treat E_μ and B_μ as independent fields in the full list of two derivative data (the left column in table 7) and to impose the Bianchi identity (which relates some of the derivatives of these fields to others) at the same stage that we impose the equations

i	1	2	3	4	5	6	7
$\tilde{\mathcal{V}}_i^{\mu}$	$*V_{1\mu}$	$*(\zeta^\alpha \sigma_{\alpha\beta})$	$*\nabla_\sigma T$	$*\nabla_\sigma \left(\frac{\mu}{T}\right)$	$*\nabla_\sigma \left(\frac{\zeta^2}{T^2}\right)$	$\tilde{P}^{\mu\nu} \omega_\nu$	$\tilde{P}_\sigma^\mu B^\sigma$
\mathcal{V}_i^{μ}	$\tilde{P}^{\mu\sigma} V_{1\sigma}$	$\tilde{P}^{\mu\beta} (\zeta^\alpha \sigma_{\alpha\beta})$	$\tilde{P}^{\mu\sigma} \nabla_\sigma T$	$\tilde{P}^{\mu\sigma} \nabla_\sigma \left(\frac{\mu}{T}\right)$	$\tilde{P}^{\mu\sigma} \nabla_\sigma \left(\frac{\zeta^2}{T^2}\right)$	$\frac{\mathcal{V}_2^{\mu} - \tilde{P}^{\mu\alpha} \zeta^\nu \partial_\alpha u_\nu}{\zeta^2}$	$-\frac{P^{\mu\nu} E_{\nu\alpha} \zeta^\alpha}{\zeta^2}$
S^c	$\zeta_\mu V_1^\mu$	$u \cdot \partial T$	$u \cdot \partial \frac{\mu}{T}$	$u \cdot \partial \frac{\zeta^2}{T^2}$	$\zeta^\mu P^{\mu\alpha} \partial_\alpha T$	$\zeta^\mu E_\mu$	$\zeta^\mu P^{\mu\alpha} \partial_\alpha \frac{\zeta^2}{T^2}$
$\tilde{\mathcal{T}}_{i\mu\nu}$	$*\sigma_{\mu\nu}^u$	$*\sigma_{\mu\nu}^\xi$	-	-	-	-	-
$\tilde{\mathcal{S}}_i$	$\omega \cdot \xi$	$B \cdot \xi$	-	-	-	-	-

Table 6: Parity odd independent one derivative data which we use in this section. The $*$ operation that takes us from vectors \mathcal{V} and tensors \mathcal{T} to pseudovectors $\tilde{\mathcal{V}}$ and pseudotensors, $\tilde{\mathcal{T}}$ is given in (4.1). We list a basis for pseudovectors, and a new basis for scalars and vectors which we will work with in this section. Note that $\tilde{\mathcal{V}}_i^c = *\mathcal{V}_i^c$.

of motion. In other words, in our listings below we will pretend that E_μ and B_μ are independent fields constrained by “equations of motion” which are given by the Bianchi identities. In the last column we list a basis of independent two derivative pseudo scalars. In Appendix C we have demonstrated that the two derivative pseudo scalars presented in Table 7 are independent.

All data	equations of motion	Two derivative data
$\zeta^\mu u^\nu \nabla_\nu B_\mu$		
$\zeta^\mu \zeta^\nu \nabla_\nu B_\mu$	$e^{\lambda\sigma} (\zeta \cdot \nabla) C_{\lambda\sigma} = 0$	
$\tilde{P}^{\mu\nu} \nabla_\nu B_\mu$	$e^{\lambda\sigma} (\nabla_\theta T_\sigma^\theta - F_{\sigma\theta} j^\theta) = 0$	$e^{\lambda\sigma} \nabla_\lambda E_\sigma$
$e^{\lambda\sigma} \nabla_\lambda E_\sigma$	$e^{\lambda\sigma} u^\theta \nabla_\lambda C_{\theta\sigma} = 0$	$e^{\lambda\sigma} (\zeta \cdot \nabla) \nabla_\lambda u_\sigma$
$e^{\lambda\sigma} (u \cdot \nabla) \nabla_\lambda u_\sigma$	$\epsilon^{\mu\nu\lambda\sigma} \zeta_\mu \nabla_\nu F_{\lambda\sigma} = 0$	$\zeta^\mu \zeta^\nu \nabla_\mu B_\nu$
$e^{\lambda\sigma} (\zeta \cdot \nabla) \nabla_\lambda u_\sigma$	$\epsilon^{\mu\nu\lambda\sigma} u_\mu \nabla_\nu F_{\lambda\sigma} = 0$	
$e^{\lambda\sigma} (u \cdot \nabla) \nabla_\lambda \zeta_\sigma$		
$e^{\lambda\sigma} (\zeta \cdot \nabla) \nabla_\lambda \zeta_\sigma$		

Table 7: Parity odd two derivative scalar data. The first column lists all possible independent two derivative scalars with B^μ and E^μ treated as independent, the second column lists the equations of motion (first three rows) and Bianchi identities (last two rows) and the last column the independent data. We have used the shorthand notation $e^{\lambda\sigma} = \epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu$ and $C_{\mu\nu} \equiv \nabla_\mu \xi_\nu - \nabla_\nu \xi_\mu - F_{\mu\nu}$.

4.2 Constraints from vanishing of two derivative and curvature terms

We denote the total entropy current by

$$J_S^\mu = J_{S_{\text{canon}}}^\mu + J_{S_{\text{new}}}^\mu + \partial_\nu (c_0 Q^{\mu\nu}), \quad (4.2)$$

where $J_{S_{\text{new}}}^\mu$ is a parity odd contribution to the entropy current and $\partial_\nu (c_0 Q^{\mu\nu})$ is a residual ambiguity in the definition of the entropy current in the parity preserving sector described in section 3. The most general contribution to $J_{S_{\text{new}}}^\mu$ is given by

$$J_{S_{\text{new}}}^\mu = \sum_{i=1}^7 \tilde{v}_i \tilde{\mathcal{V}}_i^{c\mu} + u^\mu \sum_{i=1}^2 \tilde{s}_i^a \tilde{\mathcal{S}}_i + \zeta^\mu \sum_{i=1}^2 \tilde{s}_i^b \tilde{\mathcal{S}}_i. \quad (4.3)$$

This entropy current has eleven new coefficients, each of which is an arbitrary function of T , μ and ξ . In what follows we will swap the variables μ , and ξ with

$$\nu = \frac{\mu}{T} \quad \text{and} \quad \chi = \frac{\zeta^2}{T^2}. \quad (4.4)$$

As in previous sections, the eleven parameter set of currents is significantly constrained by the requirement that two derivative and curvature terms in its divergence vanish. It will prove convenient to rewrite the entropy current in the form

$$\begin{aligned} J_{S_{\text{new}}}^\mu = & \epsilon^{\mu\nu\rho\sigma} \partial_\nu (\sigma_1 T u_\rho \zeta_\sigma) + \sigma_3 \tilde{\mathcal{V}}_3^{c\mu} + T \sigma_4 \tilde{\mathcal{V}}_4^{c\mu} + T \sigma_5 \tilde{\mathcal{V}}_5^{c\mu} \\ & + \frac{\sigma_8}{2} \epsilon^{\mu\nu\rho\sigma} \zeta_\nu F_{\rho\sigma} + T^2 \sigma_9 \omega^\mu + T \sigma_{10} B^\mu \\ & + \alpha_1 \tilde{\mathcal{V}}_1^{c\mu} + \alpha_2 \tilde{\mathcal{V}}_2^{c\mu} + \zeta^\mu [\alpha_3 (\omega \cdot \zeta) + \alpha_4 (B \cdot \zeta)] \end{aligned} \quad (4.5)$$

The coefficient functions σ_i in (4.9) are linearly related to the coefficient functions in (4.3). Although we will never need these relations, for the sake of completeness we present them in Appendix C.

The first term in (4.9) is the divergent-less combination

$$\frac{1}{2} \epsilon^{\mu\nu\lambda\sigma} \nabla_\nu (\sigma_1 T u_\lambda \zeta_\sigma) \quad (4.6)$$

with σ_1 an undetermined function of T, ν and χ . This expression contributes to the entropy current but does not contribute to its divergence. We've observed a similar expression in the parity even sector of the entropy current in equation (3.24). As was the case in the previous section, this term is physically irrelevant (it does not contribute in any way to the constraints on dissipative terms) and can be ignored in what follows.

We now proceed to compute the correction to the entropy current (4.5). It is not difficult to verify that the two derivative and curvature contribution to the divergence of the entropy current is given by

$$\begin{aligned} \nabla_\mu J_{\text{new}}^\mu = & \alpha_1 [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] \nabla_\lambda E_\sigma + \frac{(\alpha_2 + \zeta \alpha_3)}{2} [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] (\zeta \cdot \nabla) \nabla_\lambda u_\sigma \\ & + \alpha_4 \zeta^\mu \zeta^\nu \nabla_\mu B_\nu + \frac{\alpha_2}{2} \epsilon^{\mu\nu\lambda\sigma} \zeta_\mu u_\nu u^\rho \zeta^\theta \left[R_{\rho\sigma\lambda\theta} - \frac{R_{\rho\theta\sigma\lambda}}{2} \right] + \dots \end{aligned} \quad (4.7)$$

where the dots denote the derivative squared terms. The expression on the right hand side of (4.7) is a sum over a set of linearly independent two derivative data and curvature terms (see table 7). Positivity of the divergence of the entropy current requires that the coefficients of the two derivative data and curvature terms vanish. Thus,

$$\alpha_1 = \alpha_2 = \alpha_3 = \alpha_4 = 0. \quad (4.8)$$

Had we not demanded the vanishing of curvature terms above, we would have erroneously concluded that α_2 is non zero. Inserting (4.8) into (4.5) we conclude that the most general parity odd addition to the entropy current whose divergence has no two derivative data or curvature terms is given by

$$J_{S_{\text{new}}}^\mu = \epsilon^{\mu\nu\rho\sigma} \partial_\nu (\sigma_1 T u_\rho \zeta_\sigma) + \sigma_3 \tilde{\mathcal{V}}_3^{c\mu} + T \sigma_4 \tilde{\mathcal{V}}_4^{c\mu} + T \sigma_5 \tilde{\mathcal{V}}_5^{c\mu} + \frac{\sigma_8}{2} \epsilon^{\mu\nu\rho\sigma} \xi_\nu F_{\rho\sigma} + T^2 \sigma_9 \omega^\mu + T \sigma_{10} B^\mu. \quad (4.9)$$

Expression (4.9) is the final result of this subsection.

Note that positivity of the divergence of the entropy current implies a correction to the entropy density of the form

$$-u_\mu J_S^\mu = s + \sigma_8 \tilde{\mathcal{S}}_2 + \sigma_1 T \tilde{\mathcal{S}}_1 + \partial_\mu (c_0 T \zeta^\mu) - c_0 T \zeta^\mu u \cdot \partial u_\mu. \quad (4.10)$$

Equation (4.10) implies that in a parity violating system the entropy density, as defined via the Gibbs Duhem relation (i.e., the log of the partition function in equilibrium), would receive gradient corrections if c_0 , σ_1 or σ_8 are non zero.

4.3 Constraints from positivity of the quadratic form

The divergence of the sum of the canonical entropy current and the the new piece (4.9) has no two derivative or curvature terms, and so is a quadratic form in one derivative data. In this subsection we will determine the corresponding quadratic form. The requirement that this quadratic form is positive will enable us to deduce the restrictions on the σ_i 's in (4.9) and on the parameters describing the equations of motion of the theory.

From (3.9) and taking into account the anomaly, we find that

$$\partial_\mu J_S^\mu = -\partial_\mu \left(\frac{u_\nu}{T} \right) T_{diss}^{\mu\nu} + \left(\frac{E_\mu}{T} - \partial_\mu \left(\frac{\mu}{T} \right) \right) J_{diss}^\mu + \frac{\mu_{diss}}{T} \partial_\mu (f \xi^\mu) + \partial_\mu J_{S_{\text{new}}}^\mu - c\nu E \cdot B. \quad (4.11)$$

On general grounds the divergence of the non canonical part of the entropy current, together with the contribution from the anomaly, takes the form:¹⁶

$$\partial_\mu J_{S_{\text{new}}}^\mu - c\nu E \cdot B = \sum_{i=1}^7 \sum_{j=1}^7 \tilde{\mathcal{V}}_i^c \cdot B_{ij}^v \mathcal{V}_j^c + \sum_{i=1}^7 \sum_{j=1}^2 \tilde{\mathcal{S}}_i^c B_{ij}^s \mathcal{S}_j. \quad (4.12)$$

Recall that $\tilde{\mathcal{V}}^{c\mu} = *\mathcal{V}_\mu^c$ so that B_{ij}^v is an antisymmetric 7×7 matrix.

The expressions for B_{ij}^v and B_{ij}^s are straightforward but rather tedious to compute. We provide some details relevant to the computation in Appendix C. We will not completely list these matrices here, but will unveil relevant aspects of their explicit form as we go along.

To complete our analysis, we need to reexpress the divergence of the canonical entropy current (3.9) in terms of on-shell independent data. We carry out this computation in a frame invariant manner, similar to that of section 2.1.

We find it convenient to decompose the explicit fluid first derivative data that appears in (3.9), namely $\nabla_\mu \left(\frac{u_\nu}{T}\right)$ and $E_\mu/T - \partial_\mu \nu$ into $SO(2)$ invariant scalars, vectors, and tensors as follows

$$\begin{aligned} \nabla_\mu \left(\frac{u_\nu}{T}\right) &= \left(\frac{1}{3T} \nabla \cdot u - \frac{1}{2\zeta^2 T} \zeta \cdot \sigma \cdot \zeta\right) \tilde{P}_{\mu\nu} + \frac{1}{\zeta^2} \left(\frac{1}{3T} \nabla \cdot u + \frac{1}{\zeta^2 T} \zeta \cdot \sigma \cdot \zeta\right) \zeta_\mu \zeta_\nu \\ &\quad + \frac{u \cdot \partial T}{T^2} u_\mu u_\nu - \frac{1}{\zeta^2 T} \left(\zeta^\alpha u \cdot \nabla u_\alpha + \frac{\zeta^\alpha \partial_\alpha T}{T}\right) u_{(\mu} \zeta_{\nu)} \\ &\quad - \frac{1}{T} \left(u \cdot \nabla u_\alpha + \frac{\partial_\alpha T}{T}\right) u_{(\mu} \tilde{P}_{\nu)}^\alpha + \frac{2}{\zeta^2 T} \zeta_{(\mu} \mathcal{V}_{2\nu)} + \frac{1}{T} \mathcal{T}_{1\mu\nu} \\ \frac{E_\mu}{T} - \partial_\mu \nu &= \mathcal{V}_{1\mu} + u_\mu u \cdot \partial \nu + \zeta_\mu \frac{\zeta \cdot V_1}{\zeta^2}. \end{aligned} \quad (4.13)$$

We will find it convenient to organize the 7 scalars and three vectors above into two columns \mathbf{s}_d and \mathbf{v}_d ,

$$\mathbf{s}_d = \begin{pmatrix} \frac{1}{3T} \nabla \cdot u - \frac{1}{2\zeta^2 T} \zeta \cdot \sigma \cdot \zeta \\ \frac{1}{3T} \nabla \cdot u + \frac{1}{\zeta^2 T} \zeta \cdot \sigma \cdot \zeta \\ \frac{u \cdot \partial T}{T^2} \\ -\frac{1}{\zeta^2 T} \left(\zeta^\alpha u \cdot \partial u_\alpha + \frac{\zeta^\alpha \partial_\alpha T}{T}\right) \\ -u \cdot \nabla \nu \\ -\frac{\zeta V_1}{\zeta^2} \\ \frac{\partial_\mu (f \xi^\mu)}{T} \end{pmatrix}, \quad \mathbf{v}_d = \begin{pmatrix} -\frac{1}{T} \left(u \cdot \nabla u_\alpha + \frac{\partial_\alpha T}{T}\right) \tilde{P}^\alpha{}_\nu \\ \frac{2}{\zeta^2 T} \mathcal{V}_{2\nu}^c \\ -\mathcal{V}_{1\nu}^c \end{pmatrix}. \quad (4.14)$$

¹⁶This follows from the observation that the final result is a pseudo scalar. The absence of a tensor times pseudo tensor term in this expression is not completely obvious but may be checked.

The pieces of scalar and vector data in (4.14) multiply particular scalar and vector components of $T_{diss}^{\mu\nu}$ and J_{diss}^μ . It is convenient to classify the scalar, vector and tensor combinations as

$$\begin{aligned}
\mathbf{s}_1 &= T_{diss}^{\mu\nu} \tilde{P}_{\mu\nu} & \zeta^2 \mathbf{s}_2 &= \zeta \cdot T_{diss} \cdot \zeta \\
\mathbf{s}_3 &= u \cdot T_{diss} \cdot u & \mathbf{s}_4 &= u \cdot T_{diss} \cdot \zeta \\
\mathbf{s}_5 &= u \cdot J_{diss} & \mathbf{s}_6 &= \zeta \cdot J_{diss} \\
\mathbf{s}_7 &= -\mu_{diss} \\
\mathbf{v}_1^\nu &= u_\mu T_{diss}^{\mu\alpha} \tilde{P}^\nu_\alpha & \mathbf{v}_2^\nu &= \zeta_\mu T_{diss}^{\mu\alpha} \tilde{P}^\nu_\alpha \\
\mathbf{v}_3^\nu &= \tilde{P}^\nu_\alpha J^\alpha \\
\mathbf{t} &= \tilde{P}_\mu^\alpha \tilde{P}_\nu^\beta T_{diss\ \alpha\beta} - \frac{1}{2} \tilde{P}^{\mu\nu} \tilde{P}^{\alpha\beta} T_{diss,\alpha\beta},
\end{aligned} \tag{4.15}$$

and to group these into row vectors

$$\begin{aligned}
\mathbf{s} &= (\mathbf{s}_1 \ \mathbf{s}_2 \ \mathbf{s}_3 \ \mathbf{s}_4 \ \mathbf{s}_5 \ \mathbf{s}_6 \ \mathbf{s}_7) \\
\mathbf{v} &= (\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3) .
\end{aligned} \tag{4.16}$$

These definitions permit a very simple formal expression for the divergence of the canonical entropy current:

$$\partial_\mu J_{S_{\text{canon}}}^\mu = -\mathbf{s} \ \mathbf{s}_d - \mathbf{v}_\mu \ \mathbf{v}_d^\mu - \frac{1}{T} \mathbf{t}_{\mu\nu} \mathcal{T}_1^{\mu\nu} . \tag{4.17}$$

In presenting (4.17) we have used matrix notation: the expressions on the right hand side of (4.17) are each the product of a row and a column.

Equation (4.17) is not our final expression for the divergence of the canonical entropy current for an important reason; the entries in the columns \mathbf{s}_d and \mathbf{v}_d are not independent on-shell. In fact, on-shell, each of the 7 \mathbf{s}_d is a linear combination of only four independent scalar terms. Similarly, on-shell, each of the three entries in \mathbf{v}_d is a linear sum over two on-shell vectors. Specifically

$$\mathbf{s}_d = A^s \mathcal{S}, \quad \mathbf{v}_d = A^v \mathcal{V} \tag{4.18}$$

where

$$\mathcal{S} = \begin{pmatrix} \mathcal{S}_1^c \\ \mathcal{S}_2^c \\ \mathcal{S}_3^c \\ \mathcal{S}_4^c \end{pmatrix} \quad \mathcal{V} = \begin{pmatrix} \mathcal{V}_{1\nu}^c \\ \mathcal{V}_{2\nu}^c \end{pmatrix} \tag{4.19}$$

and

$$A^s = \begin{pmatrix} \frac{Rs}{2q_n T\chi} & \frac{B_3}{3T} - \frac{A_3}{2T\chi} & \frac{B_2}{3T} - \frac{A_2}{2T\chi} & \frac{B_1}{3T} - \frac{A_1}{2T\chi} \\ -\frac{Rs}{q_n \chi T} & \frac{B_3}{3T} + \frac{A_3}{T\chi} & \frac{B_2}{3T} + \frac{A_2}{T\chi} & \frac{B_1}{3T} + \frac{A_1}{T\chi} \\ 0 & \frac{1}{T^2} & 0 & 0 \\ -\frac{R}{T^2\chi} & \frac{K_3}{T} & \frac{K_2}{T} & \frac{K_1}{T} \\ 0 & 0 & -1 & 0 \\ -\frac{1}{T^2\chi} & 0 & 0 & 0 \\ 0 & \frac{(\rho+P)K_3}{T} & \frac{(\rho+P)K_2}{T} & \frac{(\rho+P)K_1}{T} \end{pmatrix}, \quad A^v = \begin{pmatrix} -R & 0 \\ 0 & \frac{2}{T^3\chi} \\ -1 & 0 \end{pmatrix} \quad (4.20)$$

with

$$R = \frac{q}{\rho + P}. \quad (4.21)$$

Note that \mathcal{S} is a column composed of 4 of the 7 on-shell independent scalars listed in table 6, while \mathcal{V} is a column composed of 2 of the 6 on-shell independent vectors listed in the same table. Note also that A^s is a 7×4 matrix while A^v is a 3×2 matrix. The computation of the entries of the matrix A^s and A^v is outlined in Appendix C where we also provide expressions for A_i , B_i and K_i . It follows that when reexpressed in terms of on-shell independent data, the divergence of the canonical part of the entropy current takes the form

$$\partial_\mu J_{S \text{ canon}}^\mu = -\mathbf{s} A^s \mathcal{S} - \mathbf{v}_\mu A^v \mathcal{V}^\mu - \frac{1}{T} \mathbf{t}_{\mu\nu} \mathcal{T}_1^{\mu\nu}. \quad (4.22)$$

Equation (4.22) depends only on those 4 linear combinations of the 7 scalars \mathbf{s}_i that appear in the 4 columns of $\mathbf{s} A^s$. Similarly (4.22) depends on only the 2 linear combinations of the three vectors \mathbf{v}_i that appear in the 2 columns of $\mathbf{v} A^v$. These numbers could have been anticipated on general grounds. As in the case of ordinary fluid dynamics (see section 2) the equations of superfluid dynamics suffer from a field redefinition ambiguity. The field redefinitions in question are u^μ , T and μ , and may be decomposed into 3 $SO(2)$ scalars and one $SO(2)$ vector. All of the seven scalars \mathbf{s}_i described above transform under the scalar field redefinitions. As these redefinitions depend on three parameters, $7 - 3 = 4$ independent linear combinations of the scalars are field redefinition invariant. Similarly we expect to find $3 - 1 = 2$ field redefinition independent linear combinations of the three vectors \mathbf{v}_i . Since the canonical entropy current is field redefinition invariant, the expression for its divergence is necessarily also frame invariant. It follows that $\mathbf{s}_i A_{ij}^s$ and $\mathbf{v}_i A_{ij}^v$ must be linear combinations of the four frame invariant scalars and the two frame invariant vectors respectively. We have explicitly checked (see Appendix C) that the the four linear combinations $\mathbf{s} A^s$ and the 2 linear combinations $\mathbf{v} A^v$ are indeed field redefinition invariant.

Putting together (4.12) and (4.22), positivity of the entropy current implies that

$$-\frac{1}{T}\mathbf{t}_{\mu\nu}\mathcal{T}_1^{\mu\nu} \geq 0 \quad (4.23a)$$

$$-\mathbf{v}_i^\mu A_{ij}^v \mathcal{V}_j^c + \tilde{\mathcal{V}}_i^c \cdot B_{ij}^v \mathcal{V}_j^c \geq 0 \quad (4.23b)$$

$$-\mathbf{s}_i A_{ij}^s \mathcal{S}_j^c + \tilde{\mathcal{S}}_i^c B_{ij}^s \mathcal{S}_j^c \geq 0. \quad (4.23c)$$

In the rest of this section we will deduce the constraints imposed by (4.23) on the one derivative constitutive relations for the tensor $\mathbf{t}_{\mu\nu}$, the frame invariant vectors $\mathbf{v}_\mu A^v$ and the frame invariant scalars $\mathbf{s}A^s$.

4.3.1 Positivity in the Tensor Sector

The most general one derivative constitutive relation for the frame invariant tensor \mathbf{t} (on symmetry grounds) is given by

$$\mathbf{t}_{\mu\nu} = \tilde{P}_\mu^\alpha \tilde{P}_\nu^\beta T_{diss\alpha\beta} - \frac{1}{2}\tilde{P}^{\mu\nu}\tilde{P}^{\alpha\beta}T_{diss,\alpha\beta} = -\eta_i \mathcal{T}_i{}_{\mu\nu} - \tilde{\eta}_i \tilde{\mathcal{T}}_i{}_{\mu\nu} \quad (4.24)$$

where $\mathcal{T}_1 = \sigma^u$, $\mathcal{T}_2 = \sigma^\xi$ and $\tilde{\mathcal{T}}_i = *\mathcal{T}_i$ and the η 's and $\tilde{\eta}$'s are four possible coefficients. Positivity of the entropy current requires that

$$\eta_1 > 0 \quad \tilde{\eta}_1 \in \mathbf{R} \quad \eta_2 = \tilde{\eta}_2 = 0. \quad (4.25)$$

In other words, the tensor part of $T_{diss}^{\mu\nu}$ has no term proportional to \mathcal{T}_2 and to $*\mathcal{T}_2$. The latter result could have been anticipated. The divergence of the entropy current does not have a piece proportional to \mathcal{T}_2 . Therefore terms in the divergence of the entropy current which are linear in \mathcal{T}_2 must vanish. On the other hand $\mathbf{t}_{\mu\nu}$ does, in general, have a term proportional to \mathcal{T}_1 ; the coefficient of this term is the shear viscosity of the fluid and is constrained to be positive. Positivity also allows a term proportional to $*\mathcal{T}_1$. The coefficient of this term, $\tilde{\eta}_1$, is a new parity odd coefficient (it is of course an arbitrary function of the thermodynamical scalars). Note that $\tilde{\eta}_1$ drops out of the formula for entropy production. This is because the contraction of any tensor with its own star vanishes. Consequently, $\tilde{\eta}_1$ is non dissipative; its coefficient is unconstrained by the requirement of positivity of entropy production.

4.3.2 Positivity in the Vector Sector

The most general constitutive relations allowed by symmetries for the two frame invariant vectors takes the form

$$\mathbf{v}_i^\mu A_{ij}^v = -\mathcal{V}_i \kappa_{ij} - \tilde{\mathcal{V}}_i \tilde{\kappa}_{ij}. \quad (4.26)$$

The matrix of possible transport coefficients κ is a 7×2 matrix. The same is true for $\tilde{\kappa}$. The first index runs over a basis of on-shell inequivalent vectors, while the second index runs over a basis of field redefinition invariant combinations of vectors in $T_{diss}^{\mu\nu}$ and J_{diss}^μ .

A useful observation is that the quadratic form in the vector sector contains no terms proportional to the square of \mathcal{V}_i^c for $i = 3 \dots 7$. The first expression in (4.23b) has an explicit factor of either \mathcal{V}_1^c or \mathcal{V}_2^c . Since B_{ij}^v is antisymmetric, the second expression in (4.23b) has no terms proportional to the square of any of the on-shell independent vectors. Positivity thus demands that the quadratic form in the vector sector be completely independent of \mathcal{V}_i^c for $i = 3 \dots 7$. This implies that

$$B_{ij}^v = 0 \quad i = 3, \dots, 7, \quad j = 3, \dots, 7 \quad (4.27a)$$

$$\tilde{\kappa}_{ij} = -B_{ij}^v \quad i = 3, \dots, 7, \quad j = 1, 2 \quad (4.27b)$$

$$\kappa_{ij} = 0 \quad i = 3, \dots, 7, \quad j = 1, 2 \quad (4.27c)$$

We now proceed to investigate the implications of (4.27).

Let us first focus on (4.27a). A priori, this equality gives us a set of $5 \times 4/2 = 10$ partial differential equations that constrain the six independent σ_i coefficients in (4.9). One of these equations is a tautology. The remaining nine equations are as follows

$$\begin{aligned} B_{34}^v = 0 &\Rightarrow \frac{\partial(\sigma_4 - \sigma_8)}{\partial T} - \frac{1}{T} \frac{\partial\sigma_3}{\partial\nu} = 0 & B_{35}^v = 0 &\Rightarrow \frac{\partial\sigma_5}{\partial T} - \frac{1}{T} \frac{\partial\sigma_3}{\partial\chi} = 0 \\ B_{45}^v = 0 &\Rightarrow \frac{\partial(\sigma_4 - \sigma_8)}{\partial\chi} - \frac{\partial\sigma_5}{\partial\nu} = 0 & B_{37}^v = 0 &\Rightarrow \frac{\partial(\sigma_{10} - \nu\sigma_8)}{\partial T} + \frac{\sigma_3}{T} = 0 \\ B_{57}^v = 0 &\Rightarrow \frac{\partial(\sigma_{10} - \nu\sigma_8)}{\partial\chi} + \sigma_5 = 0 & B_{46}^v = 0 &\Rightarrow \frac{\partial\sigma_9}{\partial\nu} - 2\sigma_{10} + 2\nu\sigma_4 = 0 \\ B_{36}^v = 0 &\Rightarrow \frac{\partial\sigma_9}{\partial T} + 2\nu\frac{\sigma_3}{T} = 0 & B_{56}^v = 0 &\Rightarrow \frac{\partial\sigma_9}{\partial\chi} + 2\nu\sigma_5 = 0 \\ B_{47}^v = 0 &\Rightarrow \frac{\partial(\sigma_{10} - \nu\sigma_8)}{\partial\nu} + (\sigma_4 - \sigma_8) = c\nu. \end{aligned} \quad (4.28)$$

Equations (4.28) are far from independent. They admit the following two parameter set of solutions

$$\begin{aligned} \sigma_3 &= -T \frac{\partial(\sigma_{10} - \nu\sigma_8)}{\partial T} \\ \sigma_4 &= \sigma_8 + c\nu - \frac{\partial(\sigma_{10} - \nu\sigma_8)}{\partial\nu} \\ \sigma_5 &= -\frac{\partial(\sigma_{10} - \nu\sigma_8)}{\partial\chi} \\ \sigma_9 &= s_9 - \frac{2}{3}c\nu^3 + 2\nu(\sigma_{10} - \nu\sigma_8) \end{aligned} \quad (4.29)$$

In (4.29) we have chosen σ_8 and σ_{10} as our free parameters. The term s_9 in the last line of (4.29) is an integration constant. It turns out that the requirement of CPT invariance forces

$$s_9 = 0.$$

The argument for this conclusion is very similar to that given below (2.27). While we present all formulae below at nonzero s_9 , the reader should keep in mind that s_9 actually vanishes in any superfluid that enjoys invariance under CPT invariance, i.e. any superfluid that arises from a quantum field theory (and so presumably for any superfluid in the real world, or obtained via the AdS/CFT correspondence).

Equation (4.29) is the most general solution to (4.28). As an aside we note that (4.29) also implies that $B_{ij}^v = 0$ for $i, j = 1, 2$. We will use this information shortly.

We now turn to the implications of (4.27b). In this case, the coefficients $\tilde{\kappa}$ whose second index lies between 3 and 7 are completely determined in terms of B_{ij} , which is, in turn, a function of the two free parameters σ_8 and σ_{10} . We find that $B_{i2} = 0$ for $i = 3 \dots 7$. Thus,

$$\tilde{\kappa}_{i2} = 0 \quad i = 3, \dots, 7.$$

However $\tilde{\kappa}_{i1}$ are in general nonzero. They are given by

$$\begin{aligned} \tilde{\kappa}_{31} &= -B_{31}^v = -RT\sigma_3 - T\partial_T\sigma_8 \\ \tilde{\kappa}_{41} &= -B_{41}^v = -RT^2\sigma_4 - T\partial_\nu\sigma_8 \\ \tilde{\kappa}_{51} &= -B_{51}^v = -RT^2\sigma_5 - T\partial_\chi\sigma_8 \\ \tilde{\kappa}_{61} &= -B_{61}^v = -2RT^3\sigma_9 + 2T^2\sigma_{10} \\ \tilde{\kappa}_{71} &= -B_{71}^v = -RT^2\sigma_{10} + 2T\sigma_8 + cT\nu. \end{aligned} \tag{4.30}$$

To be clear we reiterate (4.27c)

$$\kappa_{ij} = 0 \quad i = 3, \dots, 7, \quad j = 1, 2. \tag{4.31}$$

Once we have implemented all these conditions, the quadratic form in the vector sector takes the form $\mathcal{V}_i^c \cdot \kappa_{ij} \mathcal{V}_j^c + \tilde{\mathcal{V}}_i^c \cdot \tilde{\kappa}_{ij} \mathcal{V}_j^c$ where $i, j = 1, 2$. Positivity of the entropy current implies

$$\sum_{i=1,2} \sum_{j=1,2} \left(\mathcal{V}_i^c \cdot \kappa_{ij} \mathcal{V}_j^c + \tilde{\mathcal{V}}_i^c \cdot \tilde{\kappa}_{ij} \mathcal{V}_j^c \right) \geq 0. \tag{4.32}$$

The Onsager relations (see [17]) imply that $\kappa_{12} = \kappa_{21}$ and $\tilde{\kappa}_{12} = -\tilde{\kappa}_{21}$. We are left with 6 independent transport coefficients κ_{11} , κ_{22} , $\kappa_{12} = \kappa_{21}$, $\tilde{\kappa}_{11}$, $\tilde{\kappa}_{22}$ and $\tilde{\kappa}_{12} = -\tilde{\kappa}_{21}$, each of which is an arbitrary function of μ , T and ξ , but is constrained by the inequality

that the matrix

$$\mathcal{K} = \begin{pmatrix} \kappa_{11} & 0 & \kappa_{12} & -\zeta\tilde{\kappa}_{12} \\ 0 & \kappa_{11} & \zeta\tilde{\kappa}_{12} & \kappa_{12} \\ \kappa_{12} & \zeta\tilde{\kappa}_{12} & \kappa_{22} & 0 \\ -\zeta\tilde{\kappa}_{12} & \kappa_{12} & 0 & \kappa_{22} \end{pmatrix} \quad (4.33)$$

is a positive semi-definite matrix. Owing to the high degree of symmetry of \mathcal{K} , this condition reads

$$\kappa_{11}\kappa_{22} \geq \kappa_{12}^2 + \zeta^2\tilde{\kappa}_{12}^2. \quad (4.34)$$

Note that the 3 κ transport coefficients occur even in parity preserving superfluids, while the 3 $\tilde{\kappa}$ coefficients violate parity and are new. Note also that two of the three new parity violating transport coefficients, $\tilde{\kappa}_{11}$ and $\tilde{\kappa}_{22}$, do not appear in the expression for entropy production, and so are non dissipative. However the third new coefficient $\tilde{\kappa}_{12}$ causes dissipation.

4.3.3 Positivity in the Scalar Sector

The most general expansion of the constitutive relations in the scalar sector takes the form

$$\mathbf{s}_i A_{ij}^s = - \sum_{i=1}^7 \mathcal{S}_i^c \beta_{ij} - \sum_{i=1}^2 \tilde{\mathcal{S}}_i \tilde{\beta}_{ij}, \quad (4.35)$$

implying the inequality

$$\sum_{i=1}^7 \sum_{j=1}^4 \mathcal{S}_i^c \beta_{ij} \mathcal{S}_j^c + \sum_{i=1}^2 \sum_{j=1}^4 \tilde{\mathcal{S}}_i \tilde{\beta}_{ij} \mathcal{S}_j^c + \sum_{i=1}^2 \sum_{j=1}^7 \tilde{\mathcal{S}}_i^c B_{ij}^s \mathcal{S}_j \geq 0. \quad (4.36)$$

Terms in (4.36) involving products of scalars and pseudo-scalars must vanish since they can not be positive definite for an arbitrary flow. Thus, focusing on the parity odd sector we must impose

$$\sum_{i=1}^2 \sum_{j=1}^4 \tilde{\mathcal{S}}_i \tilde{\beta}_{ij} \mathcal{S}_j^c + \sum_{i=1}^2 \sum_{j=1}^7 \tilde{\mathcal{S}}_i^c B_{ij}^s \mathcal{S}_j = 0. \quad (4.37)$$

Only the second piece on the right hand side of (4.37) involves \mathcal{S}_j^c with $j = 4 \dots 7$. We immediately deduce the equation

$$B_{ij}^s = 0 \quad = 1, 2, \quad j = 4, \dots, 7. \quad (4.38)$$

Equations (4.38) are automatically satisfied once we impose the solution (4.29); we have no further restrictions on the free parameters σ_8 and σ_{10} . For $j = 1, \dots, 4$, equation (4.37) implies

$$\tilde{\beta}_{ij} = -B_{ij}^s. \quad (4.39)$$

It follows that there are no undetermined parity odd transport coefficients in the scalar sector; all parity odd contributions to frame invariant scalar combinations of $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} are completely determined in terms of σ_8 and σ_{10} by (4.39) and the explicit listing

$$B_{ij}^s = \begin{pmatrix} \frac{2RT\sigma_9}{\chi} - \frac{2\sigma_{10}}{\chi} & -2\sigma_3 - 2T^2K_3\sigma_9 & -2T\sigma_4 - 2T^2K_2\sigma_9 & -2T\sigma_5 - 2K_1T^2\sigma_9 \\ -\frac{c\nu}{T\chi} - \frac{2\sigma_8}{T\chi} + \frac{R\sigma_{10}}{\chi} & \partial_T\sigma_8 - K_3T\sigma_{10} & \partial_\nu\sigma_8 - K_2T\sigma_{10} & \partial_\chi\sigma_{10} - K_1T\sigma_{10} \end{pmatrix}. \quad (4.40)$$

Note that $B_{11}^s = -\tilde{\kappa}_{61}/\zeta^2$ and $B_{21}^s = -\tilde{\kappa}_{71}/\zeta^2$ with $\tilde{\kappa}_{i1}$ defined in (4.30).

The remaining undetermined constitutive relations in the scalar sector are associated with the parity even coefficients. These are parameterized by β_{ij} where i and j both range between 1 and 4. The Onsager relations imply that the antisymmetric part of β_{ij} is zero. So we have a total of 10 parity even dissipative coefficients in the scalar sector. These 10 coefficients parameterize a symmetric matrix that is constrained to be positive

$$\sum_{i=1}^4 \sum_{j=1}^4 \mathcal{S}_i^c \beta_{ij} \mathcal{S}_j^c \geq 0. \quad (4.41)$$

5. Summary of results and special limits

In sections 3 and 4 we have built up a complete theory of relativistic superfluid hydrodynamics that may or may not enjoy a parity invariance. Our work generalizes the recent work of [10] (see also [16]) for parity conserving superfluids, and the work of [20] for charged gauge-theory fluids with triangle anomalies (see also [21, 22]).

The computations presented in sections 3 and 4 are lengthy and algebraically rather intensive. In this section we present the final results of our calculations in a manner that is self contained and makes no reference to the method of derivation.

5.1 The problem addressed

As discussed in section 3, the energy momentum tensor $T_{\mu\nu}$ and charged current J_μ of an s wave superfluid are given by

$$\begin{aligned} T^{\mu\nu} &= (\rho + P)u^\mu u^\nu + P\eta^{\mu\nu} + f\xi^\mu \xi^\nu + T_{diss}^{\mu\nu} \\ J^\mu &= q_n u^\mu - f\xi^\mu + J_{diss}^\mu \\ \mu \cdot \xi &= \mu + \mu_{diss} \end{aligned} \quad (5.1)$$

where ξ_μ is related to the background gauge potential A_μ and Goldstone Boson ψ through

$$\xi_\mu = -\partial_\mu \psi + A_\mu. \quad (5.2)$$

(ψ is the phase of the charged scalar condensate.) We often use the projected variable

$$\zeta^\mu = (\eta^{\mu\nu} + u^\mu u^\nu) \xi_\nu \equiv P^{\mu\nu} \xi_\nu. \quad (5.3)$$

in what follows. The equations of motion of superfluidity are given by

$$\begin{aligned} \partial_\mu T^{\mu\nu} &= F^{\nu\mu} J_\mu \\ \partial_\mu J^\mu &= c E_\mu B^\mu \\ \partial_\mu \xi_\nu - \partial_\nu \xi_\mu &= F_{\mu\nu} \end{aligned} \quad (5.4)$$

where we have allowed for the presence of triangle anomalies via the non conservation of the charged current. In writing (5.1) we have not specified a frame—a particular definition of the temperature T , chemical potential μ and four-velocity u^μ once we deviate from thermodynamic equilibrium. The thermodynamic functions ρ , P and f are functions of the variables

$$T, \quad \nu = \frac{\mu}{T}, \quad \chi = \frac{\mu^2 - \xi^2}{T^2}. \quad (5.5)$$

The equations (5.4) and (5.1) together specify a complete set of equations for the 9 fluid dynamical fields $u^\mu(x)$, $T(x)$, $\mu(x)$, $\xi^\mu(x)$ once $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} are specified as functions of the fluid dynamical fields and their derivatives. The equations that determine $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} in terms of derivatives of fluid dynamical fields are called constitutive relations. The main result of this paper is the determination of the most general form of the constitutive relations consistent with Lorentz invariance, positivity of the divergence of the entropy current and the Onsager relations.

The most general allowed constitutive relations turn out to depend on 20 unspecified functions of T , ν and χ which we will refer to as free parameters. In this section we will present our final results for the constitutive relations in terms of these 20 parameters.

Superfluid flows are also accompanied by an entropy current. The requirement that this entropy current be of positive divergence for every fluid flow played a key role in our derivation of the allowed form of the constitutive relations. At first order in the derivative expansion the entropy current takes the form

$$J_S = J_{S_{\text{canon}}}^\mu + J_{S_{\text{new}}}^\mu. \quad (5.6)$$

where $J_{S_{\text{canon}}}^\mu$ is the so called canonical entropy current (in a fluid frame)

$$J_{S_{\text{canon}}}^\mu = s u^\mu - \frac{\mu}{T} J_{diss}^\mu - \frac{u_\nu T_{diss}^{\mu\nu}}{T} \quad (5.7)$$

and $J_{S_{\text{new}}}^\mu$ is a correction proportional to a sum of single derivatives of fluid fields. In this section we also present our results for the correction $J_{S_{\text{new}}}^\mu$ to the entropy current.

After summarizing our results we proceed to explain the specialization of these results to the case of conformal or Weyl invariant superfluids, and certain simplified but often used limit where the superfluid velocity is relatively small—in liquid Helium once the superfluid velocity is too large superfluidity breaks down.

5.2 Listing our main result

The most general correction to the entropy current is given by

$$J_{S_{\text{new}}}^\mu = \epsilon^{\mu\nu\rho\sigma} \partial_\nu (\sigma_1 T u_\rho \zeta_\sigma) + \partial_\nu (c_0 \mathcal{Q}^{\nu\mu}) + \sigma_3 \tilde{\mathcal{V}}_3^{c\mu} + T \sigma_4 \tilde{\mathcal{V}}_4^{c\mu} + T \sigma_5 \tilde{\mathcal{V}}_5^{c\mu} + \frac{\sigma_8}{2} \epsilon^{\mu\nu\rho\sigma} \xi_\nu F_{\rho\sigma} + T^2 \sigma_9 \omega^\mu + T \sigma_{10} B^\mu. \quad (5.8)$$

where

$$\mathcal{Q}_{\mu\nu} = T(\xi_\mu u_\nu - \xi_\nu u_\mu), \quad (5.9)$$

and σ_3 , σ_4 , σ_5 and σ_9 are determined in terms of σ_8 and σ_{10} by

$$\begin{aligned} \sigma_3 &= -T \frac{\partial}{\partial T} (\sigma_{10} - \nu \sigma_8) \\ \sigma_4 &= \sigma_8 + c\nu - \frac{\partial}{\partial \nu} (\sigma_{10} - \nu \sigma_8) \\ \sigma_5 &= -\frac{\partial}{\partial \chi} (\sigma_{10} - \nu \sigma_8) \\ \sigma_9 &= s_9 - \frac{2}{3} c\nu^3 + 2\nu (\sigma_{10} - \nu \sigma_8) \end{aligned} \quad (5.10)$$

In the above relations ω and B were defined in (5.24), c is the anomaly coefficient and s_9 is an integration constant. In any CPT invariant theory

$$s_9 = 0.$$

The terms in the entropy current proportional to c_0 and σ_1 are divergence free and so are physically irrelevant (they have no effect on the equations of motion). Ignoring these terms, the final result for the entropy current is expressed in terms of σ_8 and σ_{10} which are undetermined functions of T , ν and χ . As we will see below, σ_8 and σ_{10} enter the constitutive relations, and so are free parameters.¹⁷

¹⁷In a parity preserving theory all the σ_i 's are forced to vanish. As a consequence, the correction to the entropy current in a parity even theory is restricted to the second term in (5.8) which is physically unimportant. For many practical purposes the entropy current in parity even theories may be taken to be uniquely fixed to its canonical form.

We now turn to our results for the constitutive relations. We express our results in a field redefinition invariant manner. Since the expressions $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} are not separately field redefinition invariant, we first list field redefinition invariant linear combinations of these quantities. Let

$$\begin{aligned}
\mathbf{s}_1 &= T_{diss}^{\mu\nu} \tilde{P}_{\mu\nu} & \zeta^2 \mathbf{s}_2 &= \zeta \cdot T_{diss} \cdot \zeta \\
\mathbf{s}_3 &= u \cdot T_{diss} \cdot u & \mathbf{s}_4 &= u \cdot T_{diss} \cdot \zeta \\
\mathbf{s}_5 &= u \cdot J_{diss} & \mathbf{s}_6 &= \zeta \cdot J_{diss} \\
\mathbf{s}_7 &= -\mu_{diss} \\
\mathbf{v}_1^\nu &= u_\mu T_{diss}^{\mu\alpha} \tilde{P}^\nu{}_\alpha & \mathbf{v}_2^\nu &= \zeta_\mu T_{diss}^{\mu\alpha} \tilde{P}^\nu{}_\alpha \\
\mathbf{v}_3^\nu &= \tilde{P}^\nu{}_\alpha J^\alpha \\
\mathbf{t} &= \tilde{P}_\mu{}^\alpha \tilde{P}_\nu{}^\beta T_{diss\ \alpha\beta} - \frac{1}{2} \tilde{P}^{\mu\nu} \tilde{P}^{\alpha\beta} T_{diss,\alpha\beta},
\end{aligned} \tag{5.11}$$

where

$$P^{\mu\nu} = \eta^{\mu\nu} + u^\mu u^\nu \quad \tilde{P}^{\mu\nu} = P^{\mu\nu} + \frac{\zeta^\mu \zeta^\nu}{\zeta^2}. \tag{5.12}$$

We define the row vectors

$$\begin{aligned}
\mathbf{s} &= (\mathbf{s}_1 \ \mathbf{s}_2 \ \mathbf{s}_3 \ \mathbf{s}_4 \ \mathbf{s}_5 \ \mathbf{s}_6 \ \mathbf{s}_7) \\
\mathbf{v} &= (\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3) .
\end{aligned} \tag{5.13}$$

We also define the matrices

$$A^s = \begin{pmatrix} \frac{Rs}{2q_n T \chi} & \frac{B_3}{3T} - \frac{A_3}{2T \chi} & \frac{B_2}{3T} - \frac{A_2}{2T \chi} & \frac{B_1}{3T} - \frac{A_1}{2T \chi} \\ -\frac{Rs}{q_n \chi T} & \frac{B_3}{3T} + \frac{A_3}{T \chi} & \frac{B_2}{3T} + \frac{A_2}{T \chi} & \frac{B_1}{3T} + \frac{A_1}{T \chi} \\ 0 & \frac{1}{T^2} & 0 & 0 \\ -\frac{R}{T^2 \chi} & \frac{K_3}{T} & \frac{K_2}{T} & \frac{K_1}{T} \\ 0 & 0 & -1 & 0 \\ -\frac{1}{T^2 \chi} & 0 & 0 & 0 \\ 0 & \frac{(\rho+P)K_3}{T} & \frac{(\rho+P)K_2}{T} & \frac{(\rho+P)K_1}{T} \end{pmatrix}, \quad A^v = \begin{pmatrix} -R & 0 \\ 0 & \frac{2}{T^3 \chi} \\ -1 & 0 \end{pmatrix} \tag{5.14}$$

where

$$R = \frac{q}{\rho + P} \quad V_1^\mu = \frac{E^\mu}{T} - P^{\mu\nu} \partial_\nu \nu$$

and the A_i 's B_i 's, C_i 's and K_i 's defined in appendix 4.

In terms of (5.11)-(5.14), the frame invariant scalar, vector and tensor combinations of $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} are given by the row vectors

$$\mathbf{s} A^s, \quad \mathbf{v}_\mu A^v, \quad \mathbf{t}_{\mu\nu}. \tag{5.15}$$

i	1	2	3	4	5	6	7
\mathcal{V}_i^μ	$\tilde{P}^{\mu\sigma} V_{1\sigma}$	$\tilde{P}^{\mu\beta} (\zeta^\alpha \sigma_{\alpha\beta})$	$\tilde{P}^{\mu\sigma} \nabla_\sigma T$	$\tilde{P}^{\mu\sigma} \nabla_\sigma \left(\frac{\mu}{T}\right)$	$\tilde{P}^{\mu\sigma} \nabla_\sigma \left(\frac{\zeta^2}{T^2}\right)$	$\frac{\mathcal{V}_2^{\zeta^\mu} - \tilde{P}^{\mu\alpha} \zeta^\nu \partial_\alpha u_\nu}{\zeta^2}$	$-\frac{P^{\mu\nu} F_{\nu\alpha} \zeta^\alpha}{\zeta^2}$
\mathcal{S}	$\zeta_\mu V_1^\mu$	$u \cdot \partial T$	$u \cdot \partial \frac{\mu}{T}$	$u \cdot \partial \frac{\zeta^2}{T^2}$	$\zeta^\mu P^{\mu\alpha} \partial_\alpha T$	$\zeta^\mu E_\mu$	$\zeta^\mu P^{\mu\alpha} \partial_\alpha \frac{\zeta^2}{T^2}$
$\mathcal{T}_{i\mu\nu}$	$*\sigma_{\mu\nu}^u$	$*\sigma_{\mu\nu}^\xi$	–	–	–	–	–
$\tilde{\mathcal{S}}_i$	$\omega \cdot \xi$	$B \cdot \xi$	–	–	–	–	–

Table 8: The basis in which we present our main results. The pseudo vectors and pseudo tensors $\tilde{\mathcal{V}}$ and $\tilde{\mathcal{T}}$ are given through $\tilde{\mathcal{V}}^\mu = \epsilon^{\mu\nu\alpha\beta} u_\nu \xi_\alpha \mathcal{V}_\beta$ and $\tilde{\mathcal{T}}^{\mu\nu} = \epsilon^{\mu\rho\alpha\beta} u_\rho \xi_\alpha \mathcal{T}_\beta^\nu + \epsilon^{\nu\rho\alpha\beta} u_\rho \xi_\alpha \mathcal{T}_\beta^\mu$. Here $\tilde{P}^{\mu\nu} = \eta^{\mu\nu} + u^\mu u^\nu - \zeta^\mu \zeta^\nu / \zeta^2$.

By scalars, vectors and tensors we mean expressions which transform as spin 0, ± 1 and ± 2 representations of the $SO(2)$ symmetry that is left invariant by the two vectors u^μ and ξ^μ at each point in spacetime.

We have 4 frame invariant scalars, 2 frame invariant vectors and one frame invariant tensor. The constitutive relations determine these quantities as functions of first derivative fluid expressions through the equations

$$\begin{aligned}
\mathbf{t}^{\mu\nu} &= -\eta \mathcal{T}_1^{\mu\nu} - \tilde{\eta} \tilde{\mathcal{T}}_1^{\mu\nu} \\
\mathbf{v}_i^\mu A_{ij}^v &= -\sum_{i=1}^2 \mathcal{V}_i \kappa_{ij} - \sum_{i=1}^2 \tilde{\mathcal{V}}_i \tilde{\kappa}_{ij} - \delta_{j1} \left(\sum_{i=3}^7 \tilde{\mathcal{V}}_i \tilde{\kappa}_{i1} \right) \\
\mathbf{s}_i A_{ij}^s &= -\sum_{i=1}^4 \sum_{j=1}^4 \mathcal{S}_i \beta_{ij} - \left(\sum_{i=1}^2 \sum_{j=1}^4 \tilde{\mathcal{S}}_i \tilde{\beta}_{ij} \right)
\end{aligned} \tag{5.16}$$

with \mathcal{T} , $\tilde{\mathcal{T}}$, \mathcal{V} , $\tilde{\mathcal{V}}$, \mathcal{S} and $\tilde{\mathcal{S}}$ a basis of onshell independent $SO(2)$ invariant tensors, vectors and scalars given in table 8. In equation (5.16) κ and β are symmetric matrices and $\tilde{\kappa}_{12} = -\tilde{\kappa}_{21}$. Those coefficients that occur in the big brackets on the right hand side of that equation, namely $\tilde{\kappa}_{1i}$ for $i = 3 \dots 7$ and $\tilde{\beta}_{ij}$ for $i = 1, 2$ and $j = 1 \dots 4$, are not free but are determined in terms of σ_8 and σ_{10} through the equations

$$\begin{aligned}
\tilde{\kappa}_{31} &= -RT\sigma_3 - T\partial_T\sigma_8 \\
\tilde{\kappa}_{41} &= -RT^2\sigma_4 - T\partial_\nu\sigma_8 \\
\tilde{\kappa}_{51} &= -RT^2\sigma_5 - T\partial_\chi\sigma_8 \\
\tilde{\kappa}_{61} &= -2RT^3\sigma_9 + 2T^2\sigma_{10} \\
\tilde{\kappa}_{71} &= -RT^2\sigma_{10} + 2T\sigma_8 + cT\nu
\end{aligned} \tag{5.17}$$

$$-\tilde{\beta}_{ij} = \begin{pmatrix} \frac{2RT\sigma_9}{\chi} - \frac{2\sigma_{10}}{\chi} & -2\sigma_3 - 2T^2K_3\sigma_9 & -2T\sigma_4 - 2T^2K_2\sigma_9 & -2T\sigma_5 - 2K_1T^2\sigma_9 \\ -\frac{c\nu}{T\chi} - \frac{2\sigma_8}{T\chi} + \frac{R\sigma_{10}}{\chi} & \partial_T\sigma_8 - K_3T\sigma_{10} & \partial_\nu\sigma_8 - K_2T\sigma_{10} & \partial_\chi\sigma_{10} - K_1T\sigma_{10} \end{pmatrix}. \quad (5.18)$$

The remaining coefficients in (5.16), those that occur outside the big brackets on the right hand side of (5.16), are free parameters. The symmetric 2×2 matrix κ , 4×4 matrix β , η and $\tilde{\kappa}_{12} = -\tilde{\kappa}_{21}$ are constrained to obey

$$\begin{aligned} \eta &> 0 \\ \kappa_{11}\kappa_{22} &\geq \kappa_{12}^2 + \zeta^2\tilde{\kappa}_{12}^2 \\ \beta_{ij} &\text{ is symmetric positive semi-definite.} \end{aligned} \quad (5.19)$$

The remaining parameters

$$\tilde{\kappa}_{11} \quad \tilde{\kappa}_{22} \quad \tilde{\eta} \quad \sigma_8 \quad \sigma_{10} \quad (5.20)$$

are unconstrained.

In the introduction we counted twenty free parameters. Let us enumerate them explicitly. The parity even parameters consist of a 4×4 symmetric matrix β_{ij} in the scalar sector (10 parameters), a 2×2 symmetric matrix κ_{ij} in the vector sector (3 parameters) and the shear viscosity η in the tensor sector (1 parameter) yielding a total of 14 parity even free parameters. These are the only parameters in a theory that conserves parity. The parity odd parameters consist of the three parameters $\tilde{\kappa}_{11}$, $\tilde{\kappa}_{22}$, $\tilde{\kappa}_{21} = -\tilde{\kappa}_{12}$, in the vector sector and the one parameter $\tilde{\eta}$ in the tensor sector, amounting to a total of 4 free parameters. The two undetermined functions σ_8 and σ_{10} are two additional free parameters.

Five of the six parity odd free parameters, namely $\tilde{\kappa}_{11}$, $\tilde{\kappa}_{22}$, $\tilde{\eta}$, σ_8 and σ_{10} are non dissipative, in the sense that they do not result in entropy production. The one derivative fluid data that these parameters multiply can be nonzero even in stationary solutions. On the other hand the remaining parity odd parameter $\tilde{\kappa}_{21} = -\tilde{\kappa}_{12}$ is dissipative, and this is also true of all 14 parity even constitutive parameters. These parameters cause entropy production. The requirement that these parameters produce rather than destroy entropy constrains them to satisfy (5.19).

After completing the work reported in this paper we received the preprint [33] which addresses the same question dealt with in this section. While we have not yet performed a detailed comparison between our results and those of [33], a superficial comparison of final results reveals several differences. We list some of these differences here.

1. The author of [33] finds a unique entropy current, while the entropy current in this paper has a two parameter (function) freedom.

2. The author of [33] finds a total of 2 new parity odd dissipative parameters, while we find 6 such parameters in this paper. The parameter we refer to as $\tilde{\eta}$ seems unaccounted for in [33].
3. The author of [33] reports that his new parameters are all functions of a single thermodynamical variable, while the six new dissipative parameters in this paper are all unconstrained functions of all three thermodynamic scalars.
4. The author of [33] finds no mixing between parity even and parity odd sectors, while in this paper we find nontrivial mixing between parity even and parity odd vector contributions.

We do not yet understand the reason for the discrepancy between our results and those of [33]. We leave an investigation of this issue to future work.

5.3 Weyl invariant superfluid dynamics

The equations of superfluidity simplify somewhat in conformal field theories, i.e. theories that enjoy invariance under Weyl transformations. The source of the simplification is twofold: first the stress tensor of a CFT is always traceless. Second, the dependence of all thermodynamic functions on temperature can be deduced by dimensional analysis if we take the remaining variables to be the dimensionless quantities ν and χ . In the rest of this subsection we briefly outline the special simplifications in Weyl invariant superfluid hydrodynamics.

Let us first consider the entropy current. The coefficient σ_3 in the expression (5.8) for the entropy current of a Weyl invariant theory must vanish. This follows because $\tilde{\mathcal{V}}_3$ does not transform homogeneously under Weyl transformations. Since σ_8 and σ_{10} are dimensionless and therefore independent of T , equation (5.10) does not pose an extra restriction on these two parameters.

Let us now turn to the consequences of Weyl invariance on constitutive relations. It turns out that (beside implying that all constitutive parameters are independent of T) Weyl invariance implies no special simplifications for constitutive relations in the tensor sector. In the vector sector we expect that $\tilde{\kappa}_{13} = 0$ since, like σ_3 , it multiplies $\tilde{\mathcal{V}}_3$ which does not transform homogeneously under Weyl transformations. Since σ_3 vanishes and σ_8 is independent of the temperature, this condition is automatic from equation (5.17), i.e. it does not pose an extra restriction on σ_8 . The remaining terms in the vector sector remain unchanged.

In the scalar sector dimensional analysis implies that

$$A_3 = K_3 = 0 \quad B_3 = -\frac{3}{T}. \quad (5.21)$$

The fact that the scalar \mathcal{S}_2 does not transform homogeneously under Weyl transformations implies that both indices of β in the third equation in (5.16) run only over the indices 1, 3, 4. Consequently β is effectively a 3×3 symmetric matrix (with 6 parameters) in the Weyl invariant case. More importantly, the tracelessness of the stress tensor implies that the number of scalar terms in $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} is six rather than seven. In other words, the seven terms \mathbf{s}_i listed in (5.11) are restricted by the linear relation

$$\mathbf{s}_1 + \mathbf{s}_2 - \mathbf{s}_3 = 0. \quad (5.22)$$

Since there are still three field redefinitions in the scalar sector, this implies that we have three rather than four field redefinition invariant scalar terms which can be constructed out of $T_{diss}^{\mu\nu}$, J_{diss}^μ and μ_{diss} . Accordingly we find that

$$\begin{aligned} \mathbf{s}_i A_{i2}^s &= 0 \\ \tilde{\beta}_{i2} &= 0 \end{aligned} \quad (5.23)$$

so that the row vector $\mathbf{s}_i A_{ij}^s$ has three rather than four independent entries. Equation (5.23) follows from (5.21), (5.22) and $\sigma_3 = 0$.

It follows that Weyl invariant superfluid hydrodynamics is governed by 10 constitutive parameters (when it preserves parity) and 16 constitutive parameters (when it violates parity).

5.4 The $\zeta \rightarrow 0$ limit

In liquid Helium, superfluidity breaks down once the velocity of the superfluid component is too large relative to the velocity of the normal component. The projected superfluid velocity $\zeta^\mu = P^{\mu\nu} \xi_\nu$ can be thought of as a control parameter for that limit. Once $\zeta = 0$ one can locally boost a fluid element to a frame where neither its normal component nor its superfluid component are in motion.

When ζ is non zero, the velocity of the normal component u^μ together with ζ^μ break the $SO(3, 1)$ symmetry to $SO(2)$. In the $\zeta \rightarrow 0$ limit (the collinear limit) this symmetry is enhanced to $SO(3)$. The number of possible expressions for $SO(3)$ invariant scalars, tensors, vectors and their parity odd counterparts is significantly smaller than that of the $SO(2)$ symmetric case. As discussed in [16, 10] there are two possible scalars, $\partial \cdot u$ and $\partial_\mu (f \xi^\mu)$ (which we can swap with $u \cdot \partial T$ and $u \cdot \partial \nu$ via the equations of motion), one vector $V_1^\mu = -P^{\mu\nu} \partial_\nu \nu + E^\mu / T$ and two tensors in the parity even sector, $\sigma_{\mu\nu}$ and the symmetric traceless projection of $\partial_\mu \zeta_\nu$ onto the space orthogonal to u_μ . In the parity odd sector we have two pseudo vectors

$$\omega^\mu = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} u_\nu \partial_\rho u_\sigma \quad \text{and} \quad B^\mu = \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} u_\nu F_{\rho\sigma}. \quad (5.24)$$

Thus, in the collinear limit the transport coefficients of the theory reduce to a 2×2 symmetric matrix β_{ij} (in the parity even scalar sector), a diffusion coefficient κ (in the parity even vector sector), a shear viscosity η (in the parity even tensor sector), and two parity odd terms $\tilde{\kappa}_\omega$ and $\tilde{\kappa}_B$. These enter the constitutive relations through

$$\begin{aligned}
-\frac{\partial \cdot u}{3T} T_{diss}^{\mu\nu} P_{\mu\nu} - \frac{u \cdot \partial T}{T^2} u \cdot T_{diss} \cdot u + u \cdot \partial \nu u \cdot J_{diss} + \frac{\mu_{diss}}{T} \partial_\mu (f \xi^\mu) &= S_i \beta_{ij} S_j \\
(R P_\alpha^\mu T_{diss}^{\alpha\nu} u_\nu + P_\alpha^\mu J^\alpha) &= \kappa V_1^\mu - \tilde{\kappa}_B B^\mu - \tilde{\kappa}_\omega \omega^\mu \\
P_\alpha^\mu P_\beta^\nu T_{diss}^{\alpha\beta} - \frac{1}{3} P^{\mu\nu} P_{\alpha\beta} T_{diss}^{\alpha\beta} &= -\eta \sigma^{\mu\nu}.
\end{aligned} \tag{5.25}$$

It must also be true that

$$J_S^\mu{}_{new} = \sigma_\omega \omega^\mu + \sigma_B B^\mu + \dots \tag{5.26}$$

where we have written only the odd part of the entropy current, omitting the total derivative term in the parity even sector (the second term on the right hand side of (5.8)). This omission is represented by the \dots in (5.26).

As we have explained, we expect (5.25) and (5.26) to hold on general grounds. We should, therefore, be able to verify these equations—and read off the values of $\tilde{\kappa}_B$, $\tilde{\kappa}_\omega$ —by studying the $\zeta \rightarrow 0$ limit of our general results. We now describe how to take this limit.

Let us first consider the entropy current. The third, fourth and fifth terms in (5.8) each have an explicit factor of ζ and so simply vanish in the $\zeta \rightarrow 0$ limit. (We assume that all physical quantities like the entropy current—and therefore all all coefficient functions that appear in the entropy current—are analytic functions of ζ .) The last two terms in (5.8) are already proportional to ω^μ and B^μ respectively. The sixth term in (5.8) is proportional to B^μ as $\zeta \rightarrow 0$, while the first term has a piece proportional to ω^μ and a piece proportional to B^μ . Putting it all together we recover the form (5.26) with

$$\begin{aligned}
\sigma_\omega &= T^2 \left(s_9 - \frac{2}{3} c \nu^3 + 2\nu \left(\sigma_{10} - \nu \sigma_8 - \frac{1}{2} \sigma_1 \right) \right) \\
\sigma_B &= T \left(\sigma_{10} - \nu \sigma_8 - \frac{1}{2} \sigma_1 \right).
\end{aligned} \tag{5.27}$$

where all functions are evaluated, of course, at $\zeta^2 = 0$.

Let us now turn to the constitutive relations. Notice that $-(\mathbf{v}A^v)_1$ is simply the $SO(2)$ vector part of the $SO(3)$ frame invariant vector presented on the left hand side

of the second equation in (5.25). Note also that at leading order in small ζ

$$\frac{2T(\mathbf{s}A^s)_2}{A_3} \sim \frac{2T(\mathbf{s}A^s)_2}{A_2} \sim \frac{2T(\mathbf{s}A^s)_1}{A_1} \sim \frac{2\mathbf{s}_2 - \mathbf{s}_1}{\chi}.$$

Since the constitutive relations (5.16) equate this combination to an analytic function of ζ , it follows that in the $\zeta \rightarrow 0$ limit $2\mathbf{s}_2 - \mathbf{s}_1$ is frame invariant and, in fact, vanishes at $\zeta = 0$.¹⁸ Using this fact it follows that $-(\mathbf{s}A^s)_1$ is proportional to the scalar part of the vector on the left hand side of the second equation in (5.25). Thus,

$$\begin{aligned} \kappa &\geq 0 \\ \eta &\geq 0 \\ \beta_{ij} &\text{ is positive semi-definite} \\ \tilde{\kappa}_\omega = -\tilde{\kappa}_{61} &= -\zeta^2\beta_{11} = 2RT^3s_9 - \frac{4}{3}cRT^3\nu^3 - 4RT^3\nu^2\sigma_8 + 2T^2(2RT\nu - 1)\sigma_{10} \\ \tilde{\kappa}_B = -\tilde{\kappa}_{62} &= -\zeta^2\beta_{21} = RT^2\sigma_{10} - 2T\sigma_8 - cT\nu. \end{aligned} \tag{5.28}$$

It was necessary that $-\tilde{\kappa}_{61} = -\zeta^2\beta_{11}$ and that $\tilde{\kappa}_{62} = -\zeta^2\beta_{21}$ in order that our results group into an $SO(3)$ vector in the $\zeta \rightarrow 0$ limit. This gives us a mild consistency check on our results.

Let us summarize. The parity odd contributions to the entropy current and constitutive relations are much simpler in the collinear limit than in the general case. The only constitutive relation that receives corrections in this limit is the expression for the frame invariant vector listed in the left hand side of the second equation of (5.25) (equal to the dissipative part of the charge current in a Landau like frame). The corrections to both this vector and the entropy current are linear combinations of B^μ and ω^μ . These linear combinations are specified in terms of 3 free functions, σ_8 , σ_{10} and σ_1 and one integration constant (which vanishes on assuming CPT symmetry). As 4 coefficient parameters are determined in terms of 3 parameters, the coefficients obey a single linear identity. This identity is given by

$$\frac{1}{2}\sigma_\omega - \mu\sigma_B = -\frac{\mu^3}{3T}c + \frac{1}{2}s_9T^2 \tag{5.29}$$

Under the eminently reasonable assumption of CPT invariance $s_9 = 0$ and

$$\frac{1}{2}\sigma_\omega - \mu\sigma_B = -\frac{\mu^3}{3T}c \tag{5.30}$$

¹⁸This could have been anticipated on general grounds, as this combination is proportional to the $SO(2)$ scalar in the $SO(3)$ stress tensor, i.e. it is proportional to $\zeta \cdot \sigma \cdot \zeta$.

The relation (5.30) is the only real prediction in the $SO(3)$ invariant sector. Note that it constrains coefficients in the entropy current alone. In the next section we will use the AdS/CFT correspondence to test this relationship (of course $s_9 = 0$ in the holographic context).

The $\zeta \rightarrow 0$ for Weyl invariant superfluids

As we have seen above, Weyl invariance removes one of scalar field redefinition invariant combinations of one derivative corrections to the constitutive relations (because the trace of the stress tensor is set to zero) and also removes one of the scalars in terms of which the parity even part of these constitutive relations is expressed (because $u \cdot \partial T$ and $\partial \cdot u$ do not transform homogeneously under Weyl rescalings). In more detail, since in a conformal field theory, $q_n \left(\frac{\partial}{\partial \nu} \ln \frac{q_n}{s} \right) u \cdot \partial \nu = \partial_\mu (f \xi^\mu)$, there is only one permissible on-shell scalar. In the $\zeta \rightarrow 0$ limit the 2×2 matrix β described above collapses to a 1×1 matrix. Thus Weyl invariant superfluid dynamics is characterized by 3 constitutive parameters in the $\zeta \rightarrow 0$ limit, one in the tensor sector (the shear viscosity), one in the vector sector (conductivity) and one in the scalar sector (a new coefficient).

Apart from determining the dependence of $\tilde{\kappa}_\omega$, $\tilde{\kappa}_B$, σ_ω and σ_B on the temperature (using dimensional analysis), Weyl invariance does not impact the discussion presented above for parity odd corrections to constitutive relations and the entropy current. In particular (5.25)-(5.30) continue to hold, and do not significantly simplify in a theory that enjoys Weyl invariance.

6. A holographic computation

The AdS/CFT correspondence [35, 36, 37] can be used to study the superfluid phase of appropriate large N gauge theories in the limit of infinite 't Hooft coupling [38, 39, 34, 40, 41]. We will refer to such phases of gauge theories as holographic superfluids. Various features of holographic superfluids have been studied in the literature. These include sound modes [42, 43, 41, 44, 16], critical superfluid velocities [34, 45] and vortex structure [46, 47, 48]. Holographic superfluids with a non uniform temperature, chemical potential and velocity fields have been recently constructed in [16, 10] (see also [9]). These dynamical superfluids were constructed for a certain specialized bulk action which, following the work of [30], allows for an analytic treatment.

In this subsection we will demonstrate that the prediction (5.30), that follows from our general analysis, is indeed true for superfluids that admit a dual description via the AdS/CFT correspondence. Our demonstration uses the technology of the so called fluid gravity correspondence [24, 25, 26, 27, 28, 21, 22, 29] but allows for the implementation of this map in an abstract manner that does not require us to know the explicit form

of the solutions dual even to stationary fluid flows. In this respect our analysis is reminiscent of the thermodynamical analysis of Sonner and Withers [9].

Our starting point for the analysis is the bulk action

$$S = S_{EH} + S_{matter} + S_{CS} \quad (6.1)$$

where

$$\begin{aligned} S_{EH} &= \frac{1}{2\kappa^2} \int \sqrt{-g} (R + 12) \\ S_{matter} &= \frac{1}{2\kappa^2} \int \sqrt{-g} \left(-\frac{1}{4} V_F(|\psi|) F_{mn} F^{mn} - V_\psi(|\psi|) |\partial_m \psi - iq A_m \psi|^2 - V(|\psi|) \right) \\ S_{CS} &= \frac{c}{24} \int \sqrt{-g} \epsilon^{mnpqr} A_m F_{np} F_{qr} \end{aligned} \quad (6.2)$$

with $\epsilon^{01234} = 1/\sqrt{-g}$, $F = dA$, $V_F(0) = 1$, $V_\psi(0) = 1$ and $V(0) = 0$. We will often use $D_m \psi = \partial_m \psi - iq A_m \psi$. Roman indices run from 0 to 3 and 5. Later we will use Greek indices to denote the boundary coordinates $\mu = 0, \dots, 3$ and $i, j = 1, \dots, 3$ will denote the spatial coordinates along the boundary. We have chosen the action (6.1) for several reasons. As has been discussed in detail in [38, 39, 34, 40], to construct a holographic superfluid one needs to spontaneously break a gauge symmetry in an asymptotically AdS geometry. The minimal required fields for such a construction involve a gauge field A_μ , a charged scalar ψ and a metric $g_{\mu\nu}$. The parity conserving part of the action (6.1), i.e., $S_{EH} + S_{matter}$, is the most general action one can write down which involves the fields described above. The parity violating sector consists of a Chern-Simons term, S_{CS} , which naturally appears in consistent truncations of type IIB supergravity. For example, the actions of the consistent supergravity truncations described in [31, 32] can be described by the action (6.1).

The equations of motion which follow from (6.1) are

$$\begin{aligned} \frac{1}{\sqrt{-g}} ((\partial_m - iq A_m) V_\psi \sqrt{-g} D^m \psi) &= \frac{1}{4} \frac{\partial V_F}{\partial \psi^*} F^2 + \frac{\partial V_\psi}{\partial \psi^*} |D\psi|^2 + \frac{\partial V}{\partial \psi^*} \\ \frac{1}{\sqrt{-g}} \partial_m (\sqrt{-g} V_F F^{mn}) &= iq V_\psi (\psi^* (D^n \psi) - \psi (D^n \psi)^*) - \frac{c\kappa^2}{4} \epsilon^{nmpqs} F_{mp} F_{qs} \\ R_{mn} - \frac{1}{2} R g_{mn} - 6g_{mn} &= \mathcal{T}_{mn} \end{aligned} \quad (6.3)$$

with

$$\begin{aligned} \mathcal{T}_{mn} = & \frac{1}{2}V_F F_{mp} F^{np} - \frac{1}{8}g_{mn} F_{pq} F^{pq} - \frac{1}{2}g_{mn} V - \frac{1}{2}g_{mn} V_\psi |D\psi|^2 \\ & + \frac{1}{2}V_\psi ((D_m\psi)(D_n\psi)^* + (D_n\psi)(D_m\psi)^*) . \end{aligned} \quad (6.4)$$

Our strategy for computing the parity odd corrections to the entropy current and the parity odd transport coefficients closely follows that of [24, 26, 21, 22, 16, 10]. We first construct a stationary solution to the equations of motion (6.3) in which neither the superfluid nor the normal component are in motion. Being interested in the limit of small relative superfluid velocity, we then allow for the gauge field to have a small non vanishing spatial component. By boosting and rotating the latter solution, we find a generic bulk configuration involving six free parameters which are associated with the temperature, chemical potential, velocity of the normal component and the expectation value of the goldstone boson of the dual superfluid. We then promote these six parameters to be spacetime dependent fields. Once we do so the equations of motion (6.3) will no longer be satisfied. Therefore, we add corrections to the our previous solution so that the equations of motion (6.3) will be satisfied at least up to second order gradients of the six parameters. We can then compute the one point functions of the energy momentum tensor, charged current and gradient of the Goldstone boson in the boundary theory and from them read off the transport coefficients associated with the first order terms in a gradient expansion.¹⁹

6.1 A stationary solution

The ansatz for a stationary solution to the equations of motion (6.3)—corresponding to a configuration in which the normal fluid and superfluid are both at rest—is

$$ds^2 = -r^2 f(r) dt^2 + r^2 dx^2 + \sigma(r) dt dr \quad (6.5)$$

$$\psi = \rho(r) e^{iq\varphi(r)} \quad (6.6)$$

$$A_m = (A_0(r), 0, 0, 0, A_5(r)) . \quad (6.7)$$

In this ansatz we have not completely fixed a gauge. In what follows we will present our analysis in the gauge invariant variables

$$G_m = A_m - \partial_m \varphi . \quad (6.8)$$

In this section we will not attempt to find an explicit solution for the unknown functions in (6.5), such a solution will not be needed in what follows below. We will

¹⁹The authors of [49] recently argued that the transport coefficients associated with the parity odd sector can also be computed using a Kubo formula.

instead note certain abstract aspects of this solution that can be obtained without explicitly solving the equations.

- The equation for G_5 is algebraic and is given by

$$G_5 = -\frac{G_0\sigma}{r^2 f}. \quad (6.9)$$

- The remaining equations of motion can be chosen to be a set of 4 linear differential equations in the variables f , σ , ρ and G_0 . The equations have the property that they involve only first derivatives (in the variable r) of f and σ and second order derivatives (in the variable r) of ρ and G_0 .
- Assuming that the solution is a black hole, the fields σ and ρ are non vanishing at r_h , the black hole horizon, while f and G_0 have a simple zero at r_h . The fact that G_0 vanishes at r_h follows from regularity of G_5 (defined in (6.9)) near the horizon. The temperature, T , and entropy density s of the black hole are given by

$$T = \frac{r_h^2 f'(r_h)}{4\pi\sigma(r_h)} \quad s = \frac{\pi r_h^3}{\kappa^2}. \quad (6.10)$$

- One linear combination of the 4 equations can be shown to express the constancy (in the radial direction) of the ‘Noether’ charge Q_1 [50],

$$2\kappa^2 Q_1 = \frac{r^5 f'}{\sigma} - \frac{r^3 V_F G G'}{\sigma}. \quad (6.11)$$

The equation in question asserts that

$$Q_1' = 0$$

where a prime denotes a derivative in the r direction. Using (6.10) we find that

$$Q_1 = sT. \quad (6.12)$$

- It is not difficult to work out a Graham Fefferman style solution of the equations of motion at large r . Putting in the physical constraint that the non normalizable mode dual to the scalar is turned off (i.e. that the operator dual to the scalar field is not sourced in the dual boundary theory), we can use (6.11) and the equations of motion for f , σ , ρ and G_0 to show that near the asymptotically AdS boundary

($r \rightarrow \infty$)

$$\begin{aligned}
f &= 1 + \frac{1}{b^4 r^4} + \mathcal{O}(r^{-5}) \\
\sigma &= 1 - \frac{1}{6} C_\Delta |\langle O_\psi \rangle|^2 \Delta r^{-2\Delta} + \mathcal{O}(r^{-2\Delta-2}) \\
\rho &= r^{-\Delta} (C_\Delta |\langle O_\psi \rangle| + \mathcal{O}(r^{-2})) \\
G_0 &= \mu - \kappa^2 q_t r^{-2} + \mathcal{O}(r^{-3})
\end{aligned} \tag{6.13}$$

where C_Δ is a real number, O_ψ is the operator dual to ψ , and q_t represents the total charge density as seen in the rest frame of the normal component; using the standard AdS/CFT prescription [36, 35] to convert bulk to boundary quantities we find that

$$q_t = -u^\mu J_\mu$$

with J_μ the boundary theory charged current. Using (6.11) together with (6.12) we find

$$sT + \mu q_t = \frac{2}{b^4 \kappa^4} \equiv 4P \tag{6.14}$$

which is the Gibbs Duhem relation for a conformal superfluid (see (3.6)).

6.2 Adding in a small uniform superfluid velocity

As we have explained above, in this section we focus on the collinear limit, i.e. the limit in which the normal and superfluid velocities are equal. Following a long tradition we do, however, wish to allow the derivatives of the normal and superfluid velocities to be independent variables. Thus while we can set the normal and superfluid velocities equal at a given point, it would be inconsistent to do the same in the neighborhood of that point. In order to implement the fluid gravity map, even in our limited context, we need more general stationary background solutions than the ones described in the previous paragraph. In particular, since we are interested only in first derivative corrections, we need control over background solutions in which the fluid is at rest and the superfluid in motion to linear order in the superfluid velocity. In this section we describe the relevant solutions.

A constant superfluid velocity is an $SO(3)$ vector. The solution that describes infinitesimal superfluid motion (at first order) is a regular, normalizable, translationally invariant fluctuation about the static background described in the previous subsection. A vector fluctuation involves linear fluctuations of the vector modes g_{ti} and G_i . We find it more convenient to work in terms of the variables g and γ defined by

$$G_i = -g \partial_i \phi \quad g_{ti} = -r^2 \gamma \partial_i \phi$$

where $\partial_i\phi$ is the expectation value of the spatial component of the Goldstone boson in the dual field theory (i.e. $\partial_\mu\phi$ is a function of the four boundary rather than the 5 bulk dimensions).

From the Einstein equations we obtain the following equation of motion for γ at linear order

$$\left(\frac{\gamma'r^5 + G'_0gr^3V_F}{\sigma}\right)' = 0. \quad (6.15)$$

Integrating (6.15) we obtain

$$2\kappa^2Q_2 = \frac{\gamma'r^5 + G'_0gr^3V_F}{\sigma} \quad (6.16)$$

with Q_2 an integration constant. The asymptotic behavior of γ can be related to thermodynamic quantities in the solution via

$$\gamma = \frac{1}{2}(q_t - q)r^{-4} + O(r^{-5}). \quad (6.17)$$

Integrating (6.16) near the boundary, it is not difficult to verify the form (6.17) and to check that

$$Q_2 = q. \quad (6.18)$$

The equation of motion for g is obtained from the spatial components of the Maxwell equation and can also be written as a total derivative,

$$\left(\frac{fr^3(gG'_0 - g'G_0)V_F}{\sigma} + 2\kappa^2\gamma Q_1\right)' = 2\kappa^2f'Q_2. \quad (6.19)$$

Integrating it once we get

$$\frac{fr^3(gG'_0 - g'G_0)V_F}{\sigma} + 2\kappa^2\gamma Q_1 = Q_3 + 2\kappa^2fQ_2 \quad (6.20)$$

with Q_3 an integration constant. We will now argue that Q_3 vanishes. The argument follows by demanding regularity at the event horizon. The fact that r_h is an event horizon (and so that dr is a null one-form at $r = r_h$) implies that

$$\gamma(r_h) = 0. \quad (6.21)$$

Every term in (6.20) vanishes at $r = r_h$ except Q_3 , so we conclude that $Q_3 = 0$ for $T > 0$. The near boundary expansion of g then takes the form

$$g = 1 - \frac{(q_t - q)\kappa^2}{\mu}r^{-2} + O(r^{-3}). \quad (6.22)$$

The second order linearized equations (6.15) and (6.19) have four linearly independent solutions. One of these is

$$g = 0 \quad \gamma = \text{constant} \quad (6.23)$$

which corresponds to a deformation of the dual field theory metric. Another solution is

$$g = G_0 \quad \gamma = \frac{Q_2}{Q_1} f \quad (6.24)$$

which corresponds to a boost of the stationary solution. Of the remaining two linearly independent solutions only one can be chosen to be regular at the horizon. In terms of the two first order integrated equations (6.16) and (6.20), imposing $Q_2 \neq 0$ precludes (6.23) and imposing $Q_3 = 0$ removes the solution which diverges at the horizon.

6.3 Boosting

So far we have worked in a frame where $u^\mu = (1, 0, 0, 0)$. As explained in [16] we can always boost to a frame where the metric and gauge field take the form

$$\begin{aligned} ds^2 &= -r^2 f(r) u_\mu u_\nu dx^\mu dx^\nu + r^2 P_{\mu\nu} dx^\mu dx^\nu - 2\sigma u_\mu dx^\mu dr + 2r^2 \gamma u_{(\mu} \bar{\zeta}_{\nu)} dx^\mu dx^\nu \\ A_\mu &= G u_\mu dx^\mu + g \bar{\zeta}_\mu dx^\mu + G_5 dr, \end{aligned} \quad (6.25)$$

with

$$\begin{aligned} G &= G_0 \\ \lim_{r \rightarrow \infty} A_\beta &= a_\beta \\ \bar{\zeta}^\alpha &= P^{\alpha\beta} (\partial_\beta \phi - a_\beta), \end{aligned} \quad (6.26)$$

a_β the external gauge field and ϕ the Goldstone boson. Our conventions here slightly differ from the main text where we've defined $\bar{\zeta} = -\zeta$, see equations (3.1), (3.2).

In our current holographic formulation, it is convenient to use a frame where the energy momentum tensor, charged current and Josephson condition are given by

$$\begin{aligned} T^{\mu\nu} &= (\rho + P + f\mu^2) u^\mu u^\nu + \eta^{\mu\nu} P + 2f\mu \bar{\zeta}_{(\mu} u_{\nu)} + f \bar{\zeta}_\mu \bar{\zeta}_\nu + \bar{T}_{diss}^{\mu\nu} \\ J^\mu &= q_t u^\mu + f \bar{\zeta}^\mu + \bar{J}_{diss}^\mu \\ u^\mu (\partial_\mu \phi - a_\mu) &= -\mu + \bar{\mu}_{diss} \equiv -\mu_T \end{aligned} \quad (6.27)$$

where

$$q_t = q + f\mu, \quad (6.28)$$

and our frame choice is $u_\mu \bar{T}_{diss}^{\mu\nu} = 0$ and $u_\mu \bar{J}_{diss}^\mu = 0$.²⁰ One can check that expanding (6.27) to linear order in $\bar{\zeta}^\mu$ and neglecting $\bar{T}_{diss}^{\mu\nu}$, \bar{J}_{diss}^μ and $\bar{\mu}_{diss}$, the boundary theory energy momentum tensor, charged current and chemical potential associated with the solution (6.25) are obtained.

6.4 The gradient expansion

We now have in hand all stationary solutions that will be needed for our analysis. As explained at the beginning of this section, the next step in our analysis is to allow the thermodynamic variables to depend on the spacetime coordinates and look for corrections to the metric, δg_{mn} , gauge field, δA_m and scalar $\delta\psi$, so that the equations of motion (6.3) are satisfied. Since we will be interested in the collinear limit, we will set $\bar{\zeta} = 0$ but keep its derivatives non zero. In this limit, the equations of motion for the corrections to the metric, gauge field and scalar naturally decompose themselves into scalar, vector and tensor equations under the $SO(3) \subset SO(3,1)$ symmetry retained by the background. We write the corrections to the metric in the form

$$\begin{aligned} \delta g_{mn} dx^m dx^n = & -2u_\mu dx^\mu r \left(u^\alpha \partial_\alpha u_\nu + \frac{1}{3} \partial_\alpha u^\alpha u_\nu \right) dx^\nu - 2\delta\sigma u_\mu dx^\mu dr \\ & - r^2 \delta f u_\mu dx^\mu u_\nu dx^\nu + 2r^2 \delta V_\mu u_\nu dx^\mu dx^\nu + r^2 \delta \pi_{\mu\nu} dx^\mu dx^\nu, \end{aligned} \quad (6.29)$$

the corrections to the gauge invariant combination G_m in the form

$$\delta A_m dx^m = -\delta G u_\mu dx^\mu + \delta g_\mu dx^\mu + \delta G_5 dr, \quad (6.30)$$

and corrections to the magnitude of the scalar as $\delta\rho$.

The equation of motion for δG_5 turns out to be an algebraic equation which can be solved. The remaining kinematic equations for δG , $\delta\rho$, δf and $\delta\sigma$ are coupled. The vector sector contains two coupled equations involving δV_μ and δg_μ and the equation of motion for the tensor mode $\delta\pi_{\mu\nu}$ can be solved for since it is decoupled from the

²⁰Actually, in our computation we are using a frame where

$$\begin{aligned} T^{\mu\nu} &= (\rho + P + f\mu^2)u^\mu u^\nu + \eta^{\mu\nu}P + 2f\mu\bar{\zeta}_{(\mu}u_{\nu)} + \frac{f\mu}{\mu_T}\bar{\zeta}_\mu\bar{\zeta}_\nu + \bar{T}_{diss}^{\mu\nu} \\ J^\mu &= q_t u^\mu + \frac{f\mu}{\mu_T}\bar{\zeta}^\mu + \bar{J}_{diss}^\mu \\ u^\mu (\partial_\mu\phi - a_\mu) &= -\mu + \bar{\mu}_{diss} \equiv -\mu_T. \end{aligned}$$

The dissipative corrections \bar{J}_{diss} , \bar{T}_{diss} and $\bar{\mu}_{diss}$ in this frame are different from J_{diss} , T_{diss} and μ_{diss} used throughout this work. Since in this section we will only be computing the transport coefficients in the vector sector, the difference between the two frames is immaterial. The reader is referred to [16, 10] for an extensive discussion.

rest of the equations. We have verified that the five constraint equations imply energy momentum conservation and current conservation in the dual field theory. We have also checked that the ratio of the shear viscosity to entropy density retains its universal value as has been alluded to in [51, 52, 16].

Since we are interested only in the parity odd sector, and since in the collinear limit this sector receives contributions only from the vector modes, we can focus entirely on the equations of motion for δV_μ and δg_μ which read

$$\left(\frac{r^5 \delta V'_\mu}{\sigma} - \frac{G' \delta g_\mu r^3 V_F}{\sigma} \right)' = \mathbf{S}_\mu^{\delta V} \quad (6.31a)$$

$$\left(\frac{V_F f r^3 (\delta g_\mu G' - \delta g'_\mu G)}{\sigma} - 2\kappa^2 Q_1 \delta V_\mu \right)' = \mathbf{S}_\mu^{\delta g}. \quad (6.31b)$$

(Recall that $G = G_0$ from (6.26).) Note that the homogeneous parts of these equations agree exactly with (6.15) as expected.

After integrating these once we can eliminate δV in place of δg and write the resulting differential equation in Sturm-Liouville form,

$$\left(\frac{r^3 f V_F}{\sigma} \delta g'_\mu \right)' - \left(\frac{r V_F^2 G'^2}{\sigma} + 2q^2 r \sigma V_\psi \rho^2 \right) \delta g_\mu = -\frac{\mathbf{S}_\mu^{\delta g}}{G} + \frac{2\kappa^2 Q_1 \sigma}{r^5 g} \int_r^\infty \mathbf{S}_\mu^{\delta V}(x) dx \equiv \mathbf{S}^{\text{total}}. \quad (6.32)$$

In order to simplify the term linear in δg_μ we used the equation of motion for G . The limits of integration on the right hand side of (6.32) have been determined by the boundary values of δg_μ and δV_μ and the requirement that we are in the frame described in (6.27).

Since (6.32) is a linear second order differential equation we can use the method of Green's functions to solve it. Consider first the homogeneous version of (6.32). We denote the two linearly independent solutions of the homogeneous equation by a and \hat{a} such that

$$a = 1 + \mathcal{O}(r^{-2}) \quad a(r_h) = a_h \quad (6.33)$$

$$\hat{a} = r^{-2} + \dots \quad \hat{a}(r_h) = \infty. \quad (6.34)$$

We find that

$$a = \frac{Q_1 g + Q_2 G}{Q_1 + Q_2 \mu}. \quad (6.35)$$

The solution (6.35) corresponds to a shift in the superfluid velocity described in the previous subsection. (One can use the method of variation of parameters to find \hat{a} in integral form though we will not be needing it in our analysis.) The solution to

$$\frac{d}{dr} \left(\frac{r^3 f V_F}{\sigma} \frac{d\mathcal{G}(r, \tilde{r})}{dr} \right) - \left(\frac{r V_F^2 G'^2}{\sigma} + 2q^2 r \sigma V_\psi \rho^2 \right) \mathcal{G}(r, \tilde{r}) = \delta(r - \tilde{r}) \quad (6.36)$$

which vanishes at the asymptotically AdS boundary and also at the horizon is given by

$$\mathcal{G}(r, \tilde{r}) = \begin{cases} -\frac{1}{2}a(\tilde{r})\hat{a}(r) & r > \tilde{r} \\ -\frac{1}{2}a(r)\hat{a}(\tilde{r}) & r < \tilde{r}. \end{cases} \quad (6.37)$$

Thus, if $S^{\text{total}}(\infty) = 0$, a near boundary series expansion of δg_μ is given by

$$\delta g_\mu = -\frac{1}{2r^2} \int_{r_h}^{\infty} a \mathbf{S}_\mu^{\text{total}} dx + \frac{1}{2r^2} \lim_{x \rightarrow \infty} \left(x^2 \frac{d}{dx} \mathbf{S}_\mu^{\text{total}}(x) \right) + \dots \quad (6.38)$$

Then, according to the standard AdS/CFT prescription, the dissipative corrections to the charged current in the boundary theory take the form

$$\bar{J}_{diss\mu} = -\frac{1}{2\kappa^2} \left(\int_{r_h}^{\infty} a \mathbf{S}_\mu^{\text{total}} dx - \lim_{x \rightarrow \infty} \left(x^2 \frac{d}{dx} \mathbf{S}_\mu^{\text{total}}(x) \right) \right). \quad (6.39)$$

Let us focus on parity odd contributions to $\mathbf{S}^{\text{total}}$. we find that

$$\mathbf{S}_\mu^{\text{total}} = 4c\kappa^2(G - \mu g)G'\omega_\mu - 2c\kappa^2 gG'B_\mu + \left(\text{parity even terms} \right) \quad (6.40)$$

where ω^μ and B^μ were defined in Table 1. We reproduce their expression here for convenience

$$\omega^\mu \equiv \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} u_\nu \partial_\rho u_\sigma \quad B^\mu \equiv \frac{1}{2} \epsilon^{\mu\nu\rho\sigma} u_\nu F_{\rho\sigma}. \quad (6.41)$$

Hence

$$\bar{J}_{diss}^\mu = \frac{c}{Q_1 + Q_2\mu} \left(\int_{r_h}^{\infty} (Q_1 g + Q_2 G) g G' dr B^\mu + \int_{r_h}^{\infty} 2(\mu g - G)(Q_1 g + Q_2 G) G' dr \omega^\mu \right) + \left(\text{parity even terms} \right). \quad (6.42)$$

Following the work of [26], the contribution of the parity odd terms to the entropy current come from the horizon value of δV_μ , i.e., if we denote the entropy current by J_S^μ then

$$J_S^\mu = s\delta V^\mu(r_h) + \left(\text{parity even terms} \right). \quad (6.43)$$

It remains to evaluate the parity odd contributions to δV^μ at the horizon. Integrating (6.31b) once we find that

$$\frac{V_F f r^3 (\delta g_\mu G' - \delta g'_\mu G)}{\sigma} - 2\kappa^2 Q_1 \delta V_\mu = \int_{r_h}^r \mathbf{S}_\mu^{\delta g} + \delta Q \quad (6.44)$$

with δQ an integration constant. Evaluating (6.44) at the horizon and at the boundary, we find that

$$\begin{aligned}\delta V^\mu(r_h) &= -\frac{\mu \bar{J}_{diss}^\mu}{Q_1} + \frac{\int_{r_h}^\infty \mathbf{S}^{\delta g} dr}{2Q_1 \kappa^2} + \left(\begin{array}{c} \text{parity even} \\ \text{terms} \end{array} \right) \\ &= -\frac{\mu}{T} \bar{J}_{diss}^\mu + \frac{c}{T} \left(\int_{r_h}^\infty 2(\mu g - G) G G' dr \omega^\mu + \int_{r_h}^\infty g G G' dr B^\mu \right) + \left(\begin{array}{c} \text{parity even} \\ \text{terms} \end{array} \right).\end{aligned}\quad (6.45)$$

Restricting ourselves to parity odd contributions, the first term on the right hand side of the first line of (6.45) represents the contribution of the canonical part of the entropy current $J_{S\text{ canon}}^\mu$. The second term on the right hand side of the first line of (6.45) will give us the corrections to $J_{S\text{ canon}}^\mu$.

Following the notation in (4.21), we denote

$$R = \frac{q}{\rho + P} \quad (6.46)$$

and write the parity odd transport coefficients and the corrections to the entropy current as in (5.25)

$$\begin{aligned}\bar{J}_{diss}^\mu &= -\tilde{\kappa}_\omega \omega^\mu - \tilde{\kappa}_B B^\mu + \left(\begin{array}{c} \text{parity even} \\ \text{terms} \end{array} \right) \\ J_S^\mu &= -\frac{\mu}{T} \bar{J}_{diss}^\mu + \sigma_B \omega^\mu + \sigma_\omega \omega^\mu + \left(\begin{array}{c} \text{parity even} \\ \text{terms} \end{array} \right).\end{aligned}\quad (6.47)$$

Then, (6.45) and (6.42) imply

$$\tilde{\kappa}_B = -c \int_{r_h}^\infty g^2 G' + R(G - g\mu) g G' dr \quad (6.48a)$$

$$\tilde{\kappa}_\omega = 2c \int_{r_h}^\infty (G - \mu g) g G' + R(G - \mu g)^2 G' dr \quad (6.48b)$$

$$\sigma_\omega = \frac{2c}{T} \int_{r_h}^\infty (G - \mu g) G G' dr \quad (6.48c)$$

$$\sigma_B = \frac{c}{T} \int_{r_h}^\infty g G G' dr. \quad (6.48d)$$

While we can not solve the integrals in (6.48) explicitly, we find that the following relations are satisfied:

$$\frac{1}{2} \sigma_\omega - \mu \sigma_B = -\frac{c\mu^3}{3T} \quad (6.49a)$$

$$\frac{1}{2} \tilde{\kappa}_\omega - \mu \tilde{\kappa}_B + T(1 - \mu R) \sigma_B = -\frac{cR\mu^3}{3}. \quad (6.49b)$$

(To obtain (6.49) from (6.48) we used $G(\infty) = \mu$ and $G(r_h) = 0$.) The first of these relations is precisely the universal prediction (5.27) between entropy current coefficients predicted in the last section. Using (5.28) and (5.27) we can also obtain holographic expressions for s_9 , σ_1 , σ_8 and σ_{10}

$$s_9 = 0 \tag{6.50a}$$

$$\sigma_1 = \frac{c}{T^2} \left(\mu^2 - \int_{r_h}^{\infty} g(4G - \mu g)G' dr \right) \tag{6.50b}$$

$$\sigma_8 = \frac{c}{2T} \left(-\mu + \int_{r_h}^{\infty} g^2 G' dr \right) \tag{6.50c}$$

$$\sigma_{10} = -\frac{c}{T^2} \int g(G - \mu g)G' dr. \tag{6.50d}$$

To study the values of $\tilde{\kappa}_\omega$, $\tilde{\kappa}_B$, σ_ω and σ_B at the phase transition we set $g = 1$. Then, all the integrals in (6.48) can be carried out explicitly. We find that the resulting expressions exactly match the zero superfluid density values of $\tilde{\kappa}_\omega$, $\tilde{\kappa}_B$, σ_ω and σ_B studied in [21, 22, 20], implying that these coefficients are continuous across the phase transition.

7. Discussion

In this paper we have described a framework for describing superfluid hydrodynamics at first order in the derivative expansion. We have determined the most general form of the hydrodynamical equations that are consistent with Lorentz invariance, the Onsager principle, and the requirement that the second law of thermodynamics apply in every conceivable situation. We begin this section by summarizing our results, and then turn to a discussion of possible applications and extensions.

We have found that the most general equations of Lorentz invariant but non-parity conserving superfluids requires the specification of twenty parameters. Fourteen of these are associated with the parity even sector and were described already in [10], generalizing an earlier 13 parameter framework spelt out in [18, 19]). In this work we have found that six more parameters need to be specified in the parity odd sector of the theory. Five of these six parameters are non dissipative; they drop off from the expression of the divergence of the entropy current, and are unconstrained by inequalities. Three of these five parameters ($\tilde{\eta}$, $\tilde{\kappa}_{11}$ and $\tilde{\kappa}_{22}$ in the language of the main text) are relatively simple. They are the transport coefficients which are associated with the pseudo-tensor mode

$$\tilde{\mathcal{T}}^{\mu\alpha} = \epsilon^{\mu\nu\rho\sigma} u_\nu \zeta_\nu \sigma_\sigma^{\mu\alpha} + \epsilon^{\alpha\nu\rho\sigma} u_\nu \zeta_\nu \sigma_\sigma^{\mu\alpha} \tag{7.1a}$$

and the pseudo vector modes

$$\tilde{\mathcal{V}}_1 = \epsilon^{\mu\nu\rho\sigma} u_\nu \zeta_\rho \left(\partial_\sigma \frac{\mu}{T} - \frac{E_\sigma}{T} \right) \quad (7.1b)$$

$$\tilde{\mathcal{V}}_2 = \epsilon^{\mu\nu\rho\sigma} u_\nu \zeta_\rho \zeta^\alpha \sigma_{\alpha\sigma}. \quad (7.1c)$$

Note that $\tilde{\eta}$ closely resembles the hall viscosity in 2+1 dimensional theories [53]. There, when parity is violated it is possible to have a transport coefficient η_H associated with the tensor mode

$$\epsilon^{\mu\nu\rho} u_\nu \sigma_\rho^\alpha + \epsilon^{\alpha\nu\rho} u_\nu \sigma_\rho^\mu. \quad (7.2)$$

The hall viscosity is known to be associated with the ground state's intrinsic angular momentum. It would be interesting to see whether $\tilde{\eta}$ is also associated with similar quantities.

The remaining two nondissipative constitutive parameters, σ_8 and σ_{10} , multiply relatively complicated expressions in the constitutive relations. These two parameters determine the eight transport coefficients associated with the pseudo scalars $\omega \cdot \xi$ and $B \cdot \xi$ which appear in the constitutive relations, and the transport coefficients of the pseudo vectors $*\partial_\mu T$, $*\partial_\mu \frac{\mu}{T}$, $*\partial_\mu \frac{\zeta^2}{T^2}$, ω^μ and B^μ . It is possible that these expressions admit significant simplifications when expressed in terms of variables that might turn out to be more natural than those adopted in this paper. σ_8 and σ_{10} also parameterize deviations of the entropy current away from its canonical form.

The sixth parity odd parameter is dissipative since it enters into the formula for entropy production. In the main text we have denoted this parameter by $\tilde{\kappa}_{21}$ (and also $\tilde{\kappa}_{12} = -\tilde{\kappa}_{21}$). It is associated with the pseudo vector modes (7.1b) and (7.1c). While undetermined, it, and the transport coefficients κ_{ij} $i, j = 1, 2$ associated with the parity even sector, need to satisfy the inequality

$$\kappa_{11}\kappa_{22} \geq \kappa_{12}^2 + \zeta^2 \tilde{\kappa}_{12}^2.$$

Note that this equation mixes parity odd and parity even coefficients.

The triangle anomaly, if it exists, enters the expression for the pseudo scalar and pseudo vector coefficients whose value depends on σ_8 and σ_{10} . It is interesting to compare this result with that of [20] which studied (non-superfluid) hydrodynamics in the presence of triangle anomalies. There, the only addition to the parity-odd sector are the two pseudo vectors ω^μ and B^μ whose transport coefficients are completely fixed in terms of the anomaly and a single integration constants. When considering superfluids the functional dependence of the transport coefficients of ω^μ and B^μ on T , μ and ζ is completely arbitrary. This novel feature opens the possibility for interesting physical

phenomenon in a parity violating theory, even in the absence of an anomaly which is an intrinsically relativistic effect.

The most immediate application of our work would be to the modelling of the long distance behaviors of real world superfluids (or superconductors) whose hydrodynamics violates parity. We do not yet know of any candidate experimental systems of this sort. We do note, however, that non centro symmetric superconductors live on parity violating lattices. It would be exciting to investigate whether parity violation in these (or analogous systems) could lead to experimentally observable nonzero values for any of the 6 parameters described above.

In this paper we have worked out the general theory of superfluidity in 4 dimensions. Several aspects of our analysis depend on the existence of an ϵ tensor with 4 indices. It seems likely that the theory of superfluidity in 5 and higher dimensions will differ in qualitative aspects from the theory we have worked out. It would be interesting to flesh this out.

We have already verified certain aspects of our general construction of the equations of superfluid hydrodynamics using explicit computations within the AdS/CFT framework. However all the computations reported in this paper work in the collinear limit of vanishing superfluid velocity. It would be instructive to demonstrate that all the numerous other relations, implied by our work, between transport coefficients that do not preserve $SO(3)$ invariance, are also borne out by AdS/CFT computations. In this context we point out that the generality of the expected results suggests that their derivation may also be carried out for a generic holographic superfluid (perhaps along the lines of Section 6) and should not involve the details of the corresponding dynamical systems.

An interesting feature of our AdS/CFT analysis is that in the collinear limit all the transport coefficients in the parity odd sector of the theory are continuous at the phase transition (i.e., they match their value in the uncondensed phase where $f = 0$). This is similar to the behavior of the shear viscosity η and diffusion coefficient κ of the parity even sector, but differs from the divergent behavior of the transport coefficient associated with the scalar $\partial_\mu(f\xi^\mu)$ [16]. It would be interesting to understand this result in terms of the theory of dynamical critical exponents [54].

It would also be interesting to work out the properties of stationary superfluid flows (like rotating superfluids on an S^3) and to investigate how the properties of such configurations are affected by the parity odd non dissipative terms that can be ‘turned on’ in such configurations. Such configurations would be dual, under the AdS/CFT correspondence, to rotating hairy black holes in global AdS space.

To end this paper let us highlight a structural aspect of our analysis that we find quite remarkable. For this purpose it is sufficient to focus on the case of parity invariant

superfluids. The total number of constitutive coefficients allowed, merely on symmetry grounds, is 50. In this paper we have shown that the requirement that the equations that follow from these constitutive relations are consistent with positivity of divergence of *any* entropy current (plus the Onsager relations) cuts down the number of constitutive coefficients from 50 to 14. In other words, the second law of thermodynamics gives very powerful and precise constraints on dynamical equations. Given the duality between fluid dynamics and gravity, it is natural to wonder whether similar results may also be true for extensions of the theory of gravity. We leave detailed investigation of this exciting possibility to future work.

Acknowledgments

We would like especially to thank D. Son for very useful discussions and correspondence. We would like to acknowledge useful discussions and correspondences with K. Damle, D. Dhar, R. Gopakumar, C. Herzog, R. Loganayagam, D. Son, V. Tripathy and S. Wadia. The work of S. M. was supported in part by a Swarnajayanti Fellowship. J. B., S. B., and S. M. would also like to acknowledge their debt to the steady and generous support of the people of India for research in basic science. S. M. Would like to thank the organizers of the Great Lakes Strings Meeting, Iranian school for applications of AdS/CFT and the Solvay conference for hospitality during the completion of this work. A. Y. is supported in part by the Department of Energy under Grant No. DE-FG02-91ER40671.

A. The linear independence of data for the parity even superfluid

A.1 The linear independence of first order terms

A.1.1 The vector sector

The equations of motion in the vector sector are

$$\begin{aligned}
 \tilde{P}_{\mu\beta} \partial_\nu T^{\nu\beta} &= \tilde{P}_{\mu\beta} F^{\beta\nu} J_\nu \\
 \tilde{P}^{\mu\beta} u^\nu (\partial_\beta \xi_\nu - \partial_\nu \xi_\beta) &= \tilde{P}^{\mu\beta} E_\beta \\
 \tilde{P}^{\mu\beta} \xi^\nu (\partial_\beta \xi_\nu - \partial_\nu \xi_\beta) &= \tilde{P}^{\mu\beta} F_{\beta\nu} \xi^\nu .
 \end{aligned}
 \tag{A.1}$$

A basis of ten one derivative vectors (before using the equations of motion) was listed in table 3. It is given by

$$\begin{aligned} & \tilde{P}^{\mu\beta}(u\cdot\partial)u_\beta, \tilde{P}^{\mu\beta}(u\cdot\partial)\xi_\beta, \tilde{P}^{\mu\beta}(\xi\cdot\partial)u_\beta, \tilde{P}^{\mu\beta}(\xi\cdot\partial)\xi_\beta, \tilde{P}^{\mu\beta}\partial_\beta\left(\frac{\mu}{T}\right), \\ & \tilde{P}^{\mu\beta}\partial_\beta T, \tilde{P}^{\mu\beta}\partial_\beta\left(\frac{\xi}{T}\right), \tilde{P}^{\mu\beta}\xi^\nu\partial_\beta u_\nu, \tilde{P}^{\mu\beta}E_\beta, \tilde{P}^{\mu\beta}F_{\beta\nu}\xi^\nu. \end{aligned} \quad (\text{A.2})$$

The quantities in (A.2) are not all on-shell inequivalent as they are constrained by the relations (A.1). In this subsection we will argue that it is consistent to choose the seven vectors listed in the third column of Table 3 as independent vector data. That is, we will show that it is possible to use the equations (A.1) to solve for $\tilde{P}^{\mu\beta}\partial_\beta T$, $\tilde{P}^{\mu\beta}\partial_\beta\left(\frac{\xi}{T}\right)$, $\tilde{P}^{\mu\beta}\xi^\nu\partial_\beta u_\nu$ in terms of

$$\begin{aligned} & \tilde{P}^{\mu\beta}(u\cdot\partial)u_\beta, \tilde{P}^{\mu\beta}(u\cdot\partial)\xi_\beta, \tilde{P}^{\mu\beta}(\xi\cdot\partial)u_\beta, \tilde{P}^{\mu\beta}(\xi\cdot\partial)\xi_\beta, \tilde{P}^{\mu\beta}\partial_\beta\left(\frac{\mu}{T}\right), \\ & \tilde{P}^{\mu\beta}E_\beta, \tilde{P}^{\mu\beta}F_{\beta\nu}\xi^\nu \end{aligned} \quad (\text{A.3})$$

If we rewrite the equations of motion in (A.1) in terms of the quantities in (A.2) we find

$$\begin{aligned} \tilde{P}^{\mu\beta}\left((P+\rho)(u\cdot\partial)u_\beta + \partial_T P \partial_\beta T + \partial_{\frac{\mu}{T}}\partial_\beta\frac{\mu}{T} + \partial_{\frac{\xi}{T}}\partial_\beta\frac{\xi}{T} + f(\xi\cdot\partial)\xi_\beta\right) &= \tilde{P}^{\mu\beta}(qE_\mu - fF_{\mu\nu}\xi^\nu) \\ \tilde{P}^{\mu\beta}\left((u\cdot\partial)\xi_\beta - T\partial_\beta\frac{\mu}{T} - \frac{\mu}{T}\partial_\beta T + \xi^\nu\partial_\beta u_\nu\right) &= \tilde{P}^{\mu\beta}E_\beta \\ \tilde{P}^{\mu\beta}\left((\xi\cdot\partial)\xi_\beta + T\xi\partial_\beta\frac{\xi}{T} + \frac{\xi}{T}\partial_\beta T\right) &= \tilde{P}^{\mu\beta}F_{\beta\nu}\xi^\nu. \end{aligned} \quad (\text{A.4})$$

It is possible to use (A.4) to solve for the scalars listed in (A.3) if and only if the 3×3 matrix of the three vectors $\tilde{P}^{\mu\beta}\partial_\beta T$, $\tilde{P}^{\mu\beta}\partial_\beta\left(\frac{\xi}{T}\right)$, $\tilde{P}^{\mu\beta}\xi^\nu\partial_\beta u_\nu$ in the three equations (A.4) has nonzero determinant. This 3×3 matrix is given by

$$M_{(v)} = \begin{pmatrix} \partial_T P & \partial_{\xi/T} P & 0 \\ -\mu/T & 0 & 1 \\ \xi^2/T & T\xi & 0 \end{pmatrix} \quad (\text{A.5})$$

and its determinant is given by

$$\text{Det}(M_{(v)}) = \xi \left(\frac{\xi}{T} \partial_{\frac{\xi}{T}} P - T \partial_T P \right). \quad (\text{A.6})$$

It is nonzero for a generic functional form for $P(T, \mu, \xi)$. We conclude that the vectors (A.3) form a basis for onshell independent one derivative vectors.

A.1.2 The scalar sector

The equations of motion in the scalar sector are given by

$$\begin{aligned}
\xi_\nu \partial_\mu T^{\mu\nu} &= qE \cdot \xi \\
u_\nu \partial_\mu T^{\mu\nu} &= fE \cdot \xi \\
\partial_\mu J^\mu &= cE \cdot B \\
u^\nu \xi^\mu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) &= E \cdot \xi.
\end{aligned} \tag{A.7}$$

A basis of 11 one derivative vectors (before using the equations of motion) was listed in Table 3. We denote them by $\{\mathcal{L}_j^{(a)}, S_i^{(a)}\}$ for the first set of on shell independent scalars and $\{\mathcal{L}_j^{(b)}, S_i^{(b)}\}$ for the second set. Here j runs from 1 to 4 and i runs from 1 to 7. We have used the notation in Table 4. The new quantities $\mathcal{L}_j^{(b)}, \mathcal{L}_j^{(b)}$ are defined as follows

$$\begin{aligned}
\mathcal{L}_1^{(a)} &= u \cdot \partial \Sigma_1, & \mathcal{L}_i^{(b)} &= \xi \cdot \partial \Sigma_1, \\
\mathcal{L}_2^{(a)} &= u \cdot \partial \Sigma_2, & \mathcal{L}_i^{(b)} &= \xi \cdot \partial \Sigma_2, \\
\mathcal{L}_3^{(a)} &= u \cdot \partial \Sigma_3, & \mathcal{L}_i^{(b)} &= \xi \cdot \partial \Sigma_3, \\
\mathcal{L}_4^{(a)} &= \xi^\mu u \cdot \partial u_\mu, & \mathcal{L}_i^{(b)} &= \xi^\mu \xi \cdot \partial u_\mu.
\end{aligned} \tag{A.8}$$

The quantities defined in (A.8) are the dependent data for the two choices of bases among the on-shell inequivalent quantities. These quantities are to be determined by the equation of motion (A.7) in terms of the dependent quantities S_i . Note that the sets $\{\mathcal{L}_i^{(a)}, S_i^{(a)}\}$ and $\{\mathcal{L}_i^{(b)}, S_i^{(b)}\}$ are different partitioning of the same set of quantities.

The equation of motion in (A.7) expressed in terms of the quantities in (A.8) has the form

$$\begin{aligned}
\sum_{i=1}^7 (\mathbf{e}_i^{(a)})_p S_i^{(a)} + \sum_{j=1}^4 (\ell_j^{(a)})_p \mathcal{L}_j^a &= 0. \\
\sum_{i=1}^7 (\mathbf{e}_i^{(b)})_p S_i^{(b)} + \sum_{j=1}^4 (\ell_j^{(b)})_p \mathcal{L}_j^b &= 0.
\end{aligned} \tag{A.9}$$

In the equations above the index p runs from 1 to 4 denoting the 4 equations in (A.7). Again note that both the equations in (A.9) refer to the same set of equations. We find it convenient to define the new set of quantities

$$A = -\frac{\chi^2}{T^2(\nu^2 - \chi^2)}; \quad B = -\frac{1}{T^2(\mu^2 - \xi^2)}; \quad C = -\frac{\nu}{T(\nu^2 - \xi^2)}. \tag{A.10}$$

so that the projector

$$\tilde{P}^{\mu\nu} = \eta^{\mu\nu} + Au^\mu u^\nu + B\xi^\mu \xi^\nu + C(\xi^\mu u^\nu + u^\mu \xi^\nu). \tag{A.11}$$

The coefficients in (A.9) are given by

$$\begin{aligned}
(\mathbf{e}_1^{(a)})_1 &= -(P + \rho), & (\mathbf{e}_2^{(a)})_1 &= f\nu, & (A.12) \\
(\mathbf{e}_3^{(a)})_1 &= B(P + \rho) + Cf\mu - f, & (\mathbf{e}_4^{(a)})_1 &= B\xi T f\mu + \mu\partial_\chi f, \\
(\mathbf{e}_5^{(a)})_1 &= \mu\partial_\nu f - CTf\mu + fT, & (\mathbf{e}_6^{(a)})_1 &= \mu\partial_T f + 2f\nu, \\
(\mathbf{e}_7^{(a)})_1 &= -f \\
(\mathbf{e}_1^{(a)})_2 &= (P + \rho)\mu, & (\mathbf{e}_2^{(a)})_2 &= -f\chi^2 T, \\
(\mathbf{e}_3^{(a)})_2 &= -(\mu B(\rho + P) + CTf\xi^2), & (\mathbf{e}_4^{(a)})_2 &= \partial_\chi P - \xi^2\partial_\chi f - \xi fT - B\xi^3 T f, \\
(\mathbf{e}_5^{(a)})_2 &= \partial_\nu P - \xi^2\partial_\nu f + CTf\xi^2, & (\mathbf{e}_6^{(a)})_2 &= \partial_T P - \xi^2\partial_T f - 2f\chi^2 T, \\
(\mathbf{e}_7^{(a)})_2 &= -q, \\
(\mathbf{e}_1^{(a)})_3 &= q, & (\mathbf{e}_2^{(a)})_3 &= \frac{f}{T} \\
(\mathbf{e}_3^{(a)})_3 &= -(Bq + Cf), & (\mathbf{e}_4^{(a)})_3 &= -\partial_\chi f - B\xi fT \\
(\mathbf{e}_5^{(a)})_3 &= \partial_\nu f + CTf, & (\mathbf{e}_6^{(a)})_3 &= -\partial_T f - \frac{f}{T} \\
(\mathbf{e}_7^{(a)})_3 &= 0, \\
(\mathbf{e}_1^{(a)})_4 &= 0, & (\mathbf{e}_2^{(a)})_4 &= 0, \\
(\mathbf{e}_3^{(a)})_4 &= -1, & (\mathbf{e}_4^{(a)})_4 &= 0 \\
(\mathbf{e}_5^{(a)})_4 &= T, & (\mathbf{e}_5^{(a)})_6 &= \nu, \\
(\mathbf{e}_7^{(a)})_4 &= -1.
\end{aligned}$$

and

$$\begin{aligned}
(\ell_1^{(a)})_1 &= Cf\mu\xi T - \partial_\chi \rho, & (\ell_2^{(a)})_1 &= -(Af\mu T + \partial_\nu \rho), & (A.13) \\
(\ell_3^{(a)})_1 &= -\partial_T \rho, & (\ell_4^{(a)})_1 &= Af\mu + C(P + \rho), \\
(\ell_1^{(a)})_2 &= \mu\partial_\chi(P + \rho) - Cf\xi^3 T, & (\ell_2^{(a)})_2 &= \mu\partial_\nu(P + \rho) + TA\xi^2 f, \\
(\ell_3^{(a)})_2 &= \mu\partial_T(P + \rho), & (\ell_1^{(a)})_2 &= (P + \rho) - C\mu(P + \rho) - Af\xi^2 \\
(\ell_1^{(a)})_3 &= \partial_\chi q - C\xi fT, & (\ell_2^{(a)})_3 &= \partial_\nu q + ATf, \\
(\ell_3^{(a)})_3 &= \partial_T q, & (\ell_4^{(a)})_3 &= -(Cq + Af) \\
(\ell_1^{(a)})_4 &= \xi T, & (\ell_2^{(a)})_4 &= 0, \\
(\ell_3^{(a)})_4 &= \chi^2 T, & (\ell_4^{(a)})_4 &= 0.
\end{aligned}$$

The other set of coefficients, with index (b) , can be read from (A.12) and (A.13) using

$$\begin{aligned} (\ell_1^{(b)})_i &= (\mathbf{e}_4^{(a)})_i; & (\ell_2^{(b)})_i &= (\mathbf{e}_5^{(a)})_i; & (\ell_3^{(b)})_i &= (\mathbf{e}_6^{(a)})_i; & (\ell_4^{(b)})_i &= (\mathbf{e}_3^{(a)})_i; \\ (\mathbf{e}_4^{(b)})_i &= (\ell_1^{(a)})_i; & (\mathbf{e}_5^{(b)})_i &= (\ell_2^{(a)})_i; & (\mathbf{e}_6^{(b)})_i &= (\ell_3^{(a)})_i; & (\mathbf{e}_3^{(b)})_i &= (\ell_4^{(a)})_i; \end{aligned} \quad (\text{A.14})$$

We can express all the derivatives in (A.12) and (A.13) as derivatives of a single function, say, the pressure. Thermodynamic relations that enable us to do so are

$$q = \frac{1}{T} \partial_{\mu/T} P; \quad f = \frac{1}{T\xi} \partial_{\xi/T} P; \quad \rho = -P + T \partial_T P - \frac{\xi}{T} \partial_{\frac{\xi}{T}} P. \quad (\text{A.15})$$

We make the following observations:

- a) We can use the equations of motion (A.9) to solve for the 4 scalars $\xi^\mu u \cdot \nabla u_\mu$, $u \cdot \partial \Sigma_i$ ($i = 1, \dots, 3$) in terms of the 7 independent scalars in the 3rd column of the first row of Table 3. This is possible if and only if the 4×4 matrix of coefficients of the four quantities in the first equation in (A.9) has nonzero determinant. This matrix is given by

$$M_{ij}^{(a)} = (\ell_i^{(a)})_j; \quad (\text{A.16})$$

- b) We can use the equations of motion (A.9) to solve for the quantities $\xi^\mu \xi \cdot \partial u_\mu$, $\xi \cdot \partial \Sigma_i$ in terms of the 7 independent scalars in the 3rd column of the second row of Table 3. This is possible if and only if the 4×4 matrix of coefficients of the 4 quantities in the second equation in (A.9) has nonzero determinant. This matrix is given

$$M_{ij}^{(b)} = (\ell_i^{(b)})_j; \quad (\text{A.17})$$

The relations (A.15) allow us to express the matrices (A.16) and (A.17) in terms of the pressure. Using several reasonable equations of state we have used `Mathematica` to verify that the determinant of the matrices in (A.16) and (A.17) is generically non-zero.

A.2 The linear independence of the second order terms

A list of second order scalar data, the second order equations of motion and a choice of second order independent scalar data can be found in Table 5. The second order scalar equations that follows from the first order vector equations are

$$\begin{aligned} \nabla_\alpha \left(\tilde{P}^{\alpha\mu} u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) \right) &= \nabla_\alpha \left(\tilde{P}^{\alpha\mu} E_\mu \right), \\ \nabla_\mu \left(\tilde{P}^{\mu\nu} \nabla_\beta T_\nu^\beta \right) &= \nabla_\mu \left(\tilde{P}^{\mu\nu} F_{\nu\beta} J^\beta \right) \\ \nabla_\alpha \left(\tilde{P}^{\alpha\mu} \xi^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu) \right) &= \nabla_\alpha \left(\tilde{P}^{\alpha\mu} F_{\mu\nu} \xi^\nu \right) \end{aligned} \quad (\text{A.18})$$

The two derivative terms in these equations take the form

$$\tilde{P}^{\mu\nu} \left((P + \rho) u^\beta \nabla_\mu \nabla_\beta u_\nu + \partial_T \rho \nabla_\mu \partial_\nu \frac{\mu}{T} + \partial_{\xi/T} \nabla_\mu \partial_\nu \frac{\xi}{T} + f \xi^\beta \nabla_\mu \nabla_\beta \xi_\nu \right) = \dots \quad (\text{A.19})$$

$$\tilde{P}^{\mu\nu} \left(u^\beta \nabla_\mu \nabla_\beta \xi_\nu - T \nabla_\mu \partial_\nu \frac{\mu}{T} - \frac{\mu}{T} \nabla_\mu \partial_\nu T + \xi^\beta \nabla_\mu \nabla_\nu u_\beta \right) = \dots \quad (\text{A.20})$$

$$\tilde{P}^{\mu\nu} \left(\xi^\beta \nabla_\mu \nabla_\beta \xi_\nu + T \xi \nabla_\mu \partial_\nu \frac{\xi}{T} + \frac{\xi^2}{T} \nabla_\mu \partial_\nu T \right) = \dots \quad (\text{A.21})$$

The quantities $\tilde{P}^{\mu\nu} \nabla_\mu \partial_\nu T$, $\tilde{P}^{\mu\nu} \nabla_\mu \partial_\nu \frac{\xi}{T}$, $\tilde{P}^{\mu\nu} \xi^\beta \nabla_\mu \nabla_\nu u_\beta$ can be solved for using equations (A.19), (A.20), and (A.21). Note that these two derivative scalar quantities do not appear in any other equations of motion. We can then use the remaining 8 equations of motion to solve for the other 8 dependent data,

$$u^\mu u^\nu \nabla_\mu \partial_\nu \Sigma_i, \quad u^\mu u^\nu \xi^\beta \nabla_\mu \nabla_\nu u_\beta, \quad \xi^\mu \xi^\nu \xi^\beta \nabla_\mu \nabla_\nu u_\beta, \quad \xi^\mu \xi^\nu \nabla_\mu \partial_\nu \Sigma_i \quad (\text{A.22})$$

where i runs from 1 to 3. The remaining 8 two derivative scalar equations of motion are

$$u^\beta \nabla_\beta (u_\mu \nabla_\nu T^{\mu\nu}) = u^\beta \nabla_\beta (-E_\mu J^\mu), \quad (\text{A.23a})$$

$$u^\beta \nabla_\beta (\xi_\mu \nabla_\nu T^{\mu\nu}) = u^\beta \nabla_\beta (\xi^\mu F_{\mu\nu} J^\nu), \quad (\text{A.23b})$$

$$u^\beta \nabla_\beta (\nabla_\mu J^\mu) = u^\beta \nabla_\beta (c E^\mu B_\mu), \quad (\text{A.23c})$$

$$u^\beta \nabla_\beta (\xi^\mu u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu)) = u^\beta \nabla_\beta (\xi^\mu E_\mu) \quad (\text{A.23d})$$

$$\xi^\beta \nabla_\beta (u_\mu \nabla_\nu T^{\mu\nu}) = \xi^\beta \nabla_\beta (-E_\mu J^\mu), \quad (\text{A.23e})$$

$$\xi^\beta \nabla_\beta (\xi_\mu \nabla_\nu T^{\mu\nu}) = \xi^\beta \nabla_\beta (\xi^\mu F_{\mu\nu} J^\nu), \quad (\text{A.23f})$$

$$\xi^\beta \nabla_\beta (\nabla_\mu J^\mu) = \xi^\beta \nabla_\beta (c E^\mu B_\mu), \quad (\text{A.23g})$$

$$\xi^\beta \nabla_\beta (\xi^\mu u^\nu (\partial_\mu \xi_\nu - \partial_\nu \xi_\mu)) = \xi^\beta \nabla_\beta (\xi^\mu E_\mu) \quad (\text{A.23h})$$

The matrix of coefficients of the terms in (A.22) as they appear in the equation of motion (A.23) may be expressed as

$$N_{ij} = \begin{pmatrix} M_{ij}^{(a)} & 0 \\ 0 & M_{ij}^{(b)} \end{pmatrix} \quad (\text{A.24})$$

where the rows represent the ordered equations in (A.23) and the columns represent the ordered quantities in (A.22). It follows that

$$\text{Det}[N_{ij}] = \text{Det}[M_{ij}^{(a)}] \text{Det}[M_{ij}^{(b)}]. \quad (\text{A.25})$$

In the previous section we concluded that both $\text{Det}[M_{ij}^{(b)}]$ and $\text{Det}[M_{ij}^{(a)}]$ are generically non-zero. Therefore we can infer that $\text{Det}[N_{ij}]$ is also generically non-zero.

In order to understand the structure of the matrix N we note that the first four equations in (A.23) are generated by the action of $u \cdot \partial$ on the first equation in (A.9). We then find that $u \cdot \partial$ acting on $S_i^{(a)}$ generates all the independent second order data as presented in Table 5. Likewise, the action of $u \cdot \partial$ on the $\mathcal{L}_i^{(a)}$ generates the four terms $u^\mu u^\nu \xi^\beta \nabla_\mu \nabla_\nu u_\beta$, $u^\mu u^\nu \nabla_\mu \partial_\nu \Sigma_i$ ($i = 1, \dots, 3$). In fact these dependent two derivative terms appear only in equations (A.23a), (A.23c), (A.23d), (A.23e) and is not there in the rest of the four equations in (A.23).

Similarly, we can think of the equations (A.23f), (A.23g), (A.23h), (A.23h) as being obtained by the action of $\xi \cdot \partial$ on the second equation in (A.9). Also here the terms $\xi^\mu \xi^\nu \xi^\beta \nabla_\mu \nabla_\nu u_\beta$, $\xi^\mu \xi^\nu \nabla_\mu \partial_\nu \Sigma_i$ (which constitutes the 4 remaining second order quantities which are determined by the equation of motion) are generated by $\xi \cdot \partial$ acting on the $\mathcal{L}_i^{(b)}$ terms. These dependent four second order quantities do not appear in the first four equations in (A.23). This structure justifies the block diagonal form of the coefficient matrix in (A.24).

B. Derivation of thermodynamic identities for normal fluids

In this appendix we present a derivation of the equations of motion (2.22). Our starting point is the observation that the scalar components of the equations of conservation of the stress tensor and current reduce, at first order to

$$\begin{aligned} u_\nu \nabla_\mu T^{\mu\nu} &= u_\nu F^{\mu\nu} J_\nu = 0 \\ \Rightarrow (u\partial)\rho + (\rho + P)\Theta &= 0 \end{aligned} \tag{B.1}$$

$$\nabla_\mu J^\mu = 0 \Rightarrow (u \cdot \partial)q + q\Theta = 0 \tag{B.2}$$

where $\Theta = (\nabla \cdot u)$

Derivation of the first equation in (2.22)

Subtracting $[\frac{\partial q}{\partial \nu} \times \text{eq. (B.2)}]$ from $[\frac{\partial \rho}{\partial \nu} \times \text{eq. (B.1)}]$ we find the following relations:

$$\begin{aligned}
& \frac{\partial q}{\partial \nu}(u \cdot \partial)\rho - \frac{\partial \rho}{\partial \nu}(u \cdot \partial)q = - \left[(\rho + P)\frac{\partial q}{\partial \nu} - q\frac{\partial \rho}{\partial \nu} \right] \Theta \\
\Rightarrow & \left[\frac{\partial q}{\partial \nu} \frac{\partial \rho}{\partial T} - \frac{\partial \rho}{\partial \nu} \frac{\partial q}{\partial T} \right] (u \cdot \partial)T = - \left[(\rho + P)\frac{\partial q}{\partial \nu} - q\frac{\partial \rho}{\partial \nu} \right] \Theta \\
\Rightarrow & \left[\frac{\partial q}{\partial \nu} \frac{\partial P}{\partial T} - \frac{\partial P}{\partial \nu} \frac{\partial q}{\partial T} \right] (u \cdot \partial)T = - \left[\frac{\partial P}{\partial \rho} \right]_q \left[(\rho + P)\frac{\partial q}{\partial \nu} - q\frac{\partial \rho}{\partial \nu} \right] \Theta \\
\Rightarrow & \left[\left(\frac{\rho + P}{T} \right) \frac{\partial q}{\partial \nu} - Tq \frac{\partial q}{\partial T} \right] (u \cdot \partial)T = - \left[\frac{\partial P}{\partial \rho} \right]_q \left[(\rho + P)\frac{\partial q}{\partial \nu} - q\frac{\partial \rho}{\partial \nu} \right] \Theta \\
\Rightarrow & \frac{(u \cdot \partial)T}{T} = - \left[\frac{\partial P}{\partial \rho} \right]_q \Theta
\end{aligned} \tag{B.3}$$

In the third line we have used two identities:

$$\begin{aligned}
\left[\frac{\partial P}{\partial \rho} \right]_q \left[\frac{\partial \rho}{\partial T} \right]_\nu &= \left[\frac{\partial P}{\partial T} \right]_\nu - \left[\frac{\partial P}{\partial q} \right]_\rho \left[\frac{\partial q}{\partial T} \right]_\nu \\
\left[\frac{\partial P}{\partial \rho} \right]_q \left[\frac{\partial \rho}{\partial \nu} \right]_\nu &= \left[\frac{\partial P}{\partial \nu} \right]_\nu - \left[\frac{\partial P}{\partial q} \right]_\rho \left[\frac{\partial q}{\partial \nu} \right]_\nu
\end{aligned} \tag{B.4}$$

In the last line of (B.3) we have used the following three thermodynamic identities.

$$\begin{aligned}
\left[\frac{\partial P}{\partial T} \right]_\nu &= \frac{\rho + P}{T} \\
\left[\frac{\partial P}{\partial \nu} \right]_T &= Tq \\
\left[\frac{\partial \rho}{\partial \nu} \right]_T &= T \frac{\partial^2 P}{\partial T \partial \nu} - \frac{\partial P}{\partial \nu} = T^2 \frac{\partial}{\partial T} \left[\frac{1}{T} \frac{\partial P}{\partial \nu} \right] = T^2 \left[\frac{\partial q}{\partial T} \right]_\nu
\end{aligned} \tag{B.5}$$

Derivation of the second equation in (2.22)

Adding $[\frac{\partial P}{\partial q} \times \text{eq. (B.2)}]$ and $[\frac{\partial P}{\partial \rho} \times \text{eq. (B.1)}]$ we find

$$\begin{aligned}
& \frac{\partial P}{\partial \rho}(u \cdot \partial)\rho + \frac{\partial P}{\partial q}(u \cdot \partial)q = - \left[(\rho + P)\frac{\partial P}{\partial \rho} + q\frac{\partial P}{\partial q} \right] \Theta \\
\Rightarrow & (u \cdot \partial)P = - \left[(\rho + P)\frac{\partial P}{\partial \rho} + q\frac{\partial P}{\partial q} \right] \Theta \\
\Rightarrow & \left[(\rho + P)\frac{(u \cdot \partial)T}{T} + Tq(u \cdot \partial)\nu \right] = - \left[(\rho + P)\frac{\partial P}{\partial \rho} + q\frac{\partial P}{\partial q} \right] \Theta \\
\Rightarrow & (u \cdot \partial)\nu = - \frac{1}{T} \left[\frac{\partial P}{\partial q} \right]_\rho \Theta
\end{aligned} \tag{B.6}$$

In the third line of (B.6) we have used the first law

$$\begin{aligned} dP &= (S + \nu q) dT + T q d\nu = (\rho + P) \frac{dT}{T} + T q d\nu \\ \rho + P &= T(S + \nu q) \end{aligned} \tag{B.7}$$

and in the last line of (B.6) we have used the first equation in (2.22).

Derivation of the third equation in (2.22)

The third equation in (2.22) follows from the vector component of the stress tensor conservation equation.

$$\begin{aligned} P_{\mu\theta} \nabla_\nu T^{\nu\theta} &= P_{\mu\theta} F^{\theta\alpha} J_\alpha = q E_\mu \\ \Rightarrow P_\mu^\theta \partial_\theta P + (\rho + P)(u \cdot \nabla) u_\mu &= q E_\mu \\ \Rightarrow P_\mu^\theta \frac{\partial_\theta T}{T} + (u \cdot \nabla) u_\mu &= \frac{q}{\rho + P} V_{1\mu} \end{aligned} \tag{B.8}$$

In the last line of (B.8) we have used the first law as written in (B.7).

C. Details relating to parity violating superfluids

In this Appendix we provide several computational details relating to section 4.

C.1 All equations of motion for ideal superfluid

The equations of motion for a superfluid are listed in table 3. There are four scalar equations, one pseudo scalar equation and 3 vector equations. In this subsection, using thermodynamics, we simplify these equations so that they can be easily used to solve for the dependent fluid data in terms of the independent data.

First we note the identity

$$\begin{aligned} \nabla_\mu \xi_\nu - \nabla_\nu \xi_\mu &= F_{\mu\nu} \\ \Rightarrow -\xi \nabla_\mu \xi - (\xi \cdot \nabla) \xi_\mu &= F_{\mu\nu} \xi^\nu. \end{aligned} \tag{C.1}$$

In the second line we have contracted both sides of the equation with ξ_ν . Using (C.1) and the first law we can simplify the stress tensor conservation equation projected in

the direction perpendicular to u^μ :

$$\begin{aligned}
& P_\theta^\mu \nabla_\nu T^{\nu\theta} - P^{\mu\theta} F_{\theta\nu} J^\nu = 0 \\
\Rightarrow & P_\theta^\mu \nabla_\nu \left[(\rho + P) u^\nu u^\theta + P \eta^{\nu\theta} + f \xi^\nu \xi^\theta \right] - P_\theta^\mu F^{\theta\nu} (q u_\nu - f \xi_\nu) = 0 \\
\Rightarrow & P^{\mu\theta} [\nabla_\theta P + f(\xi \cdot \nabla) \xi_\theta + f F_{\theta\nu} \xi^\nu] + (\rho + P)(u \cdot \nabla) u^\mu + \zeta^\mu \nabla_\theta (f \xi^\theta) = q E^\mu \\
\Rightarrow & P^{\mu\theta} [\nabla_\theta P - f \xi \nabla_\theta \xi] + (\rho + P)(u \cdot \nabla) u^\mu + \zeta^\mu \nabla_\theta (f \xi^\theta) = q E^\mu \tag{C.2} \\
\Rightarrow & P^{\mu\theta} \left[(\rho + P) \frac{\nabla_\theta T}{T} + T q \nabla_\theta \nu \right] + (\rho + P)(u \cdot \nabla) u^\mu + \zeta^\mu \nabla_\mu (f \xi^\mu) = q E^\mu \\
\Rightarrow & P^{\sigma\mu} \left(\frac{\nabla_\sigma T}{T} \right) + (u \cdot \nabla) u^\mu = \left[\frac{q T}{\rho + P} \right] \left(\frac{E^\mu}{T} - P^{\mu\theta} \nabla_\theta \nu \right) - \zeta^\mu \left[\frac{\nabla_\theta (f \xi^\theta)}{\rho + P} \right]
\end{aligned}$$

In the second step of (C.2) we have used the identity (C.1) and in the third step of (C.2) we used the first law in the form:

$$dP = (\rho + P) \frac{dT}{T} + T q d\nu + f \xi d\xi.$$

Next, using (C.2) we can simplify the equation relating the curl of the phase velocity ξ^μ to the field strength (ie. $C_{\mu\nu} \equiv \nabla_\mu \xi_\nu - \nabla_\nu \xi_\mu - F_{\mu\nu} = 0$). Consider the case where one of the two free indices of this equation is projected in the direction of u^μ .

$$\begin{aligned}
& u^\beta (\nabla_\mu \xi_\beta - \nabla_\beta \xi_\mu - F_{\mu\beta}) = 0 \\
\Rightarrow & u^\beta [\nabla_\mu (-\mu u_\beta + \zeta_\beta) - \nabla_\beta (-\mu u_\mu + \zeta_\mu)] = E_\mu \\
\Rightarrow & [\nabla_\mu \mu + u^\mu (u \cdot \nabla) \mu] + \mu (u \cdot \nabla) u_\mu - (u \cdot \nabla) \zeta_\mu - \zeta^\theta \nabla_\mu u_\theta = E_\mu \\
\Rightarrow & [P_\mu^\theta \nabla_\theta \mu + \mu (u \cdot \nabla) u_\mu] - (u \cdot \nabla) \zeta_\mu - \zeta^\theta \nabla_\mu u_\theta = E_\mu \tag{C.3} \\
\Rightarrow & P_\mu^\theta \nabla_\theta \nu + \nu [P_\mu^\theta \nabla_\theta T + T (u \cdot \nabla) u_\mu] - (u \cdot \nabla) \zeta_\mu - \zeta^\theta \nabla_\mu u_\theta = E_\mu \\
\Rightarrow & (u \cdot \nabla) \zeta_\mu = -T \left(1 - \frac{\mu q}{\rho + P} \right) \left(\frac{E_\mu}{T} - P_\mu^\theta \nabla_\theta \nu \right) - \zeta^\theta \nabla_\mu u_\theta
\end{aligned}$$

In the last step we have used (C.2).

Next consider the case where both the indices of $C_{\mu\nu}$ are projected in the direction perpendicular to u^μ . This can be simply analyzed by contracting the two free indices of $C_{\mu\nu}$ with $\epsilon^{\mu\nu\lambda\sigma} u_\nu$

$$\begin{aligned}
& \epsilon^{\mu\nu\lambda\sigma} u_\nu (\nabla_\lambda \xi_\sigma - \nabla_\sigma \xi_\lambda - F_{\lambda\sigma}) = 0 \\
\Rightarrow & 2\epsilon^{\mu\nu\lambda\sigma} u_\nu \nabla_\lambda \xi_\sigma - \epsilon^{\mu\nu\lambda\sigma} u_\nu F_{\lambda\sigma} = 0 \\
\Rightarrow & -2\mu \epsilon^{\mu\nu\lambda\sigma} u_\nu \nabla_\lambda u_\sigma + 2\epsilon^{\mu\nu\lambda\sigma} u_\nu \nabla_\lambda \zeta_\sigma - \epsilon^{\mu\nu\lambda\sigma} u_\nu F_{\lambda\sigma} = 0 \tag{C.4} \\
\Rightarrow & \Omega^\mu = \frac{B^\mu}{2} + \mu \omega^\mu.
\end{aligned}$$

where Ω^μ , ω^μ and B^μ are defined as

$$\Omega^\mu = \frac{1}{2}\epsilon^{\mu\nu\lambda\sigma}u_\nu\nabla_\lambda\zeta_\sigma, \quad \omega^\mu = \frac{1}{2}\epsilon^{\mu\nu\lambda\sigma}u_\nu\nabla_\lambda u_\sigma, \quad B^\mu = \frac{1}{2}\epsilon^{\mu\nu\lambda\sigma}u_\nu F_{\lambda\sigma}$$

If we project (C.2), (C.3) and (C.4) in the direction perpendicular to ζ^μ , it will be a rewriting of the three vector equations as listed in Table 3. Contracting the free index with ζ^μ we get two of the scalar equations and one pseudo scalar equation from Table 3, rewritten in our basis.²¹

Next we rewrite the two remaining scalar equations of table 3. First, using these two remaining scalar equations, we will show that the entropy current is conserved in equilibrium. It is most useful to use the basis in Table 6

$$\begin{aligned} u_\nu\nabla_\mu T^{\mu\nu} - u_\nu F^{\nu\mu}J_\mu &= 0 \\ \Rightarrow u_\nu\nabla_\mu \left[(\rho + P)u^\mu u^\nu + P\eta^{\mu\nu} + f\xi^\mu\xi^\nu \right] + fu_\nu F^{\nu\mu}\xi_\mu &= 0 \\ \Rightarrow -(u\cdot\nabla)\rho - (\rho + P)(\nabla\cdot u) + \mu\nabla_\theta(f\xi^\theta) + fu_\nu [(\xi\cdot\nabla)\xi^\nu + F^{\nu\mu}\xi_\mu] &= 0 \\ \Rightarrow -[(u\cdot\nabla)\rho + f\xi(u\cdot\nabla)\xi] - (\rho + P)(\nabla\cdot u) + \mu\nabla_\theta(f\xi^\theta) &= 0 \\ \Rightarrow -[T(u\cdot\nabla)s + \mu(u\cdot\nabla)q] - (Ts + \mu q)(\nabla\cdot u) + \mu\nabla_\theta(f\xi^\theta) &= 0 \\ \Rightarrow -T\nabla_\mu(su^\mu) - \mu\nabla_\mu(qu^\mu - f\xi^\mu) &= 0 \\ \Rightarrow \nabla_\mu(su^\mu) &= 0. \end{aligned} \tag{C.5}$$

In the fourth line we have used the identity (C.1). In fifth line we have used

$$d\rho = T ds + \mu dq - f\xi d\xi$$

and

$$\rho + P = Ts + \mu q$$

In the last line we have used the fact that

$$\nabla_\mu J^\mu = \nabla_\mu(qu^\mu - f\xi^\mu) = \mathcal{O}(\text{Two derivatives}) \sim 0$$

Using $\nabla_\mu(su^\mu) = 0$ and $\nabla_\mu(qu^\mu - f\xi^\mu) = 0$ we find

$$\nabla_\mu[f\xi^\mu] = s(u\cdot\nabla) \left[\frac{q}{s} \right]. \tag{C.6}$$

²¹The relevant equations are

1. $\zeta_\mu\nabla_\nu T^{\mu\nu} - \zeta_\mu F^{\mu\nu}J_\nu = 0$
2. $u^\mu\zeta^\nu C_{\mu\nu} = 0$
3. $\epsilon^{\mu\nu\lambda\sigma}\zeta_\mu u_\nu C_{\lambda\sigma} = 0$

To summarize, we list all the equations in simplified form. The four scalar and one pseudo scalar equation are given by

$$\begin{aligned}
(\nabla \cdot u) \equiv \Theta &= -\frac{(u \cdot \partial)s}{s} = [B_1(u \cdot \partial)\chi + B_2(u \cdot \partial)\nu + B_3(u \cdot \partial)T] \\
\frac{\nabla_\theta(f\xi^\theta)}{(\rho + P)} \equiv K &= \frac{s(u \cdot \partial)\left(\frac{q}{s}\right)}{(\rho + P)} = [K_1(u \cdot \partial)\chi + K_2(u \cdot \partial)\nu + K_3(u \cdot \partial)T] \\
\frac{(\zeta \cdot \nabla)T}{T} + \zeta^\theta(u \cdot \nabla)u_\theta &= RT(V_1 \cdot \zeta) - \zeta^2 K \\
\zeta^\mu \zeta^\nu \sigma_{\mu\nu} &= T^2 [A_1(u \cdot \partial)\chi + A_2(u \cdot \partial)\nu + A_3(u \cdot \partial)T] - T(1 - \mu R)(V_1 \cdot \zeta) \\
\Omega \cdot \zeta &= \frac{B \cdot \zeta}{2} + \mu(\omega \cdot \zeta)
\end{aligned} \tag{C.7}$$

The three vector equations are given by

$$\begin{aligned}
\tilde{P}^{\sigma\mu} \left[\frac{\nabla_\sigma T}{T} + (u \cdot \nabla)u_\sigma \right] &= RT \tilde{P}_\sigma^\mu V_1^\sigma \\
\tilde{P}^{\sigma\mu} \Omega_\sigma &= \tilde{P}^{\sigma\mu} \left[\frac{B_\sigma}{2} + \mu\omega_\sigma \right] \\
\tilde{P}^{\sigma\mu} (u \cdot \nabla)\zeta_\sigma &= \tilde{P}_\sigma^\mu [-T(1 - \mu R)V_1^\sigma - \zeta^\nu \nabla_\sigma u_\nu]
\end{aligned} \tag{C.8}$$

where we have defined

$$\begin{aligned}
\nu &= \frac{\mu}{T}, \quad \chi = \frac{\zeta^2}{T^2}, \quad K = \frac{\nabla_\theta[f\xi^\theta]}{\rho + P}, \quad R = \frac{q}{\rho + P} \\
B_1 &= -\frac{\partial}{\partial\chi}[\log(s)], \quad B_2 = -\frac{\partial}{\partial\nu}[\log(s)], \quad B_3 = -\frac{\partial}{\partial T}[\log(s)] \\
K_1 &= \frac{s}{\rho + P} \frac{\partial}{\partial\chi} \left[\frac{q}{s} \right], \quad K_2 = \frac{s}{\rho + P} \frac{\partial}{\partial\nu} \left[\frac{q}{s} \right], \quad K_3 = \frac{s}{\rho + P} \frac{\partial}{\partial T} \left[\frac{q}{s} \right] \\
A_1 &= -\frac{1}{2} - \nu\chi(1 - \mu R) \left[\frac{\partial}{\partial\chi} \left(\frac{q}{s} \right) \right] + \frac{\chi}{3s} \frac{\partial s}{\partial\chi} \\
A_2 &= -\nu\chi(1 - \mu R) \left[\frac{\partial}{\partial\nu} \left(\frac{q}{s} \right) \right] + \frac{\chi}{3s} \frac{\partial s}{\partial\nu} \\
A_3 &= -\nu\chi(1 - \mu R) \left[\frac{\partial}{\partial T} \left(\frac{q}{s} \right) \right] + \frac{\chi}{3s} \left(\frac{\partial s}{\partial T} - \frac{3s}{T} \right) \\
V_\mu &= \frac{E_\mu}{T} - P_\mu^\sigma \nabla_\sigma \left[\frac{\mu}{T} \right] \\
\Omega^\mu &= \frac{1}{2} \epsilon^{\mu\nu\lambda\sigma} u_\nu \nabla_\lambda \zeta_\sigma, \quad \omega^\mu = \frac{1}{2} \epsilon^{\mu\nu\lambda\sigma} u_\nu \nabla_\lambda u_\sigma, \quad B^\mu = \frac{1}{2} \epsilon^{\mu\nu\lambda\sigma} u_\nu F_{\lambda\sigma}.
\end{aligned} \tag{C.9}$$

C.2 Showing the linear independence, of the first derivative scalar data

Using the scalar equations we can solve for four dependent scalars and one dependent pseudo scalar in terms of the independent ones as they appear in the list of \mathcal{S}_i^c in table 6. We choose the five dependent scalars to be:

$$\Theta, \quad K, \quad \zeta \cdot \sigma \cdot \zeta, \quad \zeta^\nu (u \cdot \nabla) u_\nu, \quad \Omega \cdot \zeta. \quad (\text{C.10})$$

Note that the last term is the pseudo scalar. The dependence of the scalars in (C.10) on the \mathcal{S}_i^c 's from table 6 takes the following form

$$\begin{aligned} \Theta &= B_3 \mathcal{S}_2^c + B_2 \mathcal{S}_3^c + B_1 \mathcal{S}_4^c \\ K &= K_3 \mathcal{S}_2^c + K_2 \mathcal{S}_3^c + K_1 \mathcal{S}_4^c \\ \zeta \cdot \sigma \cdot \zeta &= A_3 \mathcal{S}_2^c + A_2 \mathcal{S}_3^c + A_1 \mathcal{S}_4^c - T(1 - \mu R) \mathcal{S}_1^c \\ \zeta^\nu (u \cdot \nabla) u_\nu &= RT \mathcal{S}_1^c + \zeta^2 [K_3 \mathcal{S}_2^c + K_2 \mathcal{S}_3^c + K_1 \mathcal{S}_4^c] - \frac{1}{T} \mathcal{S}_5^c \\ \Omega \cdot \zeta &= \mu(\omega \cdot \zeta) + \frac{(B \cdot \zeta)}{2} \end{aligned} \quad (\text{C.11})$$

where B_i , K_i and A_i are defined in equation (C.9). Using the first three of the above equations one can form the A^s matrix in (4.20).

C.3 Showing the linear independence, of the first derivative vector data

We wish to argue that the vectors \mathcal{V}^c form a set of independent vectors. To do so we solve for the three vectors that do not appear in the list of $\mathcal{V}_i^{c\mu}$ in terms of $\mathcal{V}_i^{c\mu}$ using the three vector equations of motion. The three vectors that are not part of the set $\{\mathcal{V}_i^c\}$ are

$$\tilde{P}^{\theta\mu}(u \cdot \nabla) u_\theta, \quad \tilde{P}^{\theta\mu}(u \cdot \nabla) \zeta_\theta, \quad \tilde{P}^{\theta\mu}(\zeta \cdot \nabla) \zeta_\theta \quad (\text{C.12})$$

From the first vector equation (C.2) we can solve for $\tilde{P}^{\theta\mu}(u \cdot \nabla) u_\theta$,

$$\begin{aligned} \tilde{P}^{\theta\mu}(u \cdot \nabla) u_\theta &= \tilde{P}^{\theta\mu} \left(-\frac{\nabla_\theta T}{T} + RT V_{1\theta} \right) \\ &= RT \mathcal{V}_1^{c\mu} - \frac{\mathcal{V}_3^{c\mu}}{T}. \end{aligned} \quad (\text{C.13})$$

This equation has been used to form the matrix A^v in (4.20).

Taking the star of the second vector equation (C.4) we can solve for $\tilde{P}^{\theta\mu}(\zeta \cdot \nabla) \zeta_\theta$

$$\begin{aligned} \tilde{P}^{\theta\mu}(\zeta \cdot \nabla) \zeta_\theta &= \tilde{P}^{\theta\mu} \left[\mu \zeta^\alpha (\partial_\alpha u_\theta - \partial_\theta u_\alpha) + \frac{T^2}{2} \partial_\theta \chi + T \chi \partial_\theta T - F_{\theta\lambda} \zeta^\lambda \right] \\ &= T \chi \mathcal{V}_3^{c\mu} + \frac{T^2}{2} \mathcal{V}_5^{c\mu} + \mu \zeta^2 \mathcal{V}_6^{c\mu} + \zeta^2 \mathcal{V}_7^{c\mu}. \end{aligned} \quad (\text{C.14})$$

From the last vector equation (C.3) we can solve for $\tilde{P}^{\theta\mu}(u \cdot \nabla)\zeta_\theta$

$$\begin{aligned}\tilde{P}^{\theta\mu}(u \cdot \nabla)\zeta_\theta &= \tilde{P}^{\theta\mu} \left[-T(1 - \mu R)V_{1\theta} - \zeta^\alpha \sigma_{\alpha\theta} - \frac{\zeta^\alpha}{2}(\partial_\theta u_\alpha - \partial_\alpha u_\theta) \right] \\ &= -T(1 - \mu R)\mathcal{V}_1^{c\mu} - \mathcal{V}_2^{c\mu} - \zeta^2 \mathcal{V}_6^{c\mu}.\end{aligned}\tag{C.15}$$

C.4 Showing the linear independence of the two derivative parity odd scalar data

Here we solve for the five two derivative parity odd scalars using the five parity odd two derivative equations of motion as listed in Table 7. The solution will be presented in terms of the three independent two derivative pseudo scalars as listed in the last column of Table 7 and squares of single derivative terms. The two derivative pseudo scalars that we shall solve for, are the following

$$\begin{array}{lll}e^{\lambda\sigma}(u \cdot \nabla)\nabla_\lambda u_\sigma & \zeta^\mu u^\nu \nabla_\nu B_\mu & \tilde{P}^{\mu\nu}\nabla_\mu B_\nu \\ e^{\lambda\sigma}(u \cdot \nabla)\nabla_\lambda \zeta_\sigma & e^{\lambda\sigma}(\zeta \cdot \nabla)\nabla_\lambda \zeta_\sigma & \end{array}$$

In the solution below we shall write the dependence on the two derivative terms explicitly but will not write the terms which are squares of a single derivative.

$$\begin{aligned}E_1 &\equiv [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] \nabla_\lambda (\nabla_\theta T_\sigma^\theta - F_{\sigma\theta} j^\theta) = 0 \\ &\Rightarrow \nabla_\lambda \left[(\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu) (\nabla_\theta T_\sigma^\theta - F_{\sigma\theta} j^\theta) \right] = \text{Products of one derivatives} \\ &\Rightarrow \nabla_\lambda \left(\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu \left[(u \cdot \nabla) u_\sigma + \frac{\partial_\sigma T}{T} - R(E_\sigma - T \partial_\sigma \nu) \right] \right) = \text{Products of one derivatives} \\ &\Rightarrow [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] (u^\theta \nabla_\theta \nabla_\lambda u_\sigma - R \nabla_\lambda E_\sigma) = \text{Products of one derivatives}\end{aligned}\tag{C.16}$$

Using E_1 we can solve $e^{\lambda\sigma}(u \cdot \nabla)\nabla_\lambda u_\sigma$ in terms of $e^{\lambda\sigma}\nabla_\lambda E_\sigma$.

$$\begin{aligned}E_2 &\equiv [\epsilon^{\mu\nu\lambda\sigma} \zeta_\mu] \nabla_\nu F_{\lambda\sigma} = 0 \\ &\Rightarrow [\epsilon^{\mu\nu\lambda\sigma} \zeta_\mu u_\nu] (u \cdot \nabla) F_{\lambda\sigma} + 2 [\epsilon^{\mu\nu\lambda\sigma} \zeta_\mu u_\lambda] u^\theta \nabla_\nu F_{\theta\sigma} = 0 \\ &\Rightarrow \zeta_\mu (u \cdot \nabla) [\epsilon^{\mu\nu\lambda\sigma} u_\nu F_{\lambda\sigma}] + 2 [\epsilon^{\mu\nu\lambda\sigma} \zeta_\mu u_\lambda] \nabla_\nu (u^\theta F_{\theta\sigma}) = \text{Products of one derivatives} \\ &\Rightarrow [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu u_\lambda] \nabla_\lambda E_\sigma - \zeta^\mu u^\nu \nabla_\nu B_\mu = \text{Products of one derivatives}\end{aligned}\tag{C.17}$$

Using E_2 we can solve $\zeta^\mu u^\nu \nabla_\nu B_\mu$ in terms of $e^{\lambda\sigma} \nabla_\lambda E_\sigma$.

$$\begin{aligned}
E_3 &\equiv [\epsilon^{\mu\nu\lambda\sigma} u_\mu] \nabla_\nu F_{\lambda\sigma} = 0 \\
&\Rightarrow \nabla_\mu [\epsilon^{\mu\nu\lambda\sigma} u_\nu F_{\lambda\sigma}] = \text{Products of one derivatives} \\
&\Rightarrow \nabla_\mu B^\mu = \text{Products of one derivatives} \\
&\Rightarrow \tilde{P}^{\mu\nu} \nabla_\mu B_\nu + \frac{\zeta^\mu \zeta^\nu}{\zeta^2} \nabla_\mu B_\nu = \text{Product of one derivatives}
\end{aligned} \tag{C.18}$$

Using E_3 we can solve $\tilde{P}^{\mu\nu} \nabla_\mu B_\nu$ in terms of $\frac{\zeta^\mu \zeta^\nu}{\zeta^2} \nabla_\mu B_\nu$.

$$\begin{aligned}
E_4 &\equiv [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] u^\theta \nabla_\lambda C_{\theta\sigma} = 0 \\
&\Rightarrow \nabla_\lambda \left([\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] [(u \cdot \nabla) \zeta_\sigma + T(1 - \mu R) V_\sigma + \zeta^\theta \nabla_\sigma u_\theta] \right) \\
&\quad = \text{Product of one derivatives} \\
&[\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] \left[(u \cdot \nabla) \nabla_\lambda \zeta_\sigma + (1 - \mu R) \nabla_\lambda E_\sigma + \frac{\zeta^\theta}{2} [\nabla_\lambda, \nabla_\sigma] u_\theta \right] \\
&\quad = \text{Product of one derivatives}
\end{aligned} \tag{C.19}$$

Using E_4 we can solve $e^{\lambda\sigma} (u \cdot \nabla) \nabla_\lambda \zeta_\sigma$ in terms of $e^{\lambda\sigma} \nabla_\lambda E_\sigma$.

$$\begin{aligned}
E_5 &\equiv [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] (\zeta \cdot \nabla) C_{\lambda\sigma} = 0 \\
&\Rightarrow [\epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu] \left[(\zeta \cdot \nabla) \nabla_\lambda \zeta_\sigma - \mu (\zeta \cdot \nabla) \nabla_\lambda u_\sigma \right] + \frac{1}{2} \zeta^\mu \zeta^\nu \nabla_\mu B_\nu \\
&\quad = \text{Product of one derivatives}
\end{aligned} \tag{C.20}$$

Using E_5 we can solve $e^{\lambda\sigma} (\zeta \cdot \nabla) \nabla_\lambda \zeta_\sigma$ in terms of $e^{\lambda\sigma} (\zeta \cdot \nabla) \nabla_\lambda u_\sigma$ and $\zeta^\mu \zeta^\nu \nabla_\mu B_\nu$. Thus, we can choose $e^{\lambda\sigma} \nabla_\lambda E_\sigma$, $e^{\lambda\sigma} (\zeta \cdot \nabla) \nabla_\lambda u_\sigma$ and $\zeta^\mu \zeta^\nu \nabla_\mu B_\nu$ as the three independent two derivative parity odd scalar data.²²

C.5 Relating the two different entropy current

The relation between the variables in (4.3) and (4.9) can be obtained by using

$$\begin{aligned}
\frac{1}{2} \epsilon^{\mu\nu\lambda\sigma} \xi_\nu F_{\lambda\sigma} &= -\mu B^\mu + (B \cdot \zeta) u^\mu - T \tilde{\mathcal{V}}_1^{c\mu} - T \tilde{\mathcal{V}}_4^{c\mu} \\
\frac{1}{2} \epsilon^{\mu\nu\lambda\sigma} \nabla_\nu [T u_\lambda \zeta_\sigma] &= T \left[u^\mu (\omega \cdot \zeta) + \frac{RT}{2} \tilde{\mathcal{V}}_1^{c\mu} \right] - T \left(\mu \omega^\mu + \frac{B^\mu}{2} \right)
\end{aligned} \tag{C.21}$$

²²Here

$$C_{\mu\nu} \equiv \nabla_\mu \xi_\nu - \nabla_\nu \xi_\mu - F_{\mu\nu}$$

and

$$e^{\lambda\sigma} = \epsilon^{\mu\nu\lambda\sigma} u_\mu \zeta_\nu$$

Explicitly, we find

$$\tilde{s}_1^a = T\sigma_1 \quad (\text{C.22})$$

$$\tilde{s}_2^a = T\sigma_8 \quad (\text{C.23})$$

$$\tilde{s}_1^b = \alpha_3 - \frac{T\mu}{\zeta^2}\sigma_1 + \frac{T^2}{\zeta^2}\sigma_9 \quad (\text{C.24})$$

$$\tilde{s}_2^b = \alpha_4 - \frac{T}{2\zeta^2}\sigma_1 - \frac{\mu}{\zeta^2}\sigma_8 + \frac{T}{\zeta^2}\sigma_{10} \quad (\text{C.25})$$

$$\tilde{v}_1 = \alpha_1 + \sigma_1 \frac{RT^2}{2} - T\sigma_8 \quad (\text{C.26})$$

$$\tilde{v}_2 = \alpha_2 \quad (\text{C.27})$$

$$\tilde{v}_3 = T\sigma_3 + T\partial_T\sigma_1 \quad (\text{C.28})$$

$$\tilde{v}_4 = T\sigma_4 + T\partial_\nu\sigma_1 - T\sigma_8 \quad (\text{C.29})$$

$$\tilde{v}_5 = T\sigma_5 + T\partial_\chi\sigma_1 \quad (\text{C.30})$$

$$\tilde{v}_6 = -T\mu\sigma_1 + T^2\sigma_9 \quad (\text{C.31})$$

$$\tilde{v}_7 = -\frac{T}{2}\sigma_1 - \mu\sigma_8 + T\sigma_{10} \quad (\text{C.32})$$

C.6 Computation of the divergence of entropy current

$\nabla_\mu J_{S_{\text{new}}}^\mu$ can be calculated term by term.

$$\nabla_\mu (\epsilon^{\mu\nu\lambda\sigma} \nabla_\nu [T\sigma_1 u_\lambda \zeta_\sigma]) = 0 \quad (\text{C.33})$$

$$\begin{aligned} \nabla_\mu [T^2\sigma_9\omega^\mu] &= -2T^2\sigma_9(\omega\cdot\zeta)K + 2RT^3\sigma_9(\omega\cdot V) \\ &\quad + T^2 \left[\frac{\partial\sigma_9}{\partial T}(\omega\cdot\nabla T) + \frac{\partial\sigma_9}{\partial\nu}(\omega\cdot\nabla\nu) + \frac{\partial\sigma_9}{\partial\chi}(\omega\cdot\nabla\chi) \right] \end{aligned} \quad (\text{C.34})$$

$$\begin{aligned} \nabla_\mu [T\sigma_{10}B^\mu] &= -T\sigma_{10}(B\cdot\zeta)K + RT^3\sigma_{10}(B\cdot V) - 2T^2\sigma_{10}(\omega\cdot V) \\ &\quad - 2T^2\sigma_{10}(\omega\cdot\nabla\nu) + T \left[\frac{\partial\sigma_{10}}{\partial T}(B\cdot\nabla T) + \frac{\partial\sigma_{10}}{\partial\nu}(B\cdot\nabla\nu) + \frac{\partial\sigma_{10}}{\partial\chi}(B\cdot\nabla\chi) \right] \end{aligned} \quad (\text{C.35})$$

$$\begin{aligned} \nabla_\mu (\sigma_3 \mathcal{V}_3^{c\mu}) &= -2\sigma_3(\omega\cdot\zeta)(u\cdot\nabla)T - \frac{\partial\sigma_3}{\partial\nu}(\mathcal{V}_4^{c\mu}\cdot\nabla T) - \frac{\partial\sigma_3}{\partial\chi}(\mathcal{V}_5^{c\mu}\cdot\nabla T) \\ &\quad + \sigma_3 [RT(V\cdot\mathcal{V}_3^{c\mu}) + 2\mu(\omega\cdot\nabla T) + (B\cdot\nabla T)] \end{aligned} \quad (\text{C.36})$$

$$\begin{aligned}\nabla_\mu(T\sigma_4\mathcal{V}_4^{c\mu}) &= -2T\sigma_4(\omega\cdot\zeta)(u\cdot\nabla)\nu + T\left[\frac{\partial\sigma_4}{\partial T}(\mathcal{V}_4^{c\mu}\cdot\nabla T) + \frac{\partial\sigma_4}{\partial\chi}(\mathcal{V}_4^{c\mu}\cdot\nabla\chi)\right] \\ &\quad + T\sigma_4[RT(V\cdot\mathcal{V}_4^{c\mu}) + 2\mu(\omega\cdot\nabla\nu) + (B\cdot\nabla\nu)]\end{aligned}\quad (\text{C.37})$$

$$\begin{aligned}\nabla_\mu(T\sigma_5\mathcal{V}_5^{c\mu}) &= -2T\sigma_5(\omega\cdot\zeta)(u\cdot\nabla)\chi + T\left[\frac{\partial\sigma_5}{\partial T}(\mathcal{V}_5^{c\mu}\cdot\nabla T) - \frac{\partial\sigma_5}{\partial\chi}(\mathcal{V}_4^{c\mu}\cdot\nabla\chi)\right] \\ &\quad + T\sigma_5[RT(V\cdot\mathcal{V}_5^{c\mu}) + 2\mu(\omega\cdot\nabla\chi) + (B\cdot\nabla\chi)]\end{aligned}\quad (\text{C.38})$$

$$\begin{aligned}\nabla_\mu\left(\frac{\sigma_8}{2}\epsilon^{\mu\nu\lambda\sigma}\xi_\nu F_{\lambda\sigma}\right) &= (B\cdot\zeta)(u\cdot\nabla)\sigma_8 - \mu(B\cdot\nabla)\sigma_8 - 2T\sigma_8[(B\cdot V) + B\cdot\nabla\nu] \\ &\quad + T\left[\frac{\partial\sigma_8}{\partial T}(V\cdot\mathcal{V}_3^{c\mu}) + \frac{\partial\sigma_8}{\partial\nu}(V\cdot\mathcal{V}_4^{c\mu}) + \frac{\partial\sigma_8}{\partial\chi}(V\cdot\mathcal{V}_5^{c\mu})\right] \\ &\quad - T\left[\frac{\partial\sigma_8}{\partial T}(\mathcal{V}_4^{c\mu}\cdot\nabla T) + \frac{\partial\sigma_8}{\partial\chi}(\mathcal{V}_4^{c\mu}\cdot\nabla\chi)\right]\end{aligned}\quad (\text{C.39})$$

In deriving these expressions we have used the following identities.

$$\begin{aligned}\frac{1}{2}\epsilon^{\mu\nu\lambda\sigma}\xi_\nu F_{\lambda\sigma} &= -\mu B^\mu + (B\cdot\zeta)u^\mu - \mathcal{V}_1^{c\mu} - \mathcal{V}_4^{c\mu} \\ \frac{1}{2}\epsilon^{\mu\nu\lambda\sigma}\nabla_\nu[Tu_\lambda\zeta_\sigma] &= T\left[u^\mu(\omega\cdot\zeta) + \frac{RT}{2}\mathcal{V}_1^{c\mu}\right] - T\left(\mu\omega^\mu + \frac{B^\mu}{2}\right)\end{aligned}\quad (\text{C.40})$$

C.7 Frame transformation formula

Under the frame transformation

$$\begin{aligned}u'_\mu &= u_\mu - \delta u_\mu \\ T' &= T - \delta T \\ \nu' &= \nu - \delta\nu\end{aligned}\quad (\text{C.41})$$

the stress tensor, current and μ_{diss} transform in the following way.

$$\begin{aligned}T'_{diss}{}^{\mu\nu} - T_{diss}{}^{\mu\nu} &= \delta T_{diss}{}^{\mu\nu} = \delta(\rho + P)u^\mu u^\nu + \delta P\eta^{\mu\nu} + \delta f\xi^\mu\xi^\nu + (\rho + P)(u^\mu\delta u^\nu + u^\nu\delta u^\mu) \\ J'_{diss}{}^\mu - J_{diss}{}^\mu &= \delta J_{diss}{}^\mu = \delta q u^\mu - \delta f\xi^\mu + q\delta u^\mu \\ \mu'_{diss} - \mu_{diss} &= \delta\mu_{diss} = \delta\mu - \xi\cdot\delta u\end{aligned}\quad (\text{C.42})$$

where δA for any scalar function A denotes

$$\delta A = \frac{\partial A}{\partial T} \delta T + \frac{\partial A}{\partial \nu} \delta \nu$$

Using these expressions one can deduce how \mathbf{s}_i transform under a change of frame

$$\begin{aligned} \delta \mathbf{s}_1 &= 2\delta P \\ \delta \mathbf{s}_2 &= \delta P + \zeta^2 \delta f \\ \delta \mathbf{s}_3 &= \delta \rho + \mu^2 \delta f \\ \delta \mathbf{s}_4 &= \mu \zeta^2 \delta f - (\rho + P)(\delta u \cdot \zeta) \\ \delta \mathbf{s}_5 &= -\delta q - \mu \delta f \\ \delta \mathbf{s}_6 &= -\zeta^2 \delta f + q(\delta u \cdot \zeta) \\ \delta \mathbf{s}_7 &= -\delta \mu + (\delta u \cdot \zeta) \end{aligned} \tag{C.43}$$

and also how the \mathbf{v}_i transform under a change of frame.

$$\begin{aligned} \delta \mathbf{v}_1^\mu &= -(\rho + P)(\tilde{P}^{\mu\nu} \delta u_\nu) \\ \delta \mathbf{v}_2^\mu &= 0 \\ \delta \mathbf{v}_3^\mu &= q(\tilde{P}^{\mu\nu} \delta u_\nu) \end{aligned} \tag{C.44}$$

where ‘ δ ’ denotes a change under field redefinition.

Using these formula one can easily form the frame invariant combinations in the vector sector. It turns out that \mathbf{v}_2^μ is frame invariant by itself. The other frame invariant combination is proportional to

$$\left(\frac{q}{\rho + P} \right) \mathbf{v}_1^\mu + \mathbf{v}_3^\mu = R \mathbf{v}_1^\mu + \mathbf{v}_3^\mu$$

These are exactly the combinations that appear in the left hand side of equation (4.26).

Checking the frame invariance of the combinations appearing in the scalar sector is more involved and require many thermodynamic identities. Here we have used Mathematica (version 7) to impose all these identities by expressing ρ , q and f in terms of a single function $P(T, \nu, \xi)$ and its derivatives. Then we explicitly checked that the four combinations appearing in the left hand side of equation (4.35) are invariant under the transformations given by (C.43).

References

- [1] L. D. Landau, *Theory of the Superfluidity of Helium II*, *Phys. Rev.* **60** (1941) 356.

- [2] L. Tisza, *The Theory of Liquid Helium*, *Phys. Rev.* **72** (1947) 838–854.
- [3] W. Israel, *Covariant Superfluid Mechanics*, *Phys. Lett.* **86A** (1981) 79.
- [4] I. M. Khalatnikov and V. V. Lebedev, *Second sound in liquid helium II*, *Phys. Lett.* **91A** (1982) 70.
- [5] W. Israel, *Equivalence of Two Theories of Relativistic Superfluid Mechanics*, *Phys. Lett.* **92A** (1982) 77.
- [6] I. K. B. Carter, *Equivalence of convective and potential variational derivations of covariant superfluid dynamics*, *Phys. Rev.* **D45** (1992) 4536.
- [7] B. Carter and I. M. Khalatnikov, *Momentum, vorticity, and helicity in covariant superfluid dynamics*, *Ann. Phys. (NY)* **219** (1992) 219.
- [8] D. T. Son, *Hydrodynamics of relativistic systems with broken continuous symmetries*, *Int. J. Mod. Phys.* **A16S1C** (2001) 1284–1286, [[hep-ph/0011246](#)].
- [9] J. Sonner and B. Withers, *A gravity derivation of the Tisza-Landau Model in AdS/CFT*, *Phys. Rev.* **D82** (2010) 026001, [[arXiv:1004.2707](#)].
- [10] J. Bhattacharya, S. Bhattacharyya, and S. Minwalla, *Dissipative superfluid dynamics from gravity*, *JHEP* **04** (2011) 125, [[arXiv:1101.3332](#)].
- [11] C. Pujol and D. Davesne, *Relativistic dissipative hydrodynamics with spontaneous symmetry breaking*, *Phys. Rev.* **C67** (2003) 014901, [[hep-ph/0204355](#)].
- [12] M. E. Gusakov and N. Andersson, *Temperature dependent pulsations of superfluid neutron stars*, *Mon. Not. Roy. Astron. Soc.* **372** (2006) 1776–1790, [[astro-ph/0602282](#)].
- [13] M. A. Valle, *Hydrodynamic fluctuations in relativistic superfluids*, *Phys. Rev.* **D77** (2008) 025004, [[arXiv:0707.2665](#)].
- [14] M. Mannarelli and C. Manuel, *Bulk viscosities of a cold relativistic superfluid: color-flavor locked quark matter*, *Phys. Rev.* **D81** (2010) 043002, [[arXiv:0909.4486](#)].
- [15] M. E. Gusakov, *Bulk viscosity of superfluid neutron stars*, *Phys. Rev.* **D76** (2007) 083001, [[arXiv:0704.1071](#)].
- [16] C. P. Herzog, N. Lisker, P. Surowka, and A. Yarom, *Transport in holographic superfluids*, [arXiv:1101.3330](#).
- [17] E. M. Lifshitz and L. D. Landau, *Course of Theoretical Physics, volume 6*, .
- [18] J. A. Clark, *Ph.D. thesis, MIT, Cambridge, MA, unpublished* (1963).

- [19] S. Putterman, *Superfluid hydrodynamics*, North-Holland, Amsterdam (1974).
- [20] D. T. Son and P. Surowka, *Hydrodynamics with Triangle Anomalies*, *Phys. Rev. Lett.* **103** (2009) 191601, [[arXiv:0906.5044](#)].
- [21] J. Erdmenger, M. Haack, M. Kaminski, and A. Yarom, *Fluid dynamics of R-charged black holes*, *JHEP* **01** (2009) 055, [[arXiv:0809.2488](#)].
- [22] N. Banerjee *et. al.*, *Hydrodynamics from charged black branes*, [arXiv:0809.2596](#).
- [23] S. Bhattacharyya, S. Lahiri, R. Loganayagam, and S. Minwalla, *Large rotating AdS black holes from fluid mechanics*, *JHEP* **09** (2008) 054, [[arXiv:0708.1770](#)].
- [24] S. Bhattacharyya, V. E. Hubeny, S. Minwalla, and M. Rangamani, *Nonlinear Fluid Dynamics from Gravity*, *JHEP* **02** (2008) 045, [[arXiv:0712.2456](#)].
- [25] M. Van Raamsdonk, *Black Hole Dynamics From Atmospheric Science*, *JHEP* **05** (2008) 106, [[arXiv:0802.3224](#)].
- [26] S. Bhattacharyya *et. al.*, *Local Fluid Dynamical Entropy from Gravity*, *JHEP* **06** (2008) 055, [[arXiv:0803.2526](#)].
- [27] S. Bhattacharyya *et. al.*, *Forced Fluid Dynamics from Gravity*, *JHEP* **02** (2009) 018, [[arXiv:0806.0006](#)].
- [28] M. Haack and A. Yarom, *Nonlinear viscous hydrodynamics in various dimensions using AdS/CFT*, *JHEP* **10** (2008) 063, [[arXiv:0806.4602](#)].
- [29] S. Bhattacharyya, R. Loganayagam, I. Mandal, S. Minwalla, and A. Sharma, *Conformal Nonlinear Fluid Dynamics from Gravity in Arbitrary Dimensions*, *JHEP* **12** (2008) 116, [[arXiv:0809.4272](#)].
- [30] C. P. Herzog, *An Analytic Holographic Superconductor*, *Phys. Rev.* **D81** (2010) 126009, [[arXiv:1003.3278](#)].
- [31] S. S. Gubser, C. P. Herzog, S. S. Pufu, and T. Tesileanu, *Superconductors from Superstrings*, *Phys.Rev.Lett.* **103** (2009) 141601, [[arXiv:0907.3510](#)].
- [32] S. Bhattacharyya, S. Minwalla, and K. Papadodimas, *Small Hairy Black Holes in $AdS_5 \times S^5$* , [arXiv:1005.1287](#).
- [33] S. Lin, *On the anomalous superfluid hydrodynamics*, [arXiv:1104.5245](#).
- [34] C. P. Herzog, P. K. Kovtun, and D. T. Son, *Holographic model of superfluidity*, *Phys. Rev.* **D79** (2009) 066002, [[arXiv:0809.4870](#)].

- [35] S. Gubser, I. R. Klebanov, and A. M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys.Lett.* **B428** (1998) 105–114, [[hep-th/9802109](#)].
- [36] E. Witten, *Anti-de Sitter space and holography*, *Adv.Theor.Math.Phys.* **2** (1998) 253–291, [[hep-th/9802150](#)].
- [37] J. M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Adv.Theor.Math.Phys.* **2** (1998) 231–252, [[hep-th/9711200](#)].
- [38] S. S. Gubser, *Breaking an Abelian gauge symmetry near a black hole horizon*, *Phys. Rev.* **D78** (2008) 065034, [[arXiv:0801.2977](#)].
- [39] S. A. Hartnoll, C. P. Herzog, and G. T. Horowitz, *Building a Holographic Superconductor*, *Phys. Rev. Lett.* **101** (2008) 031601, [[arXiv:0803.3295](#)].
- [40] P. Basu, A. Mukherjee, and H.-H. Shieh, *Supercurrent: Vector Hair for an AdS Black Hole*, *Phys. Rev.* **D79** (2009) 045010, [[arXiv:0809.4494](#)].
- [41] I. Amado, M. Kaminski, and K. Landsteiner, *Hydrodynamics of Holographic Superconductors*, *JHEP* **05** (2009) 021, [[arXiv:0903.2209](#)].
- [42] C. P. Herzog and S. S. Pufu, *The Second Sound of $SU(2)$* , *JHEP* **04** (2009) 126, [[arXiv:0902.0409](#)].
- [43] A. Yarom, *Fourth sound of holographic superfluids*, *JHEP* **07** (2009) 070, [[arXiv:0903.1353](#)].
- [44] C. P. Herzog and A. Yarom, *Sound modes in holographic superfluids*, *Phys. Rev.* **D80** (2009) 106002, [[arXiv:0906.4810](#)].
- [45] S. S. Gubser and A. Yarom, *Pointlike probes of superstring-theoretic superfluids*, *JHEP* **1003** (2010) 041, [[arXiv:0908.1392](#)].
- [46] V. Keranen, E. Keski-Vakkuri, S. Nowling, and K. Yogendran, *Inhomogeneous Structures in Holographic Superfluids: I. Dark Solitons*, *Phys.Rev.* **D81** (2010) 126011, [[arXiv:0911.1866](#)].
- [47] V. Keranen, E. Keski-Vakkuri, S. Nowling, and K. Yogendran, *Inhomogeneous Structures in Holographic Superfluids: II. Vortices*, *Phys.Rev.* **D81** (2010) 126012, [[arXiv:0912.4280](#)].
- [48] V. Keranen, E. Keski-Vakkuri, S. Nowling, and K. Yogendran, *Solitons as Probes of the Structure of Holographic Superfluids*, [arXiv:1012.0190](#).
- [49] I. Amado, K. Landsteiner, and F. Pena-Benitez, *Anomalous transport coefficients from Kubo formulas in Holography*, [arXiv:1102.4577](#). * Temporary entry *.

- [50] S. S. Gubser and A. Nellore, *Ground states of holographic superconductors*, *Phys. Rev. D* **80** (2009) 105007, [[arXiv:0908.1972](#)].
- [51] A. Buchel and S. Cremonini, *Viscosity Bound and Causality in Superfluid Plasma*, *JHEP* **1010** (2010) 026, [[arXiv:1007.2963](#)].
- [52] M. Natsuume and M. Ohta, *The Shear viscosity of holographic superfluids*, [arXiv:1008.4142](#).
- [53] J. E. Avron, R. Seiler, and P. G. Zograf, *Viscosity of quantum hall fluids*, *Phys. Rev. Lett.* **75** (Jul, 1995) 697–700.
- [54] P. C. Hohenberg and B. I. Halperin, *Theory of dynamic critical phenomena*, *Rev. Mod. Phys.* **49** (Jul, 1977) 435–479.