

2469-9

**Workshop and Conference on Geometrical Aspects of Quantum States in  
Condensed Matter**

*1 - 5 July 2013*

**On the effective hydrodynamics of Quantum Hall fluids**

Alexander Abanov

*State University of New York at Stony Brook*

# On the effective hydrodynamics of Quantum Hall fluids

Alexander Abanov

July 1, 2013

Reference: *J. Phys. A: Math. Theor.* 46 (2013) 292001

Acknowledgments: Paul Wiegmann

# Main Result

Collective behavior of the simplest FQHE fluid (Laughlin's state with  $\nu = 1/\beta$ ) is captured by an effective theory of two scalar fields  $\rho$  and  $\pi$

$$L = -\rho \partial_t \pi - H$$


$$H = \rho \left[ A_0 + \frac{\tilde{\mathbf{v}}^2}{2} + \frac{1}{2} (\nabla \times \mathbf{A}) \right]$$

with

$$\tilde{\mathbf{v}} = \nabla \pi + \mathbf{A} - \nabla^* \left( \phi + \left( \frac{\beta}{4} - \frac{1}{2} \right) \ln \rho \right)$$

$$\rho = \frac{1}{2\pi\beta} \Delta \phi \tag{1}$$

# Fractional Quantum Hall Effect (FQHE)



# FQHE

- Disorder is necessary to explain the plateaus (as in IQHE)
- Two-dimensional electron gas in magnetic field forms a *new type of quantum fluid*
- Quantum condensation of *electrons coupled to vortices* (“flux quanta”)
- Incompressible fluid
- Quasiparticles are *gapped*, have *fractional charge and statistics*
- Low energy dynamics is at the *boundary*
- Microscopics is captured by Laughlin’s wave function

$$\Psi_{1/3} = \prod_{j < k} (z_j - z_k)^3 e^{-\frac{1}{4} \sum_j |z_j|^2}, \quad z \equiv x + iy$$

# Hydrodynamics of FQHE

Goal: effective classical hydrodynamic Hamiltonian of FQHE.

- Ideal 2D fluid (no dissipation)
- Hamiltonian formulation (nonlinearity)
- Density-vorticity relation (FQHE constraint)
- Incompressibility
- Hall viscosity
- Linear response (Hall conductivity, etc)
- Chirality and dynamics of the boundary

# Very brief (and incomplete) history

- 1983, Laughlin
  - ▶ Laughlin's wave function and plasma analogy
- 1986, Girvin, MacDonald, Platzman
  - ▶ Magnetoplasmon, structure factor, single mode approximation
- 1989, Read; Zhang, Hansson, Kivelson; 1990, Stone; 1991, Lee and Zhang
  - ▶ Chern-Simons-Ginzburg-Landau theory of FQHE
  - ▶ Hydrodynamics of FQHE and FQHE constraint
- 1996, Simon, Stern, Halperin
  - ▶ Magnetization current
- 1992, Wen, Zee; 1995, Avron, Seiler, Zograf; 2007, Tokatly, Vignale; 2009, Read; Haldane
  - ▶ Hall viscosity
- 2006, Zabrodin, Wiegmann; 2005-9, Bettelheim, Wiegmann, AA
  - ▶ 2D Dyson gas
  - ▶ 1D hydrodynamics of Calogero-Sutherland model

# Density, current, velocity

Microscopic fields:

$$m = e = c = 1$$

$$\rho(\mathbf{r}) = \sum_{\alpha} \delta(\mathbf{r} - \mathbf{r}_{\alpha}) \quad (2)$$

$$\mathbf{j}(\mathbf{r}) = \sum_{\alpha} (\mathbf{p}_{\alpha} - \mathbf{A}_{\alpha}) \delta(\mathbf{r} - \mathbf{r}_{\alpha}) \quad (3)$$

Poisson brackets (commutators), Landau, 1941

$$\{\rho, v'\} = \nabla \delta(\mathbf{r} - \mathbf{r}') \quad (4)$$

$$\{v_i, v'_j\} = \epsilon_{ij} \frac{\nabla \times \mathbf{v} + B}{\rho} \delta(\mathbf{r} - \mathbf{r}') \quad (5)$$

Here the velocity is defined as  $\mathbf{j} = \rho \mathbf{v}$ .

# Hamiltonian dynamics of ideal fluid

Hamiltonian:

$$H = \int d^2r \frac{\rho v^2}{2} + U[\rho]$$

$$U[\rho] = \int d^2r [\rho \epsilon(\rho) + A(\nabla \rho)^2]$$

- the Boussinesq approximation.

Hamiltonian + Poisson brackets  $\rightarrow$  Equations of motion

$$\dot{\rho} + \nabla(\rho \mathbf{v}) = 0, \quad \text{continuity}$$

$$\dot{\mathbf{v}} + (\mathbf{v} \nabla) \mathbf{v} + \nabla w = \mathbf{E} + \mathbf{v} \times \mathbf{B}. \quad \text{Euler}$$

where  $w = \delta U / \delta \rho$  - chemical potential

Dynamics is not vorticity free.

# Vorticity and FQHE constraint

$\omega = \nabla \times \mathbf{v}$  – vorticity

$$\partial_t(\omega + B) + \nabla \cdot ((\omega + B)\mathbf{v}) = 0$$

$$\partial_t \rho + \nabla \cdot (\rho \mathbf{v}) = 0$$

Constraint  $\omega + B = \zeta \rho$  is respected by equations of motion.

$$\nabla \times \mathbf{v} = \zeta(\rho - \rho_0)$$

FQHE constraint; [Stone, 1990](#); incompressibility! dimension:  $[\zeta] = \left[\frac{\hbar}{m}\right]$

– might emerge in [quantum physics](#)!  $\zeta = \frac{\hbar}{m} \frac{2\pi}{\nu}$ .

# Stress tensor $T_{ik}$

Dynamics of conserved quantities

$$\begin{aligned} \partial_t \rho + \partial_k j_k &= 0, & j_k &= \rho v_k, \\ \partial_t j_i + \partial_k T_{ik} &= \rho F_i, & T_{ik} &= P(\rho) \delta_{ik} + \rho v_i v_k, \\ & & F_i &= E_i + \epsilon_{ik} v_k B. \end{aligned}$$

Linear expansion around  $\rho = \rho_0$ ,  $\mathbf{v} = 0$

$$\delta T_{ik} = -K u_{nn} \delta_{ik}, \quad K = \frac{\partial P}{\partial \rho} - \text{bulk modulus.}$$

**Isotropic fluid: no shear**  $\delta T_{ik} \sim u_{ik}$ .

$$u_{ik} = \frac{1}{2} (\partial_i u_k + \partial_k u_i)$$

**Ideal fluid: no viscosity**  $\sim v_{ik}$

$$v_{ik} = \frac{1}{2} (\partial_i v_k + \partial_k v_i)$$

**Hall viscosity**  $\sim (\epsilon_{in} v_{ik} + \epsilon_{kn} v_{ni})$

strain and strain rate

# Hall viscosity (Lorentz shear modulus)

Isotropic **non-dissipative** fluid

“Viscous” stress tensor (known in plasma physics)

$$\delta T_{ik} = \Lambda(\epsilon_{in}v_{nk} + \epsilon_{kn}v_{ni}).$$

**Time-reversal invariance is broken!**


For Laughlin's states

$$\Lambda = \frac{\hbar\rho_0}{4}\nu^{-1} = \frac{\hbar}{8\pi l_B^2} = \frac{1}{8\pi} \frac{eB}{c} \quad - \text{Hall viscosity}$$

Generalization (**Read, 2009**)

$$\Lambda = \frac{\hbar\rho_0}{2} \left( \frac{\nu^{-1}}{2} + h_\psi \right)$$

# Hall viscosity



Stress forces are orthogonal to the motion  $\rightarrow$  **no dissipation!**


# Hall viscosity from adiabatic metric deformation

Hall viscosity  $\rightarrow$  Berry's phase with respect to variations of the metric

$$ds^2 = \frac{1}{\tau_2} |dx + \tau dy|^2.$$

Avron, Seiler, Zograf, 1995

Read, 2009



$\Psi_\beta$  - Laughlin's function on torus  $\tau = \tau_1 + i\tau_2$ .

$\Lambda$  - adiabatic curvature!

$$\Lambda = \frac{2\hbar}{L^2} \operatorname{Im} \left\langle \frac{\partial \Psi_\beta}{\partial \tau_1} \middle| \frac{\partial \Psi_\beta}{\partial \tau_2} \right\rangle$$

# Hall viscosity and Hall conductivity

**Hall conductivity** – non-dissipative part of conductivity tensor  $\sigma_{ik}$

**Hall viscosity** – non-dissipative part of viscosity tensor  $\eta_{ikmn}$

(Odd viscosity)

(Lorentz shear modulus)

$$\Psi_{\beta}(\tau_1, \tau_2; \Phi_1, \Phi_2) = \frac{1}{\mathcal{N}} \prod_{i < j} \theta_1(z_{ij}/L_x | \tau) e^{-\frac{1}{2} \sum_j (\text{Im } z_j)^2}$$

Laughlin's function on torus  $\tau = \tau_1 + i\tau_2$  with fluxes  $\Phi_{1,2}$  through torus' cycles. ( $z = x - \Phi_2 + \tau(y + \Phi_1)$ )

$\sigma_{xy}$ ,  $\Lambda$  – adiabatic curvatures!

$$\sigma_{xy} \sim \text{Im} \left\langle \frac{\partial \Psi_{\beta}}{\partial \Phi_1} \middle| \frac{\partial \Psi_{\beta}}{\partial \Phi_2} \right\rangle \quad \Lambda \sim \text{Im} \left\langle \frac{\partial \Psi_{\beta}}{\partial \tau_1} \middle| \frac{\partial \Psi_{\beta}}{\partial \tau_2} \right\rangle$$

Avron, Seiler, Zograf, 1995

Read, 2009

# Hall viscosity (Hamiltonian)

What is the Hamiltonian generating dynamics with Hall viscosity?

$$H = \int d^2r \rho \left[ \frac{1}{2} \mathbf{v}^2 + \eta \nabla \times \mathbf{v} \right] + U[\rho].$$

Hall viscosity  $\Lambda = \eta \rho_0$ .

$$H = \int d^2r \rho \frac{1}{2} \left( \mathbf{v} + \eta \nabla^* \log \rho \right)^2 + U'[\rho].$$

Here  $\nabla^* = -\hat{z} \times \nabla$ .

$\eta$ -term can be absorbed into  $\mathbf{v}$  but Poisson's brackets will change.

$$\int d^2x (\mathbf{r} \times \rho \mathbf{v} + \mu \rho) = \int d^2x \mathbf{r} \times \left( \rho \mathbf{v} + \frac{\mu}{2} \nabla^* \rho \right)$$

# Hall viscosity (Hamiltonian)

What is the Hamiltonian generating dynamics with Hall viscosity?

$$H = \int d^2r \rho \left[ \frac{1}{2} \mathbf{v}^2 + \eta \nabla \times \mathbf{v} \right] + U[\rho].$$

Hall viscosity  $\Lambda = \eta \rho_0$ .

$$H = \int d^2r \rho \frac{1}{2} \left( \mathbf{v} + \eta \nabla^* \log \rho \right)^2 + U'[\rho].$$

Here  $\nabla^* = -\hat{z} \times \nabla$ .

$\eta$ -term can be absorbed into  $\mathbf{v}$  but Poisson's brackets will change.

$$\int d^2x (\mathbf{r} \times \rho \mathbf{v} + \mu \rho) = \int d^2x \mathbf{r} \times \left( \rho \mathbf{v} + \frac{\mu}{2} \nabla^* \rho \right)$$

# Hall viscosity (Hamiltonian)

What is the Hamiltonian generating dynamics with Hall viscosity?

$$H = \int d^2r \rho \left[ \frac{1}{2} \mathbf{v}^2 + \eta \nabla \times \mathbf{v} \right] + U[\rho].$$

Hall viscosity  $\Lambda = \eta \rho_0$ .

$$H = \int d^2r \rho \frac{1}{2} \left( \mathbf{v} + \eta \nabla^* \log \rho \right)^2 + U'[\rho].$$

Here  $\nabla^* = -\hat{z} \times \nabla$ .

$\eta$ -term can be absorbed into  $\mathbf{v}$  but Poisson's brackets will change.

$$\int d^2x (\mathbf{r} \times \rho \mathbf{v} + \mu \rho) = \int d^2x \mathbf{r} \times \left( \rho \mathbf{v} + \frac{\mu}{2} \nabla^* \rho \right)$$

## FQHE constraint + Hall viscosity

$$H = \int d^2r \frac{\rho(\mathbf{v} + \eta \nabla^* \log \rho)^2}{2} + U[\rho] \quad a_i^* \equiv \epsilon^{ij} a_j$$

with  $(\nabla^* W = -\mathbf{A})$

$$\rho = \frac{1}{2\pi\beta} \Delta\phi, \quad \mathbf{v} = \nabla\pi - \nabla^*(\phi - W), \quad \{\pi, \rho'\} = \delta(\mathbf{r} - \mathbf{r}')$$

gives equations of motion with Hall viscosity

$$\Lambda = \eta\rho_0.$$

FQHE constraint is automatically satisfied

$$\nabla \times \mathbf{v} = 2\pi\beta(\rho - \rho_0).$$

# Classical Hydrodynamics I

## Classical Hamiltonian

$$H_{cl} = \int d^2z \rho \frac{1}{2m} \left( \nabla \pi - \nabla^* \left( \phi - W + \frac{\beta - 2}{4} \ln \rho \right) \right)^2$$

produces correct static structure factor  $s(k) = \frac{1}{2}k^2 \left( 1 + \frac{\beta - 2}{4} k^2 \right)$   
and FQHE constraint but **wrong** Hall viscosity.

**Quantum fluctuations** should be incorporated into classical (effective) Hamiltonian!

Quantum zero motion  $\rightarrow$  Landau diamagnetism!

# Classical Hydrodynamics II

diamagnetic current  $\mathbf{j}_B = -\frac{1}{2}\nabla^*\rho$

$$H_{cl} = \int d^2z \rho \left[ \frac{1}{2} \left( \mathbf{v} - \boxed{\frac{\beta - 2}{4}} \nabla^* \ln \rho \right)^2 + \boxed{\frac{1}{2}} \hbar \omega_B \right]$$

$$\rho = \frac{1}{2\pi\beta} \Delta \phi \quad \{\pi, \rho'\} = \delta(\mathbf{r} - \mathbf{r}'),$$

$$\mathbf{v} \equiv \nabla \pi - \nabla^*(\phi - W) \rightarrow \text{FQHE constraint: } \nabla \times \mathbf{v} = 2\pi\beta(\rho - \rho_0)$$

Static structure factor + Hall viscosity

$$s(k) = \frac{1}{2}k^2 \left( 1 + \boxed{\frac{\beta - 2}{4}} k^2 \right) \quad \Lambda = \left( \boxed{\frac{\beta - 2}{4}} + \boxed{\frac{1}{2}} \right) \hbar \rho_0 = \frac{\beta}{4} \hbar \rho_0$$

# Classical Hydrodynamics II

diamagnetic current  $\mathbf{j}_B = -\frac{1}{2}\nabla^*\rho$

$$H_{cl} = \int d^2z \rho \left[ \frac{1}{2} \left( \mathbf{v} - \boxed{\frac{\beta - 2}{4}} \nabla^* \ln \rho \right)^2 + \boxed{\frac{1}{2}} \hbar \omega_B \right]$$

$$\rho = \frac{1}{2\pi\beta} \Delta \phi$$

$$\{\pi, \rho'\} = \delta(\mathbf{r} - \mathbf{r}'),$$

$$\mathbf{v} \equiv \nabla \pi - \nabla^*(\phi - W) \rightarrow \text{FQHE constraint: } \nabla \times \mathbf{v} = 2\pi\beta(\rho - \rho_0)$$

Static structure factor

+

Hall viscosity

$$s(k) = \frac{1}{2}k^2 \left( 1 + \boxed{\frac{\beta - 2}{4}} k^2 \right)$$

$$\Lambda = \left( \boxed{\frac{\beta - 2}{4}} + \boxed{\frac{1}{2}} \right) \hbar \rho_0 = \frac{\beta}{4} \hbar \rho_0$$

# Hydrodynamic Lagrangian

$$L_{cl} = - \int d^2x \rho \left[ \dot{\pi} + \frac{1}{2} \mathbf{v}_\alpha^2 + \frac{1}{2} (\nabla \times \mathbf{A}) \right] \quad \text{Main Result}$$

$$\mathbf{v}_\alpha \equiv \nabla \pi + \mathbf{A} - \nabla^* (\phi + \alpha \ln \rho).$$

$$\alpha = \frac{\beta - 2}{4}, \quad \eta = \frac{\beta}{4}.$$

True e/m current:  $\mathbf{j} = \rho \mathbf{v}_\eta$

FQHE constraint:  $\nabla \times \mathbf{v}_\alpha = 2\pi\beta(\rho - \rho_0) + \alpha\Delta \ln \rho.$

# Linear response

- Static structure factor  $s(k) = \frac{1}{2}k^2 \left(1 + \frac{\beta-2}{4}k^2\right)$   
 1986 Girvin, MacDonald, Platzman
- Hall viscosity  $\Lambda = \frac{\beta}{4} \hbar \rho_0$   
 1995 Avron, Seiler, Zograf;  
 2007 Tokatly, Vignale;  
 2009 Read
- Electromagnetic response, e.g.,  $\sigma_H = \frac{1}{2\pi\beta} \left(1 + \frac{\beta-4}{4}k^2\right)$   
 2011 Hoyos, Son
- other E&M and dynamic response  
 2006 Tokatly

# Linear response

Change of density under small variations of  $\mathbf{E}$  and  $B$

$$\frac{\delta\rho}{\rho_0} = \frac{\omega_0^2}{\omega^2 - \Omega_k^2} \left[ \frac{e}{m\omega_0^2} (\nabla \mathbf{E}) - \left( 1 - \frac{\eta k^2}{m\omega_0} \right) \frac{\delta\omega_c}{\omega_0} \right]$$

with magnetoplasmon dispersion

$$\frac{\Omega_k^2}{\omega_0^2} = 1 - \frac{\beta - 2}{2} \frac{k^2}{m\omega_0} + \dots$$

# Chern-Simons-Ginzburg-Landau theory

1989, Read; Zhang, Hansson and Kivelson

$$L = \Phi^* \left( i\partial_t + a_0 + A_0 - \frac{1}{2m^*} (-i\nabla - \mathbf{a} - \mathbf{A})^2 \right) \Phi + \frac{1}{4\pi\beta} \epsilon^{\mu\nu\lambda} a_\mu \partial_\nu a_\lambda + V(|\Phi|^2)$$

Hydrodynamic form:

Change  $\Phi = \sqrt{\rho} e^{i\theta}$  and solve for constraints

$$\mathbf{a} = \nabla(\pi - \theta) + \nabla^* \phi, \quad \Delta\phi = 2\pi\beta\rho$$

$$L = -\rho \left[ \dot{\pi} + \frac{1}{2} \left( \nabla\pi - \nabla^* \left( \phi - W - \frac{1}{2} \ln\rho \right) \right)^2 + A_0 + \epsilon(\rho) \right]$$

UV regularization is missing  $(-\frac{1}{2} \rightarrow \frac{\beta-2}{4})$ .

Extra term is needed:  $\sim \frac{\beta}{4} \nabla \times (\mathbf{a} + \mathbf{A}) |\Phi|^2$ .

# Quantum Hydrodynamics

Ground state: Laughlin's wave function for filling fraction  $\nu = \frac{1}{\beta}$ :

$$\Psi_\beta = \prod_{j < k} (z_j - z_k)^\beta e^{-\sum_j W(z_j, \bar{z}_j)}, \quad W = \frac{1}{4}|z|^2 + \boxed{W_1(z)}$$

written as  $\Psi_\beta[\rho] = e^{-\frac{1}{2}E_\beta[\rho]}$

(Dyson's argument, 2006 Zabrodin, Wiegmann)

$$E_\beta[\rho] = -\beta \int d^2z d^2z' \rho(z) \ln |z - z'| \rho(z') + 2 \int d^2z \rho W + \frac{2-\beta}{2} \int d^2z \rho \ln \rho.$$

$$H = \int d^2x \frac{1}{2} \bar{V} \rho V \quad \text{acts on } \Psi[\rho]$$

where

$$V = \bar{\partial} \left[ \pi + i \left( \phi - W + \frac{\beta - 2}{4} \ln \rho \right) \right], \quad \pi = -i \frac{\delta}{\delta \rho}$$

# Dyson's argument

Laughlin's plasma in collective variables

$$\sum_{j \neq k} \ln |z_j - z_k| \rightarrow \int d^2 z d^2 z' \rho(z) \ln |z - z'| \rho(z') - \int d^2 z \rho \ln \frac{1}{\sqrt{\rho}}$$

$$\prod_j d^2 z_j \rightarrow [D\rho] \prod_j \frac{1}{\rho(z_j)} \rightarrow [D\rho] \exp \left[ - \int d^2 z \rho \ln \rho \right]$$

Partition function for plasma  $\|\Psi_\beta\|^2 = \int [D\rho] e^{-E_\beta[\rho]}$  with

2d Coulomb plasma

$$E_\beta[\rho] = -\beta \int d^2 z d^2 z' \rho(z) \ln |z - z'| \rho(z')$$

$$+ \underbrace{2 \int d^2 z \rho W}_{\text{background charge}} + \frac{2-\beta}{2} \int d^2 z \rho \ln \rho. \quad \text{UV-cutoff + entropy}$$

# Equilibrium density

Electrostatic energy

$$E_\beta[\rho] = -\beta \int d^2z d^2z' \rho(z) \ln|z - z'| \rho(z') + 2 \int d^2z \rho W + \frac{2-\beta}{2} \int d^2z \rho \ln \rho.$$

Electrostatic potential

$$\phi(z) = \beta \int d^2z' \ln|z - z'| \rho(z') \quad \Delta\phi = 2\pi\beta \rho$$

Equilibrium

$$\frac{\delta E_\beta}{\delta \rho} = 0 \quad \phi = W + \frac{2-\beta}{4} \ln \rho \quad \Delta W = B = 2\pi\beta \rho_0$$

$$\rho = \rho_0 + \frac{2-\beta}{8\pi\beta} \Delta \ln \rho$$

$$\rho = \rho_0 = \frac{B}{2\pi\beta} = \frac{1}{\beta} \frac{1}{2\pi l_0^2}$$

$$\nu = 1/\beta - \text{filling fraction}$$

# QFT wave function

$$\Psi_\beta[\rho] = e^{-\frac{1}{2}E_\beta[\rho]}$$

$$\|\Psi_\beta\|^2 = \int [D\rho] e^{-E_\beta[\rho]}$$

$$\left[ \pi + i \left( \phi - W - \frac{2-\beta}{4} \ln \rho \right) \right] \Psi_\beta = 0 \quad - \text{identity for Laughlin's } \Psi_\beta$$

$$\pi = -i \frac{\delta}{\delta \rho} \quad [\pi(x), \phi(x')] = -i \delta^{(2)}(x - x')$$

**Chiral constraint** (projection to LLL) valid for any “holomorphic” wave function

$$\bar{\partial} \left[ \pi + i \left( \phi - W - \frac{2-\beta}{4} \ln \rho \right) \right] = 0$$


$$\nabla \pi - \nabla^* \left( \phi - W - \frac{2-\beta}{4} \ln \rho \right) = 0$$

# Boundary dynamics

(dispersionless case)

$$\bar{\partial} \left[ \pi + i \left( \phi - W - \frac{2 - \beta}{4} \ln \rho \right) \right] = 0 \quad - \text{chiral constraint}$$

$$\pi + i \left( \phi - W - \frac{2 - \beta}{4} \ln \rho \right) = V(z, t) \quad - \text{analytic}$$



$V(z, t) \rightarrow h(x, t)$  – boundary

$H = \int d^2r \rho A_0$  – dynamics

$h_t + A'_0 h_x + A''_0 h h_x = 0$  – incompressible droplet Iso, Rey, 1995

Hall viscosity  $\rightarrow$  boundary profile  $\rightarrow$  **dispersive corrections!**

# 1D Calogero-Sutherland model

Calogero model in harmonic potential

$$H = \frac{1}{2} \sum_j (p_j^2 + x_j^2) + \frac{1}{2} \sum_{j < k} \frac{\lambda(\lambda - 1)}{(x_j - x_k)^2}$$

The ground state wave function

$$\Psi_0 = \prod_{j < k} (x_j - x_k)^\lambda e^{-\frac{1}{2} \sum_j x_j^2}$$

Collective field theory

$$H = \int dx \left( \frac{\rho v^2}{2} + \rho \epsilon(\rho) \right)$$

$$\epsilon(\rho) = \frac{1}{2} \left[ \pi \lambda \rho^H - (\lambda - 1) \partial_x \ln \sqrt{\rho} + x \right]^2$$

# Hydrodynamics of 1D Calogero-Sutherland model

Hydrodynamics for Calogero model in harmonic potential

$$H = \int dx \rho \frac{1}{2} |\partial_x \Phi|^2$$

$$\Phi = \pi - i \left( \phi - W + \frac{\lambda - 1}{2} \ln \rho \right)$$

Here  $\phi = \int dx' \log |x - x'| \rho(x')$ ,  $W = x^2/2$  and  $\lambda$  is Calogero coupling constant ( $\lambda = 1$  for free fermions).

FQHE

$$\Phi = \pi - i \left( \phi - W + \frac{\beta - 2}{4} \ln \rho \right)$$

# Conclusions

- 1 Hamiltonian formulation for FQHE hydrodynamics is constructed
  - ▶ FQHE constraint
  - ▶ Hall viscosity
  - ▶ Linear response
  - ▶ Correspondence to Chern-Simons-Ginzburg-Landau
  - ▶ Connection to Laughlin's function and quantum hydro
- 2 Chiral constraint and boundary dynamics of FQHE droplet
- 3 Analogies with Calogero-Sutherland model

# Conclusions

- 1 Hamiltonian formulation for FQHE hydrodynamics is constructed
  - ▶ FQHE constraint
  - ▶ Hall viscosity
  - ▶ Linear response
  - ▶ Correspondence to Chern-Simons-Ginzburg-Landau
  - ▶ Connection to Laughlin's function and quantum hydro
- 2 Chiral constraint and boundary dynamics of FQHE droplet
- 3 Analogies with Calogero-Sutherland model

# Conclusions

- 1 Hamiltonian formulation for FQHE hydrodynamics is constructed
  - ▶ FQHE constraint
  - ▶ Hall viscosity
  - ▶ Linear response
  - ▶ Correspondence to Chern-Simons-Ginzburg-Landau
  - ▶ Connection to Laughlin's function and quantum hydro
- 2 Chiral constraint and boundary dynamics of FQHE droplet
- 3 Analogies with Calogero-Sutherland model

# Recent Related Works

- Paul Wiegmann
  - ▶ arXiv:1305.6893, *Anomalous Hydrodynamics of Fractional Quantum Hall States*
  - ▶ arXiv:1211.5132, *Quantum Hydrodynamics of Fractional Hall Effect: Quantum Kirchhoff Equations*
  - ▶ Phys. Rev. Lett. 108, 206810 (2012), *Non-Linear hydrodynamics and Fractionally Quantized Solitons at Fractional Quantum Hall Edge*
- Dam Son
  - ▶ arXiv:1306.0638, *Newton-Cartan Geometry and the Quantum Hall Effect*
- Eldad Bettelheim
  - ▶ arXiv:1306.3782, *Integrable Quantum Hydrodynamics in Two Dimensional Phase Space*