

Before we start: Important setup of your Computer

```
change directory:  
cd /afs/ictp/public/shared/smr2475  
.setup-config.sh  
logout  
login again
```

1st Tutorial:

The Basics of DFT

Lydia Nemec and Oliver T. Hofmann

Fritz Haber Institute of the Max Planck Society

7th August 2013



MAX-PLANCK-GESELLSCHAFT

Density functional theory and beyond:
Computational materials science for real materials, 2013

The ultimate goal!

$$H\Psi = E\Psi$$

Second order differential equation for a $3N_e$ -variable function Ψ
⇒ **Complex problem**

Unsolved issues at the simplest level of approximations (multiple solutions, ...)

Goals of this tutorial

- Familiarise with practical aspects of electronic structure theory in general and density functional theory (DFT) in particular
- Hartree-Fock (HF) method and Kohn-Sham DFT (non-periodic)
- Numerical solution of the approximate equations (tool: FHI-aims)
- Exploring potential energy surfaces (total energies at fixed nuclei, local minima, vibrational spectra)
- Electronic structure analysis (visualisation tools, electron density, Kohn-Sham orbitals and spectrum)

Solving the Kohn-Sham equations

Hohenberg-Kohn Theorem $\Psi(\mathbf{r}_1 \dots \mathbf{r}_{N_e}) \Leftrightarrow n(\mathbf{r})$

Kohn-Sham scheme

$$\left(-\frac{1}{2} \nabla^2 + \int d^3 r' \frac{n(\mathbf{r}')}{|\mathbf{r} - \mathbf{r}'|} + v_{xc} + v \right) \psi_i = \epsilon_i \psi_i$$
$$\Rightarrow n = \sum_i f_i |\psi_i|^2$$

KS Orbitals $\{\psi_i\}$ $\langle \psi_i, \psi_j \rangle = \delta_{ij}$

XC Potential v_{xc} unknown, but many approximations exist

LDA, PBE, ... \rightarrow control.in

External potential contains ionic contributions \rightarrow geometry.in

Basis set

Expand in a finite basis $\{\phi_i\}$: $\psi_j = \sum_{i=1}^N c_{ij} \phi_i$

Finite Basis

Numeric atom centered

Gaussians

Plane waves + pseudoisation

Slater type


Grid based

... many more

Basis set

Expand in a finite basis $\{\phi_i\}$: $\psi_j = \sum_{i=1}^N c_{ij} \phi_i$

Numeric atom centered (FHI-aims)



Courtesy V. Blum

Basis set

Expand in a finite basis $\{\phi_i\}$: $\psi_j = \sum_{i=1}^N c_{ij} \phi_i$



Generalized matrix eigenvalue equation in c_{ij}

$$\hat{h}^{KS} \psi = E \psi \quad \Rightarrow \quad \sum_j h_{ij}(c) c_{jl} = \epsilon_l \sum_j s_{ij} c_{jl}$$


Overlap matrix $s_{ij} = \langle \phi_i, \phi_j \rangle$

Hamilton matrix $h_{ij} = \langle \phi_i, \hat{h}^{KS} \phi_j \rangle$



Self-consistent solution

Finding the self-consistent solution



Mixing (Pulay)

(Sca)Lapack
ELPA

Criteria

- Energy
- Charge density
- Sum of eigenvalues
- Force

Problem I: The hydrogen atom

Tasks:

- Input files needed to run FHI-aims.
- Test the convergence of the total energy with basis size.
- Compare the total energy of the hydrogen atom computed with different methods implemented in FHI-aims. Do all methods converge to the same result?

Basic electronic structure with FHI-aims

FHI-aims: input files

geometry.in

control.in

Basic electronic structure with FHI-aims

FHI-aims: input files

geometry.in**control.in****# Atomic structure**

x y z

atom 0.0 0.0 0.0 H

atom 1.0 0.0 0.0 H

initial_moment 1.0

That's a comment

Units:**Positions in Å****Energies in eV****Manual, chap. 2.1**

Basic electronic structure with FHI-aims

FHI-aims: input files

geometry.in**# Atomic structure**

```
#      x     y     z
atom 0.0 0.0 0.0 H
atom 1.0 0.0 0.0 H
initial_moment 1.0

# That's a comment
```

control.in**# Physical model settings**

```
xc pw-lda
charge 0.
spin collinear
```

Units:**Positions in Å****Energies in eV****Manual, chap. 2.1**

Basic electronic structure with FHI-aims

FHI-aims: input files

geometry.in

Atomic structure

```
#      x     y     z
atom 0.0 0.0 0.0 H
atom 1.0 0.0 0.0 H
initial_moment 1.0
# That's a comment
```

control.in

Physical model settings

```
xc pw-lda
charge 0.
spin collinear
```

SCF convergence settings

```
sc_accuracy_eev 1E-2
sc_accuracy_etot 1E-5
sc_accuracy_rho 1E-4
sc_iter_limit 100
```

Units:

Positions in Å

Energies in eV

Manual, chap. 2.1

Basic electronic structure with FHI-aims

FHI-aims: input files

geometry.in

Atomic structure

x y z

atom 0.0 0.0 0.0 H

atom 1.0 0.0 0.0 H

initial_moment 1.0

That's a comment

Units:

Positions in Å

Energies in eV

control.in

Physical model settings

xc pw-lda

charge 0.

spin collinear

SCF convergence settings

sc_accuracy_eev 1E-2

sc_accuracy_etot 1E-5

sc_accuracy_rho 1E-4

sc_iter_limit 100

Species specifics

...

Basic electronic structure with FHI-aims

species_default

```
/usr/local/aimsfiles/species_default
```

Predefined species

Manual, chap. 2.2

Copy-paste into control.in

- light
- tight
- really tight

Basic electronic structure with FHI-aims

species_default

```
/usr/local/aimsfiles/species_default
```

Predefined species

Manual, chap. 2.2

Copy-paste into control.in

- **light** **Increased accuracy:**
Basis
Hartree potential
- **tight** **Basis cutoff potential**
Integration grids
- **really tight**

Basic electronic structure with FHI-aims

species_default

```
/usr/local/aimsfiles/species_default
```

Predefined species

Copy-paste into control.in

Manual, chap. 2.2

- light
- tight
- really tight



- Fast, many production tasks**
Fast pre-relaxation
- Used to verify important results**
Converged settings
- Heavily converged numerical settings**
Explicit convergence tests

species_default

```
/usr/local/aimsfiles/species_default
```

Predefined species

Manual, chap. 2.2

Copy-paste into control.in

- light
- tight
- really tight



- Fast, many production tasks**
Fast pre-relaxation
- Used to verify important results**
Converged settings
- Heavily converged numerical settings**
Explicit convergence tests

Additionally converge basis ("tiers")!

Basic electronic structure with FHI-aims

FHI-aims output

1

Invoking FHI-aims ...

Introduction

Basic electronic structure with FHI-aims

FHI-aims output

1

Invoking FHI-aims ...

2

Reading file control.in.

Summary of control.in file

Basic electronic structure with FHI-aims

FHI-aims output

1

Invoking FHI-aims ...

2

Reading file control.in.

3

Reading geometry description geometry.in.

Summary of geometry.in file

FHI-aims output

1

Invoking FHI-aims ...

2

Reading file control.in.

3

Reading geometry description geometry.in.

4

Preparing all fixed parts of the calculation.

Geometry independent preparations

Basis set generation

Basic electronic structure with FHI-aims

FHI-aims output

5

```
-----  
Begin self-consistency loop: Initialization.  
Date : 20130610, Time : 162002.389  
-----
```

Geometry dependent preparations
Integration grid
Initialization of charge density

Basic electronic structure with FHI-aims

FHI-aims output

5

```
-----  
Begin self-consistency loop: Initialization.  
Date : 20130610, Time : 162002.389  
-----
```

6

```
-----  
Begin self-consistency iteration # 1  
Date : 20130610, Time : 162002.445  
-----
```

First SCF cycle

Basic electronic structure with FHI-aims

FHI-aims output

5

```
-----  
Begin self-consistency loop: Initialization.  
Date : 20130610, Time : 162002.389  
-----
```

6

```
-----  
Begin self-consistency iteration # 1  
Date : 20130610, Time : 162002.445  
-----
```

First SCF cycle

THIS
TUTORIAL

» Energy

```
| Total energy : -13.01991124 eV  
| Total energy, T → 0 : -13.01991124 eV  
| Electronic free energy : -13.01991124 eV
```



Periodic metals only

Basic electronic structure with FHI-aims

FHI-aims output

5

```
-----  
Begin self-consistency loop: Initialization.  
Date : 20130610, Time : 162002.389  
-----
```

6

```
-----  
Begin self-consistency iteration # 1  
Date : 20130610, Time : 162002.445  
-----
```

First SCF cycle

» Self-consistency convergence accuracy

Change of charge density : 0.6753E-02
Change of sum of eigenvalues : 0.4376E+00 eV
Change of total energy : 0.1143E-01 eV

Basic electronic structure with FHI-aims

FHI-aims output

5

```
-----  
Begin self-consistency loop: Initialization.  
Date : 20130610, Time : 162002.389  
-----
```

6

```
-----  
Begin self-consistency iteration # 6  
Date : 20130610, Time : 162002.560  
-----
```

Sixth SCF cycle

» Self-consistency convergence accuracy

- | Change of charge density : 0.3163E-05
- | Change of sum of eigenvalues : -.9415E-05 eV
- | Change of total energy : 0.2388E-10 eV

Basic electronic structure with FHI-aims

FHI-aims output

7

Self-consistency cycle converged.

Basic electronic structure with FHI-aims

FHI-aims output

7

Self-consistency cycle converged.

» Energy and forces



```
| Total energy uncorrected : -0.130198526094581E+02 eV
| Total energy corrected : -0.130198526094581E+02 eV
| Electronic free energy   : -0.130198526094581E+02 eV
```

» SCF info

```
| Number of self-consistency cycles : 6
```

» Timings

Basic electronic structure with FHI-aims

FHI-aims output

7

Self-consistency cycle converged.

» Energy and forces



```
| Total energy uncorrected : -0.130198526094581E+02 eV
| Total energy corrected : -0.130198526094581E+02 eV
| Electronic free energy   : -0.130198526094581E+02 eV
```

» SCF info

```
| Number of self-consistency cycles : 6
```

» Timings

8

Have a nice day.

FHI-aims output

7

Self-consistency cycle converged.

Postprocessing

Structure optimization

- » Get next relaxation step
- » Redo SCF for new geometry

8

Have a nice day.

Problem II: Hydrofluoric acid (HF)


One of the first papers which systematically investigated the performance of DFT was published by John A. Pople and coworker in 1993.

Tasks:

- Find the equilibrium bond distance of HF.
- Compare the HF bond length for different methods.
- Calculate the atomization energy (ΔH_{at}).
- Compute the dipole moment for different methods and bond lengths.

Basic electronic structure with FHI-aims


Forces



Basic electronic structure with FHI-aims


Forces

Energy gradient



Forces


**Structure optimization:
Find local minimum on potential energy surface (PES)**



Basic electronic structure with FHI-aims


Forces

**Structure optimization:
Find local minimum on potential energy surface (PES)**



Many methods !


Industry standard: quasi Newton methods



Basic electronic structure with FHI-aims

Forces

Structure optimization:
Find local minimum on potential energy surface (PES)




Local structure optimisation

Problem IV to VIII: Hydronium cation (H_3O^+)

Tasks:

- Relax structure with two different starting points.
- Make a vibrational analysis.
- Explore the limits of the harmonic approximation.



Local structure optimisation

Harmonic molecular motion

How do atoms move in a potential V ?

⇒ Solve equations of motion!

$$\frac{d}{dt} \frac{\partial T}{\partial \dot{R}_i} + \frac{\partial V}{\partial R_i} = 0$$

Kinetic energy: T

Potential energy: V

Local structure optimisation

Harmonic molecular motion

How do atoms move in a potential V ?

⇒ Solve equations of motion!

$$\frac{d}{dt} \frac{\partial T}{\partial \dot{R}_i} + \frac{\partial V}{\partial R_i} = 0$$

Kinetic energy: T

Potential energy: V

⇒ Taylor expansion of V

around equilibrium position R_0 + harmonic approximation

$$V = V_0 \underbrace{-F(R_0)R}_{=0 \text{ equilibrium}} + \frac{1}{2} R^T H(R_0) R \underbrace{+\dots}_{=0 \text{ harmonic approximation}} \text{higher terms}$$

F : Forces

H : Hessian

Harmonic molecular motion

How do atoms move in a potential V ?

⇒ **Solve equations of motion!**

$$\frac{d}{dt} \frac{\partial T}{\partial \dot{R}_i} + \frac{\partial V}{\partial R_i} = 0$$

Kinetic energy: T

Potential energy: V

⇒ **Solution**

The Dynamic Matrix D_{ij} : $D_{ij} = \frac{1}{\sqrt{M_i}\sqrt{M_j}} H_{ij}$

$$R \sim Q e^{i\omega t}, \quad \text{with} \quad DQ - \omega^2 Q = 0$$

Eigenmodes Q

- If (1) Harmonic approximation is valid
(2) Equilibrium geometry

Local structure optimisation

Vibrations

Solve $\mathbb{R}^{3N} \times \mathbb{R}^{3N}$ eigenvalue equation

$$\det(D - \omega^2 \mathbf{1}) = 0$$

Hessian H **Dynamic matrix D**

$$H^{ij} := \frac{\partial^2 E}{\partial R_i \partial R_j} \quad D_{ij} = \frac{1}{\sqrt{M_i} \sqrt{M_j}} H_{ij}$$

In practice: finite central numerical differences (of forces)

Wrapper

```
> aims_vibrations.mpi.pl
```

Manual, chap 4.6

Local structure optimisation

Vibrations

Solve $\mathbb{R}^{3N} \times \mathbb{R}^{3N}$ eigenvalue equation

$$\det(D - \omega^2 \mathbf{1}) = 0$$

Get

- » Eigenmodes $\{Q_i, i \in 1 \dots 3N\}$
- » Eigenfrequencies

Local structure optimisation

Vibrations

Solve $\mathbb{R}^{3N} \times \mathbb{R}^{3N}$ eigenvalue equation

$$\det(D - \omega^2 \mathbf{1}) = 0$$

Get

- » **Eigenmodes** $\{Q_i, i \in 1 \dots 3N\}$
- » **Eigenfrequencies**
- » **6 (almost) zero frequency modes** (if molecule non-linear)
translations + rotations
- » **Imaginary frequency** \Rightarrow Saddle point

Local structure optimisation

Vibrations

Solve $\mathbb{R}^{3N} \times \mathbb{R}^{3N}$ eigenvalue equation

$$\det(D - \omega^2 \mathbf{1}) = 0$$

Get

- » **Eigenmodes** $\{Q_i, i \in 1 \dots 3N\}$
- » **Eigenfrequencies**
- » **6 (almost) zero frequency modes** (if molecule non-linear)
translations + rotations
- » **Imaginary frequency** \Rightarrow Saddle point
- » **Infrared intensities** (derivative of dipole moment μ)

$$I_i \sim \left| \frac{d\mu}{dQ_i} \right|^2$$

Vibrations

Based on harmonic approximation !
Limitation: Problem IX
Beyond: Tutorial 3 (MD) on Friday

Get

- » Eigenmodes $\{Q_i, i \in 1 \dots 3N\}$
- » Eigenfrequencies
- » 6 (almost) zero frequency modes (if molecule non-linear)
translations + rotations
- » Imaginary frequency \Rightarrow Saddle point
- » Infrared intensities (derivative of dipole moment μ)

$$I_i \sim \left| \frac{d\mu}{dQ_i} \right|^2$$


Visualising density differences

Visualization

Orbitals and densities

Keyword in control.in

```
output cube eigenstate homo
cube filename HOMO(cube)
output cube total_density
cube filename tot_dens_uc(cube)
```



Get: ***.cube file** - values on a regular 3D grid.

Software: **molden** (jmol, gdis, xcrysden)
⇒ Appendix of handout

Practical issues

- **Each calculation one directory**

```
> mkdir tutorial1  
> cd tutorial1  
> mkdir HF
```

- **2 input files**

```
geometry.in  
control.in
```

- **Launching FHI-aims calculation**

```
mpirun -np 4 aims.hands-on-2013.scalapack.mpi.x  
| tee aims.out
```

- **... scripting helps !**

(Sample scripts in appendix of handout)

Before we start: Important setup of your Computer

change directory:

```
cd /afs/ictp/public/shared/smr2475  
.setup-config.sh
```

logout from your KDE session

login again

Access the info-lab machine from outside via ssh:

Access the ICTP gateway

```
ssh your_user_name@ssh.ictp.it
```

From the gateway you can reach your workstation:

```
ssh hp83-inf-XX
```

replace XX by the number of your maschine (XX=1,..51)

Acknowledgements

Special Thanks!



Viktor Atalla



Adriana Supady

Your Tutors for the afternoon




Oliver T. Hofmann



Lydia Nemec



Christian Carbogno



Franz Knuth



Arvid Ihrig



Markus Schneider



Volker Blum

