



The Abdus Salam
**International Centre
for Theoretical Physics**



**Conference on Frontiers of Nanoscience
24 August - 1 September 2015, Trieste, Italy**

Aharonov-Bohm Blockade in Magneto-Polaronic Tunneling of Strongly Correlated Electrons

Gleb SKOROBAGATKO

Dahlem Center for Complex Quantum Systems,
Free University of Berlin, Germany

Abstract:

The magnetopolaron generalization of a Majorana-resonant-level model [1] in the case of single-electron transistor with symmetric tunnel couplings between $g=1/2$ -Luttinger liquid leads [2,3] and quantum dot vibrating in the transverse external magnetic field) - leads to the unexpected (Aharonov-Bohm phase-driven) type of "magnetopolaron-assisted" resonant tunneling with strong interference between different virtual "vibronic" channels. As the result, in a sharp contrast with usual "polaronic" [2] (or usual magnetopolaronic [3]) blockade at zero-temperature for ultra-small frequencies of quantum vibrations - in the $g=1/2$ -magnetopolaronic SET a complete compensation of magnetopolaronic blockade is impossible even by means of very high bias voltage (or temperatures)[1]. This phenomenon can be called as a novel "Aharonov-Bohm" type of magnetopolaronic blockade or simply as the "Aharonov-Bohm blockade" being specific for the $g=1/2$ -magnetopolaronic SETs [1].



The Abdus Salam
**International Centre
for Theoretical Physics**



References:

- [1] G.Skorobagatko, arxiv 1310.1326,(2013).
- [2] G.Skorobagatko, Phys.Rev.B,85,075310,(2012).
- [3] A.Komnik, A.O.Gogolin, Phys.Rev.Lett.,90,246403,(2003).
- [4] S.Majer, A.Komnik, Phys.Rev.B, 82, 165116,(2010).
- [5] G.Skorobagatko, S.I.Kulinich, I.V.Krive, R.I.Shekhter, M.Jonson, Low Temp.Phys., 37, 1032, (2011).