



The Abdus Salam

**International Centre  
for Theoretical Physics**



**Conference on Frontiers of Nanoscience  
24 August - 1 September 2015, Trieste, Italy**

## **Dark Modes in Open Disordered Media: Analytical, Numerical, and Experimental Results**

A. Eisenbach, Z. Shi, R. Berkovits, Yu. Bliokh, **V. Freilikher**, A.  
Genack, M. Kaveh, F. Nori

Bar-Ilan University, Department of Physics  
Ramat Gan, Israel

### Abstract:

We explore numerically, analytically, and experimentally the relationship between quasi-normal modes (QNMs) and transmission resonant peaks (TR) in the transmission spectrum of one-dimensional (1D) and quasi-1D open disordered systems, both optical and electron. It is shown for the first time that for weak disorder there exist two types of the eigenstates: ordinary QNMs which are associated with a TR, and dark QNMs which do not exhibit peaks in transmission or within the sample. In addition, unlike ordinary QNMs, the lifetimes of dark modes goes down or increase anomalously slowly as disorder increases. Over a wide range of the strength of disorder, the ratio  $N_{\text{res}}/N_{\text{mod}}$  of the number of transmission peaks  $N_{\text{res}}$  to the number of QNMs  $N_{\text{mod}}$  is insensitive to the type and degree of disorder and is close to the value  $\sqrt{\frac{2}{5}}$  which we derive analytically in the weak-scattering approximation. Dark modes have been also discovered experimentally in quasi-1D open disordered samples. Our microwave measurements and modal analysis of transmission in the crossover to localization in quasi-1D systems give a ratio of  $N_{\text{res}}/N_{\text{mod}}$  close to  $\sqrt{\frac{2}{5}}$ . In diffusive quasi-1D samples, however,  $N_{\text{res}}/N_{\text{mod}}$  falls as the effective number of transmission eigenchannels,  $M$ , increases. Once  $N_{\text{mod}}$  is divided by  $M$ , the ratio  $N_{\text{res}}/N_{\text{mod}}$  is close to the ratio found in 1D.